

Permutation groups without irreducible elements

A. M. W. Glass and H. Dugald Macpherson

May 10, 2016

Dedicated to the memory of Rüdiger Göbel

Abstract

We call a non-identity element of a permutation group irreducible if it cannot be written as a product of non-identity elements of disjoint support. We show that it is indeed possible for a sublattice subgroup of $\text{Aut}(\mathbb{R}, \leq)$ to have no irreducible elements and still be transitive on the set of pairs $\alpha < \beta$ in \mathbb{R} . This answers a question raised in “The first-order theory of ℓ -permutation groups”, a Conference talk by the first author.

$G \upharpoonright H$

Let (Ω, \leq) be a totally ordered set and G be a subgroup of $\text{Aut}(\Omega, \leq)$. Let 1 be the identity element of $\text{Aut}(\Omega, \leq)$ and $g \in G \setminus \{1\}$. Then g is said to be *irreducible* if $g = g_1 g_2$ with $g_1, g_2 \in G$ and $\text{supp}(g_1) \cap \text{supp}(g_2) = \emptyset$ implies $g_1 = 1$ or $g_2 = 1$. Note that if $G = \text{Aut}(\Omega, \leq)$, then $g \in G$ is irreducible if and only if g has a single supporting interval; *i. e.*, there is $\sigma \in \text{supp}(g)$ such that the convexification in Ω of $\{\sigma g^n \mid n \in \mathbb{Z}\}$ is $\text{supp}(g)$. We prove:

Theorem A. *There is an ℓ -subgroup of $\text{Aut}(\mathbb{R}, \leq)$ that is transitive on ordered pairs $\alpha < \beta$ and has no irreducible elements.*

Here, an ℓ -subgroup of $\text{Aut}(\mathbb{R}, \leq)$ is a subgroup G of $\text{Aut}(\mathbb{R}, \leq)$ such that $g_+ \in G$ whenever $g \in G$, where $\alpha g_+ := \alpha g$ if $\alpha g \geq \alpha$ and $\alpha g_+ = \alpha$ if $\alpha g \leq \alpha$

2010 AMS Classification: 20B22, 06F15.

Keywords: order-preserving permutation, ℓ -permutation group.

($\alpha \in \mathbb{R}$). In particular, G is a lattice-ordered group where $f \vee g = (fg^{-1} \vee 1)g$ and $f \wedge g = (f^{-1} \vee g^{-1})^{-1}$. For background on ordered permutation groups and ℓ -groups see [1].

Proof of Theorem A. Let $g \in \text{Aut}(\mathbb{R}, \leq)$. We say that g has period $n \in \mathbb{Z}_+$ if $(\alpha + n)g = \alpha g + n$ for all $\alpha \in \mathbb{R}$. Let

$$G := \{g \in \text{Aut}(\mathbb{R}, \leq) \mid (\exists m \in \mathbb{Z}_+)(g \text{ has period } m)\}.$$

Then G is transitive on ordered pairs $\alpha < \beta$ in \mathbb{R} and it is easily checked that (G, \mathbb{R}) is an ℓ -permutation group. Obviously, if $f \in G$ fixes no point in \mathbb{R} , then it must be irreducible. So G has irreducible elements. On the other hand, if $g \in G$ has period m and is not the identity but fixes $\alpha_0 \in \mathbb{R}$ (and so fixes $\alpha_0 + km$ for all $k \in \mathbb{Z}$), define $g_1, g_2 \in G$, each with periods $2m$, as follows:

$$g_1(x) = \begin{cases} g(x) & \text{if } x \in [\alpha_0 + 2km, \alpha_0 + (2k+1)m), \quad k \in \mathbb{Z} \\ x & \text{if } x \in [\alpha_0 + (2k+1)m, \alpha_0 + (2k+2)m), \quad k \in \mathbb{Z} \end{cases}$$

$$g_2(x) = \begin{cases} g(x) & \text{if } x \in [\alpha_0 + (2k+1)m, \alpha_0 + (2k+2)m), \quad k \in \mathbb{Z} \\ x & \text{if } x \in [\alpha_0 + 2km, \alpha_0 + (2k+1)m), \quad k \in \mathbb{Z} \end{cases}.$$

Then g_1 and g_2 have disjoint supports and $g = g_1 g_2$, so g is reducible. Thus if $H := \{g \in G \mid 0g = 0\}$, then H has no irreducible elements. Now H acts faithfully on \mathbb{R}_+ and $(H \upharpoonright \mathbb{R}_+, \mathbb{R}_+)$ (the permutation group induced by H on \mathbb{R}_+) is an ℓ -permutation group that is transitive on ordered pairs $\alpha < \beta$ in \mathbb{R}_+ . Consequently we obtain an ℓ -permutation group (H^*, \mathbb{R}) that is transitive on pairs $\alpha < \beta$ in \mathbb{R} and has no irreducible elements. For let $\varphi : \mathbb{R} \rightarrow \mathbb{R}_+$ be an order-preserving bijection between \mathbb{R} and \mathbb{R}_+ and $h^* \in \text{Aut}(\mathbb{R}, \leq)$ be given by $\alpha h^* = (\alpha \varphi) h \varphi^{-1}$ ($\alpha \in \mathbb{R}, h \in H$). Then the desired properties transfer from H (acting on \mathbb{R}_+) to $H^* = \{h^* : h \in H\}$ (acting on \mathbb{R}). \square

The above proof can similarly be adapted to ℓ -permutation groups (L, \mathbb{Q}) that are transitive on pairs $\alpha < \beta$ in \mathbb{Q} and have no irreducible elements.

Acknowledgment. We are most grateful to Queens' College, Cambridge and the Engineering and Physical Sciences Research Council (grant EP/K020692/1) for funding to attend the Conference in memory of Rüdiger Göbel.

References

- [1] A. M. W. Glass, *Ordered Permutation Groups*, London Math. Soc. Lecture Notes Series **55**, Cambridge University Press, Cambridge, 1981.

A. M. W. Glass,
Queens' College, Cambridge CB3 9ET, U.K.

E-mail: amwg@dpmms.cam.ac.uk

H. Dugald Macpherson, School of Mathematics, University of Leeds, Leeds LS2
9JT, U. K.

E-mail: H.D.MacPherson@leeds.ac.uk