

Calculus with Precalculus

Vladimir V. Kisil

22nd May 2003

Chapter 1

General Information

This interactive manual is designed for students attending this course in Eastern Mediterranean University. The manual is being regularly updated in order to reflect material presented during lectures.

The manual is available at the moment in HTML with frames (for easier navigation), HTML without frames and PDF formats. Each from these formats has its own advantages. Please select one better suit your needs.

There is on-line information on the following courses:

- Calculus I.
- Calculus II.
- Geometry.

There are other on-line calculus manuals. See for example E-calculus of D.P. Story.

1.1 Warnings and Disclaimers

Before proceeding with this interactive manual we stress the following:

- These Web pages are designed in order to help students as a source of *additional information*. They are **NOT** an obligatory part of the course.
- The main material introduced during *lectures* and is contained in *Text-book*. This interactive manual is **NOT** a substitution for any part of those primary sources of information.

- It is **NOT** required to be familiar with these pages in order to pass the examination.
- The entire contents of these pages is continuously improved and updated. Even for material of lectures took place weeks or months ago changes are made.

1.2 Recommended Exercises

- Section 1.1. Problems 17, 23, 25
Section 1.2. Problems 17, 27
Section 1.3. Problems 25, 31, 39, 63
Section 1.4. Problems 11, 19, 23, 33
Section 1.5. Problems 15, 49, 55, 65
Section 2.1. Problems 1, 19
Section 2.2. Problems 21, 27
Section 2.3. Problems 21, 53, 55
Section 2.4. Problems 13, 17, 19, 41, 45
Section 2.5. Problems 17, 19, 47, 55
Section 2.6. Problems 11, 17, 27, 33
Section 2.7. Problem 7
Section 2.8. Problems 1, 5
Section 3.1. Problems 7, 25
Section 3.2. Problems 11, 15, 41
Section 3.3. Problems 5, 17, 29
Section 3.4. Problems 7, 13, 23
Section 3.5. Problems 13, 17, 31, 41
Section 3.6. Problems 5, 7, 29
Section 4.1. Problems 9, 17
Section 4.2. Problems 17, 21, 23, 31, 35, 39
Section 4.3. Problems 9, 27, 35
Section 4.4. Problems 9, 27, 33
Section 4.5. Problems 13, 21, 25
Section 4.6. Problems 19, 25, 35, 47, 59
Section 5.1. Problems 15, 23
Section 5.2. Problems 11, 13, 29
Section 5.3. Problems 7, 15, 23, 27, 29
Section 5.4. Problems 3, 14
Section 5.5. Problems 7, 12, 29, 33
Section 6.1. Problems 25, 33

- Section 6.2. Problems 37, 41
- Section 6.3. Problems 33, 37
- Section 6.4. Problems 11, 23, 31, 37
- Section 6.5. Problems 35, 43, 45
- Section 6.7. Problems 5, 21, 43, 53, 61
- Section 6.8. Problems 32, 34
- Section 6.9. Problems 9, 17, 63, 65
- Section 7.1. Problems 9, 17, 29, 31
- Section 7.2. Problems 3, 5, 13, 21
- Section 7.3. Problems 5, 9, 11, 17
- Section 7.4. Problems 7, 21, 25
- Section 7.5. Problems 5, 9, 15
- Section 7.7. Problems 13, 15, 53, 63
- Section 8.1. Problems 23, 31
- Section 8.2. Problems 13, 27, 47
- Section 8.3. Problems 5, 11, 15, 25, 27, 39, 43
- Section 8.4. Problems 7, 9, 15, 17, 25, 31
- Section 8.5. Problems 11, 15, 21, 29

Contents

1	General Information	2
1.1	Warnings and Disclaimers	2
1.2	Recommended Exercises	3
2	Limits	8
2.1	Introduction to Limits	8
2.2	Definition of Limit	10
2.3	Techniques for Finding Limits	11
2.4	Limits Involving Infinity	12
2.5	Continuous Functions	13
3	Derivative	16
3.1	Tangent Lines and Rates of Changes	16
3.2	Definition of Derivative	17
3.3	Techniques of Differentiation	17
3.4	Derivatives of the Trigonometric Functions	18
3.5	The Chain Rule	19
3.6	Implicit Differentiation	19
3.7	Related Rates	20
3.8	Linear Approximations and Differentials	20
4	Applications of Derivative	22
4.1	Extrema of Functions	22
4.2	The Mean Value Theorem	24
4.3	The First Derivative Test	25
4.4	Concavity and the Second Derivative Test	26
4.5	Summary of Graphical Methods	27
4.6	Optimization Problems. Review	28
5	Integrals	29
5.1	Antiderivatives and Indefinite Integrals	29

5.2	Change of Variables in Indefinite Integrals	31
5.3	Summation Notation and Area	31
5.4	The Definite Integral	32
5.5	Properties of the Definite Integral	33
5.6	The Fundamental Theorem of Calculus	34
6	Applications of the Definite Integral	36
6.1	Area	36
6.2	Solids of Revolution	37
6.3	Volumes by Cylindrical Shells	37
6.4	Volumes by Cross Section	38
6.5	Arc Length and Surfaces of Revolution	39
7	Trancendental Functions	40
7.1	The Derivative of the Inverse Function	40
7.2	The Natural Logarithm Function	41
7.3	The Exponential Function	43
7.4	Integration Logarithm and Exponents	44
7.5	General Exponential and Logarithmic Functions	44
7.7	Inverse Trigonometric Functions	46
7.8	Hyperbolic Functions	47
7.9	l'Hospital's Rule	49
8	Techniques of Integration	51
8.1	Integration by Parts	51
8.2	Trigonometric Integrals	51
8.3	Trigonometric Substitution	52
8.4	Integrals of Rational Functions	53
8.5	Quadratic Expressions	53
8.6	Improper Integrals	54
9	Infinite Series	56
9.1	Sequences	56
9.2	Convergent or Divergent Series	57
9.3	Positive-Term Series	58
9.4	The Ratio and Root Tests	59
9.5	Alternating Series and Absolute Convergence	60
A	Algebra	62
A.1	Numbers	62
A.2	Polynomial. Factorization of Polynomials	62

A.3	Binomial Formula	63
A.4	Real Axis	63
A.5	Absolute Value	64
A.6	Inequalities	64
B	Function and Their Graph	66
B.1	Rectangular (Cartesian) Coordinates	66
B.2	Graph of an Equation	67
B.3	Line Equations	67
B.4	Symmetries and Shifts	67
B.5	Definition of a Function. Domain and Range	68
B.6	One-to-One Functions. Periodic Functions	69
B.7	Inc/Decreasing, Odd/Even Functions	69
C	Conic Section	71
C.1	Circle	71
C.2	Parabola	71
C.3	Ellipse	72
C.4	Hyperbola	74
C.5	Conclusion	75
D	Trigonometric Functions	76
E	Exponential and Logarithmic Functions	78

Chapter 2

Limits

The notion of *limits* is central for calculus almost any other notion of calculus (continuity, convergence of a sequence, derivative, integral, etc.) is based on limits. Thus it is very important to be command with limits.

2.1 Introduction to Limits

The following is a quote from a letter of Donald E. Knuth to “Notices of AMS”.

The most important of these changes would be to introduce the O notation and related ideas at an early stage. This notation, first used by Bachmann in 1894 and later popularized by Landau, has the great virtue that it makes calculations simpler, so it simplifies many parts of the subject, yet it is highly intuitive and easily learned. The key idea is to be able to deal with quantities that are only partly specified, and to use them in the midst of formulas.

I would begin my ideal calculus course by introducing a simpler “*A notation*,” which means “absolutely at most.” For example, $A(2)$ stands for a quantity whose absolute value is less than or equal to 2. This notation has a natural connection with decimal numbers: Saying that π is approximately 3.14 is equivalent to saying that $\pi = 3.14 + A(.005)$. Students will easily discover how to calculate with A :

$$\begin{aligned}10^{A(2)} &= A(100); \\(3.14 + A(.005))(1 + A(0.01)) &= 3.14 + A(.005) + A(0.0314) + A(.00005) \\ &= 3.14 + A(0.3645) = 3.14 + A(.04).\end{aligned}$$

I would of course explain that the equality sign *is not symmetric* with respect to such notations; we have $3 = A(5)$ and $4 = A(5)$ but not $3 = 4$, nor can

we say that $A(5) = 4$. We can, however, say that $A(0) = 0$. As de Bruijn points out in [1, §1.2], mathematicians customarily use the $=$ sign as they use the word “is” in English: Aristotle is a man, but a man isn’t necessarily Aristotle.

The A notation applies to variable quantities as well as to constant ones. For example,

$$\begin{aligned}\sin x &= A(1); \\ x &= A(x); \\ A(x) &= xA(1); \\ A(x) + A(y) &= A(x + y) \text{ if } x \geq 0 \text{ and } y \geq 0; \\ (1 + A(t))^2 &= 1 + 3A(t) \text{ if } t = A(1).\end{aligned}$$

Once students have caught on to the idea of A notation, they are ready for O notation, which is even less specific. In its simplest form, $O(x)$ stands for something that is $CA(x)$ for some constant C , but we don’t say what C is. We also define side conditions on the variables that appear in the formulas. For example, if n is a positive integer we can say that any quadratic polynomial in n is $O(n^2)$. If n is sufficiently large, we can deduce that

$$\begin{aligned}(n + O(\sqrt{n}))(\ln n + \gamma + O(1/n)) &= n \ln n + \gamma n + O(1) + O(\sqrt{n} \ln n) + O(\sqrt{n}) + O(1/\sqrt{n}) \\ &= n \ln n + \gamma n + O(\sqrt{n} \ln n).\end{aligned}$$

I’m sure it would be a pleasure for both students and teacher if calculus were taught in this way. The extra time needed to introduce O notation is amply repaid by the simplifications that occur later. In fact, there probably will be time to introduce the “ o notation,” which is equivalent to the taking of limits, and to give the general definition of a not-necessarily-strong derivative:

$$f(x + \epsilon) = f(x) + f'(x)\epsilon + o(\epsilon).$$

The function f is continuous at x if

$$f(x + \epsilon) = f(x) + o(1);$$

and so on. But I would not mind leaving a full exploration of such things to a more advanced course, when it will easily be picked up by anyone who has learned the basics with O alone. Indeed, I have not needed to use “ o ” in 2200 pages of *The Art of Computer Programming*, although many techniques of advanced calculus are applied throughout those books to a great variety of problems.

Students will be motivated to use O notation for two important reasons. First, it significantly simplifies calculations because it allows us to be sloppy—but in a satisfactorily controlled way. Second, it appears in the power series calculations of symbolic algebra systems like *Maple* and *Mathematica*, which today's students will surely be using.

[1]: N. G. de Bruijn, *Asymptotic Methods in Analysis* (Amsterdam: North-Holland, 1958). [2]: R. L. Graham, D. E. Knuth, and O. Patashnik, *Concrete Mathematics* (Reading, Mass.: Addison–Wesley, 1989).

2.2 Definition of Limit

Definition 2.2.1 Let a function f be defined on an open interval containing a , except possibly at a itself, and let L be a real number. The statement

$$\lim_{x \rightarrow a} f(x) = L \quad (2.2.1)$$

(L is the *limit* of function f at a) means that for every $\epsilon > 0$, there is a $\delta > 0$ such that if $0 < |x - a| < \delta$, then $|f(x) - L| < \epsilon$.

We introduce a special kind of limits:

Definition 2.2.2 We say that variable y is $o(z)$ (*o notation*) in the neighborhood of a point a if for every $\epsilon > 0$ there exists $\delta > 0$ such that if $0 < |x - a| < \delta$, then $|y| < \epsilon |z|$.

Then we could define limit of a function as follows

Definition 2.2.3 A function $f(x)$ has a limit L at point a if for $x \neq 0$

$$f(a + x) = L + o(1).$$

Theorem 2.2.4 If

$$\lim_{x \rightarrow a} f(x) = L$$

and $L > 0$, then there is an open interval $(a - \delta, a + \delta)$ containing a such that $f(x) > 0$ for every x in $(a - \delta, a + \delta)$, except possibly $x = a$.

Exercise 2.2.5 Verify limits using Definition

$$\begin{aligned} \lim_{x \rightarrow 3} (5x + 3) &= 18; \\ \lim_{x \rightarrow 2} (x^2 + 1) &= 5. \end{aligned}$$

2.3 Techniques for Finding Limits

Theorem 2.3.1 *The following basic limits are:*

$$\lim_{x \rightarrow a} c = c; \quad (2.3.1)$$

$$\lim_{x \rightarrow a} x = a. \quad (2.3.2)$$

Theorem 2.3.2 *If both limits*

$$\lim_{x \rightarrow a} f(x) \text{ and } \lim_{x \rightarrow a} g(x)$$

exist, then

$$\lim_{x \rightarrow a} [f(x) \pm g(x)] = \lim_{x \rightarrow a} f(x) \pm \lim_{x \rightarrow a} g(x); \quad (2.3.3)$$

$$\lim_{x \rightarrow a} [f(x) \cdot g(x)] = \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x). \quad (2.3.4)$$

$$\lim_{x \rightarrow a} \left[\frac{f(x)}{g(x)} \right] = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}, \quad (2.3.5)$$

provided

$$\lim_{x \rightarrow a} g(x) \neq 0.$$

Corollary 2.3.3 (i). *From formulas (2.3.1) and (2.3.4) follows*

$$\lim_{x \rightarrow a} [cf(x)] = c \lim_{x \rightarrow a} f(x).$$

(ii). *If a, m, b are real numbers then*

$$\lim_{x \rightarrow a} (mx + b) = ma + b.$$

(iii). *If n is a positive integer then*

$$\begin{aligned} \lim_{x \rightarrow a} x^n &= a^n. \\ \lim_{x \rightarrow a} [f(x)]^n &= \left[\lim_{x \rightarrow a} f(x) \right]^n, \end{aligned}$$

provided there exists the limit

$$\lim_{x \rightarrow a} f(x).$$

(iv). *If $f(x)$ is a polynomial function and a is a real number, then*

$$\lim_{x \rightarrow a} f(x) = f(a).$$

(v). If $q(x)$ is a rational function and a is in the domain of q , then

$$\lim_{x \rightarrow a} q(x) = q(a).$$

Exercise 2.3.4 Find limits

$$\begin{aligned} & \lim_{x \rightarrow 2} \sqrt{3}; & \lim_{x \rightarrow -3} x; \\ & \lim_{x \rightarrow 4} (3x + 1); & \lim_{x \rightarrow -2} (2x - 1)^{15}; \\ & \lim_{x \rightarrow 1/2} \frac{2x^2 + 5x - 3}{6x^2 - 7x + 2}; & \lim_{x \rightarrow -2} \frac{x^2 + 2x - 3}{x^2 + 5x + 6}; \\ & \lim_{x \rightarrow \pi} \sqrt[5]{\frac{x - \pi}{x + \pi}}; & \lim_{h \rightarrow 0} \left(\frac{1}{h}\right) \left(\frac{1}{\sqrt{1+h}} - 1\right) \end{aligned}$$

Theorem 2.3.5 If $a > 0$ and n is a positive integer, or if $a \leq 0$ and n is an odd positive integer, then

$$\lim_{x \rightarrow a} \sqrt[n]{x} = \sqrt[n]{a}.$$

Theorem 2.3.6 (Sandwich Theorem) Suppose $f(x) \leq h(x) \leq g(x)$ for every x in an open interval containing a , except possibly at a . If

$$\lim_{x \rightarrow a} f(x) = L = \lim_{x \rightarrow a} g(x),$$

then

$$\lim_{x \rightarrow a} h(x) = L.$$

Exercise 2.3.7 Find limits

$$\lim_{x \rightarrow 0} x^2 \sin \frac{1}{x^2}; \quad \lim_{x \rightarrow \pi/2} \left(x - \frac{\pi}{2}\right) \cos x.$$

REMARK 2.3.8 All such type of result could be modified for *one-sided* limits.

2.4 Limits Involving Infinity

Definition 2.4.1 Let a function f be defined on an infinite interval (c, ∞) (respectively $(-\infty, c)$) for a real number c , and let L be a real number. The statement

$$\lim_{x \rightarrow \infty} f(x) = L \quad \left(\lim_{x \rightarrow -\infty} f(x) = L \right)$$

means that for every $\epsilon > 0$ there is a number M such that if $x > M$ ($x < M$), then $|f(x) - L| < \epsilon$.

Theorem 2.4.2 If k is a positive rational number and c , then

$$\lim_{x \rightarrow \infty} \frac{c}{x^k} = 0 \text{ and } \lim_{x \rightarrow -\infty} \frac{c}{x^k} = 0$$

Definition 2.4.3 Let a function f be defined on an open interval containing a , except possibly at a itself. The statement

$$\lim_{x \rightarrow a} f(x) = \infty$$

means that for every $M > 0$, there is a $\delta > 0$ such that if $0 < |x - a| < \delta$, then $f(x) > M$.

Exercise 2.4.4 Find limits

$$\lim_{x \rightarrow -1} \frac{2x^2}{x^2 - x - 1} \quad \lim_{x \rightarrow 9/2} \frac{3x^2}{(2x - 9)^2} \quad (2.4.1)$$

$$\lim_{x \rightarrow \infty} \frac{-x^3 + 2x}{2x^2 - 3} \quad \lim_{x \rightarrow -\infty} \frac{2x^2 - x + 3}{x^3 + 1} \quad (2.4.2)$$

2.5 Continuous Functions

The notion of *continuity* is absorbed from our every day life. Here is its mathematical definition

Definition 2.5.1 A function f is *continuous* at a point c if

$$\lim_{x \rightarrow c} f(x) = f(c).$$

If function is not continuous at c then it is *discontinuous* at c , or that f has *discontinuity* at c . We give names to the following types of discontinuities:

(i). *Removable* discontinuity:

$$\lim_{x \rightarrow c} f(x) \neq f(c).$$

(ii). *Jump* discontinuity:

$$\lim_{x \rightarrow +c} f(x) \neq \lim_{x \rightarrow -c} f(x).$$

(iii). *Infinite* discontinuity:

$$\lim_{x \rightarrow \pm c} f(x) = \pm\infty.$$

Exercise 2.5.2 Classify discontinuities of

$$f(x) = \begin{cases} -x^2 & \text{if } x < 1 \\ 2 & \text{if } x = 1 \\ (x-2)^{-1} & \text{if } x > 1. \end{cases}$$

Theorem 2.5.3 (i). A polynomial function f is continuous at every real point c .

(ii). A rational function $q = f/g$ is continuous at every number except the numbers c such that $g(c) = 0$.

PROOF. The proof follows directly from Corollary 2.3.3. \square

Exercise 2.5.4 Find all points at which f is discontinuous

$$f(x) = \frac{x-1}{x^2+x-2}.$$

Definition 2.5.5 If a function f is continuous at every number in an open interval (a, b) we say that f is continuous on the *interval* (a, b) . We say also that f is continuous on the *interval* $[a, b]$ if it is continuous on (a, b) and

$$\lim_{x \rightarrow \pm a} f(x) = a \quad \lim_{x \rightarrow \pm b} f(x) = b.$$

Theorem 2.5.6 If two functions f and g are continuous at a real point c , the following functions are also continuous at c :

(i). the sum $f + g$.

(ii). the difference $f - g$.

(iii). the product fg .

(iv). the quotient f/g , provided $g(c) \neq 0$.

PROOF. Proof follows directly from the Theorem 2.3.2. \square

Exercise 2.5.7 Find all points at which f is continuous

$$f(x) = \frac{\sqrt{9-x}}{\sqrt{x-6}} \quad f(x) = \frac{x-1}{\sqrt{x^2-1}}.$$

Theorem 2.5.8 (i). If

$$\lim_{x \rightarrow c} g(x) = b$$

and f is continuous at b , then

$$\lim_{x \rightarrow c} f(g(x)) = f(b) = f(\lim_{x \rightarrow c} g(x)).$$

(ii). If g is continuous at c and if f is continuous at $g(c)$, then the composite function $f \circ g$ is continuous at c .

Exercise 2.5.9 Suppose that

$$f(x) = \begin{cases} c^2x, & \text{if } x < 1 \\ 3cx - 2 & \text{if } x \geq 1. \end{cases}$$

Determine all c such that f is continuous on \mathbb{R} .

Theorem 2.5.10 (Intermediate Value Theorem) If f is continuous on a closed interval $[a, b]$ and if w is any number between $f(a)$ and $f(b)$, then there is at least one point $c \in [a, b]$ such that $f(c) = w$.

Corollary 2.5.11 If $f(a)$ and $f(b)$ have opposite signs, then there is a number c between a and b such that $f(x) = 0$.

Exercise 2.5.12 Let $f(x) = x^7 + 3x + 2$ and $g(x) = -10x^6 + 3x^2 - 1$. Show that there is a solution of the equation $f(x) = g(x)$ on the interval $(-1, 0)$.

Chapter 3

Derivative

3.1 Tangent Lines and Rates of Changes

Let us construct a secant line to a graph of a function $f(x)$ through the points $(a, f(a))$ and $(a + h, f(a + h))$. Then from the formula (B.3.2) it will have a slope (see (B.3.2))

$$m = \frac{f(a + h) - f(a)}{(a + h) - a} = \frac{f(a + h) - f(a)}{h}. \quad (3.1.1)$$

If $h \rightarrow 0$ then secant line became a tangent and we obtain a formula for its slope

$$m_t = \lim_{h \rightarrow 0} \frac{f(a + h) - f(a)}{h}. \quad (3.1.2)$$

Thus the *equation of tangent line* could be written as follows (see page 67)

$$y - f(a) = m_t(x - a) \text{ or } y = f(a) + m_t(x - a). \quad (3.1.3)$$

If a body pass a distance d within time t then *average velocity* is defined as

$$v_{av} = \frac{d}{t}.$$

If the time interval $t \rightarrow 0$ then we obtain *instantaneous velocity*

$$v_a = \lim_{t \rightarrow 0} \frac{s(a + t) - s(a)}{t}. \quad (3.1.4)$$

These and many other examples lead to the notion of *derivative*.

3.2 Definition of Derivative

The following is a quote from a letter of Donald E. Knuth to “Notices of AMS”.

I would define the derivative by first defining what might be called a “strong derivative”: The function f has a strong derivative $f'(x)$ at point x if

$$f(x + \epsilon) = f(x) + f'(x)\epsilon + O(\epsilon^2)$$

whenever ϵ is sufficiently small. The vast majority of all functions that arise in practical work have strong derivatives, so I believe this definition best captures the intuition I want students to have about derivatives.

Definition 3.2.1 The *derivative* of a function f is the function f' whose value at x is given by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}, \quad (3.2.1)$$

provided the limit exists.

3.3 Techniques of Differentiation

We see immediately, for example, that if $f(x) = x^2$ we have

$$(x + \epsilon)^2 = x^2 + 2x\epsilon + \epsilon^2,$$

so the derivative of x^2 is $2x$. And if the derivative of x^n is $d_n(x)$, we have

$$\begin{aligned} (x + \epsilon)^{n+1} &= (x + \epsilon)(x^n + d_n(x)\epsilon + O(\epsilon^2)) \\ &= x^{n+1} + (xd_n(x) + x^n)\epsilon + O(\epsilon^2); \end{aligned} \quad (3.3.1)$$

hence the derivative of x^{n+1} is $xd_n(x) + x^n$ and we find by induction that $d_n(x) = nx^{n-1}$. Similarly if f and g have strong derivatives $f'(x)$ and $g'(x)$, we readily find

$$f(x + \epsilon)g(x + \epsilon) = f(x)g(x) + (f'(x)g(x) + f(x)g'(x))\epsilon + O(\epsilon^2)$$

and this gives the strong derivative of the product. The *chain rule*

$$f(g(x + \epsilon)) = f(g(x)) + f'(g(x))g'(x)\epsilon + O(\epsilon^2) \quad (3.3.2)$$

also follows when f has a strong derivative at point $g(x)$ and g has a strong derivative at x .

It is also follows that

Theorem 3.3.1 *If a function f is differentiable at a , then f is continuous at a .*

Exercise 3.3.2 Give an example of a function f which is continuous at point $x = 0$ but is not differentiable there.

It could be similarly proven the following

Theorem 3.3.3 *Let f and g be differentiable functions at point c then the following functions are differentiable at c also and derivative could be calculated as follows:*

(i). *sum and difference* $(f \pm g)'(x) = f'(x) \pm g'(x)$.

(ii). *product* $(fg)'(x) = f'(x)g(x) + f(x)g'(x)$.

(iii). *fraction*

$$\left(\frac{f}{g}\right)'(x) = \frac{f'(x)g(x) - f(x)g'(x)}{g^2(x)}.$$

3.4 Derivatives of the Trigonometric Functions

Before find derivatives of trigonometric functions we will need the following:

Theorem 3.4.1 *The following limits are:*

$$\lim_{\phi \rightarrow 0} \sin \phi = 0; \quad (3.4.1)$$

$$\lim_{\phi \rightarrow 0} \cos \phi = 1; \quad (3.4.2)$$

$$\lim_{\phi \rightarrow 0} \frac{\sin \phi}{\phi} = 1. \quad (3.4.3)$$

PROOF. Only the last limit is non-trivial. It follows from obvious inequalities

$$\sin \phi < \phi < \tan \phi$$

and the Sandwich Theorem 2.3.6. \square

Corollary 3.4.2 *The following limit is*

$$\lim_{x \rightarrow 0} \frac{1 - \cos \phi}{\phi} = 0.$$

Exercise 3.4.3 Find following limits if they exist:

$$\lim_{t \rightarrow 0} \frac{4t^2 + 3t \sin t}{t^2}; \quad \lim_{t \rightarrow 0} \frac{\cos t}{1 - \sin t};$$

$$\lim_{x \rightarrow 0} x \cot x; \quad \lim_{x \rightarrow 0} \frac{\sin 3x}{\sin 5x}.$$

We are able now to calculate derivatives of trigonometric functions

Theorem 3.4.4

$$\sin' x = \cos x; \quad \cos' x = -\sin x; \quad (3.4.4)$$

$$\tan' x = \sec^2 x; \quad \cot' x = -\csc^2 x; \quad (3.4.5)$$

$$\sec' x = \sec x \tan x; \quad \csc' x = -\csc x \cot x. \quad (3.4.6)$$

PROOF. The proof easily follows from the Theorem 3.4.1 and trigonometric identities on page 77. \square

Exercise 3.4.5 Find derivatives of functions

$$y = \frac{\sin x}{x}; \quad y = \frac{1 - \cos z}{1 + \cos z};$$

$$y = \frac{\tan x}{1 + x^2}; \quad y = \csc t \sin t.$$

3.5 The Chain Rule

The *chain rule*

$$(f \circ g)'(x) = f'(g(x))g'(x) \quad (3.5.1)$$

was proven above 3.3.2.

Exercise 3.5.1 Find derivatives

$$y = \left(z^2 - \frac{1}{z^2}\right)^3; \quad y = \frac{x^4 - 3x^2 + 1}{(2x + 3)^4};$$

$$y = \sqrt[3]{8r^3 + 27}; \quad y = (7x + \sqrt{x^2 + 3})^6.$$

3.6 Implicit Differentiation

If a function $f(x)$ is given by formula like $f(x) = 2x^7 + 3x - 1$ then we will say that it is an *explicit function*. In contrast an identity like

$$y^4 + 3y - 4x^3 = 5x + 1$$

define an *implicit function*. The derivative of implicit function could be found from an equation which it is defined. Usually it is a function both x and y . This procedure is called *implicit differentiation*.

Exercise 3.6.1 Find the slope of the tangent lines at given points

(i). $x^2y + \sin y = 2\pi$ at $P(1, 2\pi)$.

(ii). $2x^3 - x^2y + y^3 - 1 = 0$ at $P(2, -3)$.

3.7 Related Rates

If two variables x and y satisfy to some relationship then we could find their *related rates* by the implicit differentiation.

Exercise 3.7.1 (i). If $S = z^3$ and $dz/dt = -2$ when $z = 3$, find dS/dt .

(ii). If $x^2 + 3y^2 + 2y = 10$ and $dx/dt = 2$ when $x = 3$ and $y = -1$, find dy/dt .

Exercise 3.7.2 Suppose a spherical snowball is melting and the radius is decreasing at a constant rate, changing from 12 in. to 8 in. in 45 min. How fast was the volume changing when the radius was 10 in.?

3.8 Linear Approximations and Differentials

It is known from geometry that a tangent line is closest to a curve at given point among all lines. Thus equation of a tangent line (3.1.3)

$$y = f(a) + f'(a)(x - a) \quad (3.8.1)$$

gives the best approximation to a given graph of $f(x)$. We could use this *linear approximation* in order to estimate value of $f(x)$ in a vicinity of a . We denote an increment of the independent variable x by Δx and

$$\Delta y = f(x_0 + \Delta x) - f(x)$$

is the increment of dependent variable. Therefore

$$\Delta y \approx f'(x)\Delta x \text{ if } \Delta x \approx 0; \quad (3.8.2)$$

$$f(x) \approx f(x) + f'(x)\Delta x; \quad (3.8.3)$$

$$f(x) \approx f(x) + dy, \quad (3.8.4)$$

where $dy = f'(x)\Delta x$ is defined to be *differential* of $f(x)$.

An application of this formulas connected with estimation of errors of measurements:

	Exact value	Approximate value
Absolute error	$\Delta y = y - y_0$	$dy = f'(x_0)\Delta x$
Relative error	$\frac{\Delta y}{y_0}$	$\frac{dy}{y_0}$
Percentage error	$\frac{\Delta y}{y_0} \times 100\%$	$\frac{dy}{y_0} \times 100\%$

Exercise 3.8.1 Use linear approximation to estimate $f(b)$:

(i). $f(x) = -3x^2 + 8x - 7$; $a = 4$, $b = 3.96$.

(ii). $f(\phi) = \csc \phi + \cot \phi$, $a = 45^\circ$, $b = 46^\circ$.

Exercise 3.8.2 Find Δy , dy , $dy - \Delta y$ for $y = 3x^2 + 5x - 2$.

Chapter 4

Applications of Derivative

4.1 Extrema of Functions

We defined *increasing* and *decreasing* functions in Section B.7. There are more definitions

Definition 4.1.1 Let f is defined on $S \subset \mathbb{R}$ and $c \in S$.

- (i). $f(c)$ is the *maximum* value of f on S if $f(x) \leq f(c)$ for every $x \in S$.
- (ii). $f(c)$ is the *minimum* value of f on S if $f(x) \geq f(c)$ for every $x \in S$.

Maximum and minimum are called *extreme values*, or *extrema* of f . If S is the domain of f then maximum and minimum are called *global* or *absolute*.

Exercise 4.1.2 Give an examples of functions which do not have minimum or maximum values.

The important property of continuous functions is given by the following

Theorem 4.1.3 *If f is continuous on $[a, b]$, then f takes on a maximum and minimum values at least once in $[a, b]$.*

Sometimes the following notions are of great importance

Definition 4.1.4 Let c be a number in domain of f .

- (i). $f(c)$ is the *local maximum* f if there is an open interval (a, b) such that $c \in (a, b)$ and $f(x) \leq f(c)$ for every $x \in (a, b)$ in the domain of f .
- (ii). $f(c)$ is the *local minimum* f if there is an open interval (a, b) such that $c \in (a, b)$ and $f(x) \geq f(c)$ for every $x \in (a, b)$ in the domain of f .

The local extrema could be determined from values of derivative:

Theorem 4.1.5 *If f has a local extremum at a number c in an open interval, then either $f'(c) = 0$ or $f'(c)$ do not exist.*

PROOF. The proof follows from the linear approximation of Δf by $f'(c)\Delta x$: if $f'(c)$ exists and $f'(c) \neq 0$ then in an open interval around c there is values of f whose greater and less than $f(c)$. \square

The direct consequence is:

Corollary 4.1.6 *if $f'(c)$ exists and $f'(c) \neq 0$ then $f(c)$ is not a local extremum of c .*

Critical numbers of f are whose points c in the domain of f where either $f'(c) = 0$ or $f'(c)$ does not exist.

Theorem 4.1.7 *If a function f is continuous on a $[a, b]$ and has its maximum or minimum values at a number $c \in (a, b)$, then either $f'(c) = 0$ or $f'(c)$ does not exist.*

So to determine maximum and minimum values of f one should accomplish the following steps

- (i). Find all critical points of f on (a, b) .
- (ii). Calculate values of f in all critical points from step 4.1.7.(i).
- (iii). Calculate the endpoint values $f(a)$ and $f(b)$.
- (iv). The maximal and minimal values of f on $[a, b]$ are the largest and smallest values calculated in 4.1.7.(ii) and 4.1.7.(iii).

Exercise 4.1.8 Find extrema of f on the interval

(i). $y = x^4 - 5x + 4; [0, 2]$.

(ii). $y = (x - 1)^{2/3} - 4; [0, 9]$.

Exercise 4.1.9 Find the critical numbers of f

$$y = 4x^3 + 5x^2 - 42x + 7;$$

$$y = \sqrt[3]{x^2 - x - 2};$$

$$y = (4z + 1)\sqrt{z^2 - 16};$$

$$y = 8 \cos^3 t - 3 \sin 2x - 6x.$$

4.2 The Mean Value Theorem

Theorem 4.2.1 (Rolle's Theorem) *If f is continuous on a closed interval $[a, b]$ and differentiable on the open interval (a, b) and if $f(a) = f(b)$, then $f'(c) = 0$ for at least one number c in (a, b) .*

PROOF. There is two possibilities

- (i). f is constant on $[a, b]$ then $f'(x) = 0$ everywhere.
- (ii). $f(x)$ is not constant then it has at least one extremum point c (Theorem 4.1.3) which is not the end point of $[a, b]$, then $f'(c) = 0$ (Theorem 4.1.5).

□

Exercise 4.2.2 Shows that f satisfy to the above theorem and find c :

- (i). $f(x) = 3x^2 - 12x + 11$; $[0, 4]$.
- (ii). $f(x) = x^3 - x$; $[-1, 1]$.

Rolle's Theorem is the principal step to the next

Theorem 4.2.3 (Mean Value Theorem or Lagrange's Theorem) *If f is continuous on a closed interval $[a, b]$ and differentiable on the open interval (a, b) , then there exists a number $c \in (a, b)$ such that*

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

or, equivalently,

$$f(b) - f(a) = f'(c)(b - a).$$

PROOF. The proof follows from application of the Rolle's Theorem 4.2.1 to the function

$$g(x) = f(x) - \frac{f(b) - f(a)}{b - a}(x - a).$$

□

We start applications of Mean Value Theorem by two corollaries:

Corollary 4.2.4 *If $f'(x) = 0$ for all x in some interval I , then there is a constant C such that $f(x) = C$ for all x in I .*

Corollary 4.2.5 If $f'(x) = g'(x)$ for all $x \in I$, then there is a constant C such that $f(x) = g(x) + C$.

Exercise 4.2.6 Shows that f satisfy to the above theorem and find c :

- (i). $f(x) = 5x^2 - 3x + 1$; $[1, 3]$.
- (ii). $f(x) = x^{2/3}$; $[-8, 8]$.
- (iii). $f(x) = x^3 + 4x$; $[-3, 6]$.

Exercise 4.2.7 Prove: if f continuous on $[a, b]$ and if $f'(x) = c$ there, then $f(x) = cx + d$ for a $d \in \mathbb{R}$.

Exercise 4.2.8 Prove:

$$|\sin u - \sin v| \leq |u - v|.$$

4.3 The First Derivative Test

Derivative of function could provide future information on its behavior:

Theorem 4.3.1 Let f be continuous on $[a, b]$ and differentiable on (a, b) .

- (i). If $f'(x) > 0$ for every x in (a, b) , then f is increasing on $[a, b]$.
- (ii). If $f'(x) < 0$ for every x in (a, b) , then f is decreasing on $[a, b]$.

PROOF. For any numbers x_1 and x_2 in (a, b) we could write using the Mean Value Theorem:

$$f(x_1) - f(x_2) = f'(c)(x_1 - x_2).$$

Then for $x_1 > x_2$ we will have $f(x_1) > f(x_2)$ if $f'(c)$ is positive and $f(x_1) < f(x_2)$ if $f'(c)$ is negative. \square

To check the sign of continuous derivative in an interval $[a, b]$ which does not contain critical points it is enough to verify it for a single point $k \in (a, b)$ (See Intermediate Value Theorem). We shall call $f'(k)$ a *test value*.

Test 4.3.2 (First Derivative Test) Let c is a critical number for f , f is continuous in an open interval I containing c and differentiable in I , except possibly at c itself. Then from the above theorem it follows that

- (i). If f' changes from positive to negative at c , then $f(c)$ is a local maximum of f .

- (ii). If f' changes from negative to positive at c , then $f(c)$ is a local minimum of f .
- (iii). If $f'(x) > 0$ or $f'(x) < 0$ for all $x \in I$, $x \neq c$, then $f(c)$ is not a local extremum of f .

Exercise 4.3.3 Find the local extrema of f and intervals of monotonicity, sketch the graph

- (i). $y = 2x^3 + x^2 - 20x + 1$.
- (ii). $y = 10x^3(x - 1)^2$.
- (iii). $y = x(x^2 - 9)^{1/2}$.
- (iv). $y = x/2 - \sin x$.
- (v). $y = 2 \cos x + \cos 2x$.

Exercise 4.3.4 Find local extrema of f on the given interval

- (i). $y = \cot^2 x + 2 \cot x$ $[\pi/6, 5\pi/6]$.
- (ii). $y = \tan x - 2 \sec x$ $[-\pi/4, \pi/4]$.

4.4 Concavity and the Second Derivative Test

Definition 4.4.1 Let f be differentiable on an open interval I . The graph of f is

- (i). *concave upward* on I if f' is increasing on I ;
- (ii). *concave downward* on I if f' is decreasing on I .

If a graph is concave upward then it lies above any tangent line and for downward concavity it lies below every tangent line.

Test 4.4.2 (Test for Concavity) If the second derivative f'' of f exists on an open interval I , then the graph of f is

- (i). *concave upward* on I if $f''(x) > 0$ on I ;
- (ii). *concave downward* on I if $f''(x) < 0$ on I .

Definition 4.4.3 A point $(c, f(c))$ on the graph of f is a *point of inflection* if the following conditions are satisfied:

- (i). f is continuous at c .
- (ii). There is an open interval (a, b) containing c such that the graph has different types of concavity on (a, c) and (c, b) .

Test 4.4.4 (Second Derivative Test) Suppose that f is differentiable on an open interval containing c and $f'(c) = 0$.

- (i). If $f''(c) < 0$, then f has a local maximum at c .
- (ii). If $f''(c) > 0$, then f has a local minimum at c .

Exercise 4.4.5 Find the local extrema of f , intervals of concavity and points of inflections.

- (i). $y = 2x^6 - 6x^4$.
- (ii). $y = x^{1/5} - 1$.
- (iii). $y = 6x^{1/2} - x^{3/2}$.
- (iv). $y = \cos x - \sin x$.
- (v). $y = x + 2 \cos x$.

4.5 Summary of Graphical Methods

In order to sketch a graph the following steps should be performed

- (i). Find domain of f .
- (ii). Estimate range of f and determine region where f is negative and positive.
- (iii). Find region of continuity and classify discontinuity (if any).
- (iv). Find all x - and y -intercepts.
- (v). Find symmetries of f .
- (vi). Find critical numbers and local extrema (using the First or Second Derivative Test), region of monotonicity of f .
- (vii). Determine concavity and points of inflections.
- (viii). Find asymptotes.

Exercise 4.5.1 Sketch the graphs:

$$f(x) = \frac{2x^2 - x - 3}{x - 2};$$

$$f(x) = \frac{8 - x^3}{2x^2};$$

$$f(x) = \frac{-3x}{\sqrt{x^2 + 4}};$$

$$f(x) = x^3 + \frac{3}{x};$$

$$f(x) = \frac{-4}{x^2 + 1};$$

$$f(x) = |x^3 - x|;$$

$$f(x) = |\cos x| + 2.$$

4.6 Optimization Problems. Review

To solve optimization problem one need to translate the problem to a question on extrema of a function of one variable.

Exercise 4.6.1 (i). Find the minimum value of A if $A = 4y + x^2$, where $(x^2 + 1)y = 324$.

(ii). Find the minimum value of C if $C = (x^2 + y^2)^{1/2}$, where $xy = 9$.

Exercise 4.6.2 A metal cylindrical container with an open top is to hold 1 m^3 . If there is no waste in construction, find the dimension that require the least amount of material.

Exercise 4.6.3 Find the points of the graph of $y = x^3$ that is closest to the point $(4, 0)$.

Chapter 5

Integrals

5.1 Antiderivatives and Indefinite Integrals

Definition 5.1.1 A function F is an *antiderivative* of the function f on an interval I if $F'(x) = f(x)$ for every $x \in I$.

It is obvious that if F is an antiderivative of f then $F(x) + C$ is also antiderivative of f for any real constant C . C is called an *arbitrary constant*. It follows from Mean Value Theorem that *every* antiderivative is of this form.

Theorem 5.1.2 Let F be an antiderivative of f on an interval I . If G is any antiderivative of f on I , then

$$G(x) = F(x) + C$$

for some constant C and every $x \in I$.

Exercise 5.1.3 The above Theorem may be false if the domain of f is different from an interval I . Give an example.

Definition 5.1.4 The notation

$$\int f(x) dx = F(x) + C,$$

where $F'(x) = f(x)$ and C is an arbitrary constant, denotes the family of all antiderivatives of $f(x)$ on an interval I and is called *indefinite integral*.

Theorem 5.1.5

$$\int \frac{d}{dx}(f(x)) dx = f(x) + C; \quad (5.1.1)$$

$$\frac{d}{dx} \left(\int f(x) dx \right) = f(x). \quad (5.1.2)$$

The above Theorem allows us to construct a primitive table of antiderivatives from the tables of derivatives:

Derivative	Indefinite Integral	
$\frac{d}{dx}(x) = 1$	$\int 1 dx = \int dx = x + C$	
$\frac{d}{dx}\left(\frac{x^{r+1}}{r+1}\right) = x^r \quad (r \neq -1)$	$\int x^r dx = \frac{x^{r+1}}{r+1} + C \quad (r \neq -1)$	
$\frac{d}{dx}(\sin x) = \cos x$	$\int \cos x dx = \sin x + C$	
$\frac{d}{dx}(-\cos x) = \sin x$	$\int \sin x dx = -\cos x + C$	(5.1.3)
$\frac{d}{dx}(\tan x) = \sec^2 x$	$\int \sec^2 x dx = \tan x + C$	
$\frac{d}{dx}(-\cot x) = \csc^2 x$	$\int \csc^2 x dx = -\cot x + C$	
$\frac{d}{dx}(\sec x) = \sec x \tan x$	$\int \sec x \tan x dx = \sec x + C$	
$\frac{d}{dx}(-\csc x) = \csc x \cot x$	$\int \csc x \cot x dx = -\csc x + C$	

Theorem 5.1.6

$$\int cf(x) dx = c \int f(x) dx \quad (5.1.4)$$

$$\int [f(x) \pm g(x)] dx = \int f(x) dx \pm \int g(x) dx \quad (5.1.5)$$

Exercise 5.1.7 Find antiderivatives of the following functions

$$\begin{aligned}
 &9t^2 - 4t + 3; \quad 4x^2 - 8x + 1; \quad \frac{1}{z^3} - \frac{3}{z^2}; \\
 &\sqrt{u^3} - \frac{1}{2}u^{-2} + 5; \quad (3x - 1)^2; \quad \frac{(t^2 + 3)^2}{t^6}; \\
 &\frac{7}{\csc x}; \quad -\frac{1}{5}\sin x; \quad \frac{1}{\sin^2 t}.
 \end{aligned}$$

A *Differential equation* is an equation that involves derivatives or differentials of an unknown function. Additional values of f or its derivatives are called *initial conditions*.

Exercise 5.1.8 Solve the differential equations subject to the given conditions

$$f'(x) = 12x^2 - 6x + 1, \quad f(1) = 5; \quad (5.1.6)$$

$$f''(x) = 4x - 1, \quad f'(2) = -2, \quad f(1) = 3; \quad (5.1.7)$$

$$\frac{d^2y}{dx^2} = 3 \sin x - 4 \cos x, \quad y = 7, y' = 2 \text{ if } x = 0. \quad (5.1.8)$$

5.2 Change of Variables in Indefinite Integrals

One more important formula for indefinite integral could be obtained from the rules of differentiation. The chain rule implies:

Theorem 5.2.1 *If F is an antiderivative of f , then*

$$\int f(g(x))g'(x) dx = F(g(x)) + C.$$

If $u = g(x)$ and $du = g'(x) dx$, then

$$\int f(u) du = F(u) + C.$$

Exercise 5.2.2 Find the integrals

$$\int x(2x^2 + 3)^{10} dx; \quad \int x^2 \sqrt[3]{3x^3 + 7} dx; \quad (5.2.1)$$

$$\int \left(1 + \frac{1}{x}\right)^{-3} \left(\frac{1}{x^2}\right) dx; \quad \int \frac{t^2 + t}{(4 - 3t^2 - 2t^3)^4} dt; \quad (5.2.2)$$

$$\int \frac{\sin 2x}{\sqrt{1 - \cos 2x}} dx; \quad \int \sin^3 x \cos x dx. \quad (5.2.3)$$

5.3 Summation Notation and Area

Definition 5.3.1 We use the following *summation notation*:

$$\sum_{k=1}^n a_k = a_1 + a_2 + \cdots + a_n.$$

Theorem 5.3.2

$$\sum_{k=1}^n c = cn; \quad (5.3.1)$$

$$\sum_{k=1}^n (a_k \pm b_k) = \sum_{k=1}^n a_k \pm \sum_{k=1}^n b_k; \quad (5.3.2)$$

$$\sum_{k=1}^n ca_k = c \left(\sum_{k=1}^n a_k \right). \quad (5.3.3)$$

Theorem 5.3.3

$$\sum_{k=1}^n k = \frac{n(n+1)}{2}; \quad (5.3.4)$$

$$\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}; \quad (5.3.5)$$

$$\sum_{k=1}^n k^3 = \left[\frac{n(n+1)}{2} \right]^2. \quad (5.3.6)$$

This sum will help us to find inscribed rectangular polygon and circumscribed rectangular polygon.

Exercise 5.3.4 Find the area under the graph of the following functions:

(i). $y = 2x + 3$, from 2 to 4.

(ii). $y = x^2 + 1$, from 0 to 3.

5.4 The Definite Integral

There is a way to calculate an area under the graph of a function $y = f(x)$. We could approximate it by a sum of the form

$$\sum_{k=1}^n f(w_k) \Delta x_k, \quad w_k \in \Delta x_k.$$

It is a *Riemann sum*. The approximation will be precise if will come to the *limit of Riemann sums*:

$$\lim_{\delta x_k \rightarrow 0} \sum_{k=1}^n f(w_k) \delta x_k = L.$$

If this limit exists it called *definite integral* of f from a to b and denoted by:

$$\int_a^b f(x) dx = \lim_{\delta x_k \rightarrow 0} \sum_{k=1}^n f(w_k) \delta x_k = L.$$

If the limit exist we say that f is *integrable function* on $[a, b]$.

5.5 Properties of the Definite Integral

Theorem 5.5.1 *If c is a real number, then*

$$\int_a^b c dx = c(b - a).$$

Theorem 5.5.2 *If f is integrable on $[a, b]$ and c is any real number, then cf is integrable on $[a, b]$ and*

$$\int_a^b cf(x) dx = c \int_a^b f(x) dx.$$

Theorem 5.5.3 *If f and g are integrable on $[a, b]$, then $f \pm g$ is also integrable on $[a, b]$ ($a > b$) and*

$$\int_a^b [f(x) \pm g(x)] dx = \int_a^b [f(x) \pm g(x)] dx.$$

Theorem 5.5.4 *If f is integrable on $[a, b]$ and $f(x) \geq 0$ for all $x \in [a, b]$ then*

$$\int_a^b f(x) dx \geq 0.$$

Corollary 5.5.5 *If f and g are integrable on $[a, b]$ and $f(x) \geq g(x)$ for all $x \in [a, b]$ then*

$$\int_a^b f(x) dx \geq \int_a^b g(x) dx.$$

Theorem 5.5.6 (Mean Value Theorem for Definite Integrals) *If f is continuous on a closed interval $[a, b]$, then there is a number z in the open interval (a, b) such that*

$$\int_a^b f(x) dx = f(z)(b - a).$$

Definition 5.5.7 Let f be continuous on $[a, b]$. The *average value* f_{av} of f on $[a, b]$ is

$$f_{av} = \frac{1}{b - a} \int_a^b f(x) dx.$$

Once it is known that integration is the inverse of differentiation and related to the area under a curve, we can observe, for example, that if f and f' both have strong derivatives at x , then

$$\begin{aligned} f(x + \epsilon) - f(x) &= \int_0^\epsilon f'(x + t) dt \\ &= \int_0^\epsilon (f'(x) + f''(x)t + O(t^2)) dt \\ &= f'(x)\epsilon + f''(x)\epsilon^2/2 + O(\epsilon^3). \end{aligned} \quad (5.5.1)$$

5.6 The Fundamental Theorem of Calculus

There is an unanswered question from the previous section: *Why undefined and defined integrals shared their names and notations?* The answer is given by the following

Theorem 5.6.1 (Fundamental Theorem of Calculus) *Suppose f is continuous on a closed interval $[a, b]$.*

(i). *If the function G is defined by*

$$G(x) = \int_a^x f(t) dt$$

for every x in $[a, b]$, then G is an antiderivative of f on $[a, b]$.

(ii). *If F is any antiderivative of f on $[a, b]$, then*

$$\int_a^b f(x) dx = F(b) - F(a).$$

PROOF. The proof of the first statement follows from the Mean Value Theorem for Definite Integral. End the second part follows from the first and initial condition

$$\int_a^a f(x) dx = 0.$$

□

Corollary 5.6.2 (i). *If f is continuous on $[a, b]$ and F is any antiderivative of f , then*

$$\int_a^b f(x) dx = F(x) \Big|_a^b = F(b) - F(a).$$

(ii).

$$\int_a^b f(x) dx = \left[\int f(x) dx \right]_a^b.$$

(iii). Let f be continuous on $[a, b]$. If $a \leq c \leq b$, then for every x in $[a, b]$

$$\frac{d}{dx} \int_c^x f(t) dt = f(x).$$

Theorem 5.6.3 If $u = g(x)$, then

$$\int_a^b f(g(x))g'(x) dx = \int_{g(a)}^{g(b)} f(u) du.$$

Theorem 5.6.4 Let f be continuous on $[-a, a]$.(i). If f is an even function,

$$\int_{-a}^a f(x) dx = \int_{-a}^a f(x) dx.$$

(ii). If f is an odd function,

$$\int_{-a}^a f(x) dx = 0.$$

Exercise 5.6.5 Calculate integrals

$$\begin{aligned} & \int_{-2}^{-1} \left(x - \frac{1}{x}\right)^2 dx; & \int_1^4 \sqrt{5-x} dx; & \int_{-1}^1 (v^2 - 1)^3 v dv; \\ & \int_0^{\pi/2} 3 \sin\left(\frac{1}{2}x\right) dx; & \int -\pi/6^{\pi/6}(x + \sin 5x) dx; & \int_0^{\pi/3} \frac{\sin x}{\cos^2 x} dx; \\ & \int_{-1}^5 |2x - 3| dx; \end{aligned}$$

Chapter 6

Applications of the Definite Integral

6.1 Area

We know that the geometric meaning of the definite integral of a positive function is the area under the graph. We could calculate areas of more complicated figures by combining several definite integrals.

Exercise 6.1.1 Find areas bounded by the graphs:

(i). $x = y^2, x - y = 2$.

(ii). $y = x^3, y = x^2$.

(iii). $y = x^{2/3}, x = y^2$.

(iv). $y = x^3 - x, y = 0$.

(v). $x = y^3 + 2y^2 - 3y, x = 0$.

(vi). $y = 4 + \cos 2x, y = 3 \sin \frac{1}{2}x$.

Exercise 6.1.2 Express via sums of integrals areas:

(i). $y = \sqrt{x}, y = -x, x = 1, x = 4$.

6.2 Solids of Revolution

Theorem 6.2.1 Let f be continuous on $[a, b]$, and let R be the region bounded by the graph of f , the x -axis, and the vertical lines $x = a$ and $x = b$. The volume V of the solid of revolution generated by revolving R about the x -axis is

$$V = \int_a^b \pi [f(x)]^2 dx.$$

Theorem 6.2.2 (Volume of a Washer)

$$V = \int_a^b \pi ([f(x)]^2 - [g(x)]^2) dx.$$

Exercise 6.2.3 (i). $y = 1/x$, $x = 1$, $x = 3$, $y = 0$; x -axis;

(ii). $y = x^3$, $x = -2$, $y = 0$, x -axis;

(iii). $y = x^2 - 4x$, $y = 0$; x -axis;

(iv). $y = (x - 1)^2 + 1$, $y = -(x - 1)^2 + 3$; x -axis;

Exercise 6.2.4 Find volume of revolution for $y = x^3$, $y = 4x$ rotated around $x = 4$.

6.3 Volumes by Cylindrical Shells

Let a cylindrical shell has outer and inner radiuses as r_1 and r_2 then and altitude h . We introduce the average radius $r = (r_1 + r_2)/2$ and the thickness $\Delta r = r_2 - r_1$. Then its volume is:

$$V = \pi r_1^2 h - \pi r_2^2 h = 2\pi r \Delta r h.$$

Let a region bounded by a function $f(x)$ and x -axis. If we rotate it around the y -axis then it is an easy to observe that the volume of the solid will be as follow:

$$V = \int_a^b 2\pi x f(x) dx.$$

Exercise 6.3.1 Find volumes:

(i). $y = \sqrt{x}$, $x = 4$, $y = 0$, y -axis.

- (ii). $y = x^2, y^2 = 8x, y$ -axis.
- (iii). $y = 2x, y = 6, x = 0, x$ -axis.
- (iv). $y = \sqrt{x+4}, y = 0, x = 0, x$ -axis.

6.4 Volumes by Cross Section

If a plane intersects a solid, then the region common to the plane and the solid is a *cross section* of the solid. There is a simple formula to calculate volumes by cross sections:

Theorem 6.4.1 (Volumes by Cross Sections) *Let S be a solid bounded by planes that are perpendicular to the x -axis at a and b . If, for every x in $[a, b]$, the cross-sectional area of S is given by $A(x)$, there A is continuous on $[a, b]$, then the volume S is*

$$V = \int_a^b A(x) dx.$$

Corollary 6.4.2 (Cavalieri's theorem) *If two solids have equal altitudes and if all cross sections by planes parallel to their bases and at the same distances from their bases have equal areas, then the solids have the same volume.*

Exercise 6.4.3 Let R be the region bounded by the graphs of $x = y^2$ and $x = 9$. Find the volume of the solid that has R as its base if every cross section by a plane perpendicular to the x -axis has the given shape.

- (i). Rectangle of height 2.
- (ii). A quartercircle.

Exercise 6.4.4 Find volume of a pyramid if its altitude is h and the base is a rectangle of dimensions a and $2a$.

Exercise 6.4.5 A solid has as its base the region in xy -plane bounded by the graph of $y^2 = 4x$ and $x = 4$. Find the volume of the solid if every cross section by a plane perpendicular to the y -axis is semicircle.

6.5 Arc Length and Surfaces of Revolution

We say that a function f is *smooth* on an interval if it has a derivative f' that is continuous throughout the interval.

Theorem 6.5.1 *Let f be smooth on $[a, b]$. The arc length of the graph of f from $A(a, f(a))$ to $B(b, f(b))$ is*

$$L_a^b = \int_a^b \sqrt{1 + [f'(x)]^2} dx$$

We could introduce the *arc length function* s for the graph of f on $[a, b]$ is defined by

$$s(x) = \int_a^x \sqrt{1 + [f'(t)]^2} dt.$$

Theorem 6.5.2 *Let f be smooth on $[a, b]$, and let s be the arc length function for the graph of $y = f(x)$ on $[a, b]$. If Δx is an increment in the variable x , then*

$$\frac{ds}{dx} = \sqrt{1 + [f'(x)]^2}; \quad (6.5.1)$$

$$ds = \sqrt{1 + [f'(x)]^2} \Delta x. \quad (6.5.2)$$

Exercise 6.5.3 Find the arc length:

- (i). $y = 2/3x^{2/3}$; $A(1, 2/3)$, $B(8, 8/3)$;
- (ii). $y = 6\sqrt[3]{x^2} + 1$; $A(-1, 7)$, $B(-8, 25)$;
- (iii). $30xy^3 - y^8 = 15$; $A(8/15, 1)$, $B(\frac{271}{240}, 2)$;

Exercise 6.5.4 Find the length of the graph $x^{2/3} + y^{2/3} = 1$.

Theorem 6.5.5 *If f is smooth and $f(x) \geq 0$ on $[a, b]$, then the area S of the surface generated by revolving the graph of f about the x -axis is*

$$S = \int_a^b 2\pi f(x) \sqrt{1 + [f'(x)]^2} dx.$$

Exercise 6.5.6 Find the area of the surface generated by revolving of the graph

- (i). $y = x^3$; $A(1, 1)$, $B(2, 8)$;
- (ii). $8y = 2x^4 + x^{-2}$, $A(1, 3/8)$, $B(2, 129/32)$;
- (iii). $x = 4\sqrt{y}$; $A(4, 1)$, $B(12, 9)$.

Chapter 7

Transcendental Functions

There is a special sort of functions which have a strange name *transcendental*. We will explore the important rôle of these functions in calculus and mathematics in general.

7.1 The Derivative of the Inverse Function

We define *one-to-one functions* in Section B.6. For such function we could give the following definition.

Definition 7.1.1 Let f be a one-to-one function with domain D and range R . A function g with domain R and range D is the *inverse function* of f , if for all $x \in D$ and $y \in R$ $y = f(x)$ iff $x = g(y)$.

Theorem 7.1.2 Let f be a one-to-one function with domain D and range R . If g is a function with domain R and range D , then g is the inverse function of f iff both the following conditions are true:

(i). $g(f(x)) = x$ for every $x \in D$.

(ii). $f(g(x)) = x$ for every $x \in R$.

Exercise 7.1.3 Find inverse function.

(i). $f(x) = \frac{3x+2}{2x-5}$;

(ii). $f(x) = 5x^2 + 2$, $x \geq 0$;

(iii). $f(x) = \sqrt{4 - x^2}$.

Theorem 7.1.4 *If f is continuous and increasing on $[a, b]$, then f has an inverse function f^{-1} that is continuous and increasing on $[f(a), f(b)]$.*

Theorem 7.1.5 *If a differentiable function f has an inverse function $g = f^{-1}$ and if $f'(g(c)) \neq 0$, then g is differentiable at c and*

$$g'(c) = \frac{1}{f'(g(c))}. \quad (7.1.1)$$

PROOF. The formula follows directly from differentiation by the Chain rule the identity $f(g(x)) = x$ (see Theorem 7.1.2). \square

Exercise 7.1.6 Find domain and derivative of the inverse function.

- (i). $f(x) = \sqrt{2x + 3}$;
- (ii). $f(x) = 4 - x^2$, $x \geq 0$;
- (iii). $f(x) = \sqrt{9 - x^2}$, $0 \leq x \leq 3$.

Exercise 7.1.7 Prove that inverse function exists and find slope of tangent line to the inverse function in the given point.

- (i). $f(x) = x^5 + 3x^3 + 2x - 1$, $P(5, 1)$;
- (ii). $f(x) = 4x^5 - (1/x^3)$, $P(3, 1)$;
- (iii). $f(x) = x^5 + x$, $P(2, 1)$.

7.2 The Natural Logarithm Function

We know that antiderivative for a function x^n is $x^{n+1}/(n+1)$. This expression is defined for all $n \neq -1$. This case deserves a special name

Definition 7.2.1 The *natural logarithm function*, denoted by \ln , is defined by

$$\ln x = \int_1^x \frac{1}{t} dt,$$

for every $x > 0$.

From the properties of definite integral follows that

$$\begin{aligned}\ln x &> 0 & \text{if } x > 1; \\ \ln x &< 0 & \text{if } x < 1; \\ \ln x &= 0 & \text{if } x = 1.\end{aligned}$$

Theorem 7.2.2

$$\frac{d}{dx}(\ln x) = \frac{1}{x}.$$

Theorem 7.2.3 *If $u = g(x)$ and g is differentiable, then*

$$\begin{aligned}\frac{d}{dx}(\ln u) &= \frac{1}{u} \frac{du}{dx}, & \text{if } u > 0; \\ \frac{d}{dx}(\ln |u|) &= \frac{1}{u} \frac{du}{dx}, & \text{if } u \neq 0.\end{aligned}$$

Corollary 7.2.4 *The natural logarithm is an increasing function.*

This gives a new way to prove the principal laws of logarithms.

Exercise 7.2.5 Prove using laws of logarithms that

$$\lim_{x \rightarrow +\infty} = +\infty \quad \text{and} \quad \lim_{x \rightarrow 0^-} = -\infty.$$

From this Exercise and Corollary 7.2.4 follows

Corollary 7.2.6 *To every real number x there corresponds exactly one positive real number y such that $\ln y = x$.*

Exercise 7.2.7 Find implicit derivatives:

(i). $3y - x^2 + \ln xy = 2$.

(ii). $y^3 + x^2 \ln y = 5x + 3$.

Another useful application is *logarithmic differentiation* which is given by the formula:

$$\frac{d}{dx} f(x) = f(x) \frac{d}{dx} \ln(f(x)). \quad (7.2.1)$$

It is useful for functions consisting from products and powers of elementary functions.

Exercise 7.2.8 Find derivative of functions using logarithmic differentiation:

$$f(x) = (5x + 2)^3(6x + 1)^2; \quad (7.2.2)$$

$$f(x) = \sqrt{(3x^2 + 2)\sqrt{6x - 7}}; \quad (7.2.3)$$

$$f(x) = \frac{(x^2 + 3)^5}{\sqrt{x + 1}}. \quad (7.2.4)$$

7.3 The Exponential Function

Corollary 7.2.6 justify the following

Definition 7.3.1 The *natural exponential function*, denoted by $\exp x = e^x$, is the inverse of the natural logarithm function. The letter $e (= 2.718281828\dots)$ denotes the positive real number such that $\ln e = 1$.

By the definition

$$\ln e^x = x, \quad x \in \mathbb{R} \quad (7.3.1)$$

$$e^{\ln x} = x, \quad x > 0. \quad (7.3.2)$$

By the same definition we could derive laws of exponents from the laws of logarithms.

Theorem 7.3.2

$$\frac{d}{dx}(e^x) = e^x \quad (7.3.3)$$

$$\frac{d}{dx}(e^{g(x)}) = e^{g(x)} \frac{dg(x)}{dx} \quad (7.3.4)$$

$$(7.3.5)$$

PROOF. The proof of the first identity follows from differentiation of 7.3.1 by the chain rule. The second identity follows from the first one and the chain rule. \square

Exercise 7.3.3 Find implicit derivatives

(i). $xe^y + 2x - \ln(y + 1) = 3$;

(ii). $e^x \cos y = xe^y$.

Exercise 7.3.4 Find extrema and regions of monotonicity:

(i). $f(x) = x^2e^{-2x}$;

(ii). $f(x) = e^{1/x}$.

7.4 Integration Using Natural Logarithm and Exponential Functions

The following formulas are direct consequences of change of variables in definite integral and definition of logarithmic and exponential functions:

$$\int \frac{1}{g(x)} g'(x) dx = \ln |g(x)| + C; \quad (7.4.1)$$

$$\int e^{g(x)} g'(x) dx = e^{g(x)} + C. \quad (7.4.2)$$

From here we could easily derive

Theorem 7.4.1

$$\int \tan u du = -\ln |\cos u| + C; \quad (7.4.3)$$

$$\int \cot u du = \ln |\sin u| + C; \quad (7.4.4)$$

$$\int \sec u du = \ln |\sec u + \tan u| + C; \quad (7.4.5)$$

$$\int \csc u du = \ln |\csc u - \cot u| + C. \quad (7.4.6)$$

Exercise 7.4.2 Evaluate integrals

$$\int \frac{x^3}{x^4 - 5} dx; \quad \int \frac{(2 + \ln x)^{10}}{x} dx;$$

$$\int \frac{e^x}{(e^x + 2)^2} dx; \quad \int \frac{\cot \sqrt[3]{x}}{\sqrt[3]{x^2}} dx.$$

7.5 General Exponential and Logarithmic Functions

Using laws of logarithms we could make

Definition 7.5.1 The *exponential function with base a* is defined as follows:

$$f(x) = a^x = e^{\ln a^x} = e^{x \ln a}. \quad (7.5.1)$$

From this definition the following properties follows

Theorem 7.5.2

$$\begin{aligned}\frac{d}{dx}(a^x) &= a^x \ln a; \\ \frac{d}{dx}(a^u) &= a^u \ln a \frac{du}{dx}; \\ \int a^x dx &= \left(\frac{1}{\ln a}\right) a^x + C; \\ \int a^u du &= \left(\frac{1}{\ln a}\right) a^u + C.\end{aligned}$$

Exercise 7.5.3 Evaluate integrals

$$\begin{aligned}\int 5^{-5x} dx; & \quad \int \frac{(2x+1)^2}{2^x} dx; \\ \int e^e dx; & \quad \int x^5 dx; \\ \int x^{\sqrt{5}} dx; & \quad \int (\sqrt{5})^x dx;\end{aligned}$$

Exercise 7.5.4 The region under the graph of $y = 3^{-x}$ from $x = 1$ to $x = 2$ is revolved about the x -axis. Find the volume of the resulting solid.

Having a^x already defined we could give the following

Definition 7.5.5 The *logarithmic function with base a* $f(x) = \log_a x$ is defined by the condition $y = \log_a x$ iff $x = a^y$.

The following properties follows directly from the definition

Theorem 7.5.6

$$\begin{aligned}\frac{d}{dx}(\log_a x) &= \frac{d}{dx} \left(\frac{\ln x}{\ln a} \right) = \frac{1}{\ln a} \frac{1}{x}; \\ \frac{d}{dx}(\log_a |u|) &= \frac{d}{dx} \left(\frac{\ln |u|}{\ln a} \right) = \frac{1}{\ln a} \cdot \frac{1}{u} \frac{du}{dx}.\end{aligned}$$

Exercise 7.5.7 Find derivatives of functions:

$$\begin{aligned}f(x) &= \log_{\sqrt{3}} \cos 5x; \\ f(x) &= \ln \log x.\end{aligned}$$

7.7 Inverse Trigonometric Functions

We would like now to define inverse trigonometric functions. But there is a problem: inverse functions exist only for one-to-one functions and trigonometric functions are not the such.

Exercise 7.7.1 Prove that a periodic function could not be a one-to-one function.

A way out could be as follows: we restrict a trigonometric function f to an interval I in such a way that f is one-to-one on I and there is no a bigger interval $I' \supset I$ that f is one-to-one on I' .

Definition 7.7.2 (i). The *arcsine* (*inverse sine function*) denoted \arcsin is defined by the condition $y = \arcsin x$ iff $x = \sin y$ for $-1 \leq x \leq 1$ and $-\pi/2 \leq y \leq \pi/2$.

(ii). The *arccosine* (*inverse cosine function*) denoted \arccos is defined by the condition $y = \arccos x$ iff $x = \cos y$ for $-1 \leq x \leq 1$ and $0 \leq y \leq \pi$.

(iii). The *arctangent* (*inverse tangent function*) denoted \arctan is defined by the condition $y = \arctan x$ iff $x = \tan y$ for $x \in \mathbb{R}$ and $-\pi/2 \leq y \leq \pi/2$.

(iv). The *arcsecant* (*inverse secant function*) denoted arcsec is defined by the condition $y = \operatorname{arcsec} x$ iff $x = \sec y$ for $|x| > 1$ and $y \in [0, \pi/2)$ or $y \in [\pi, 3\pi/2)$.

Exercise* 7.7.3 Why there is no a much need to introduce inverse functions for cotangent and cosecant?

Applying Theorem on derivative of an inverse function we could conclude that

Theorem 7.7.4

$$\begin{aligned} \frac{d}{dx}(\arcsin u) &= \frac{1}{\sqrt{1-u^2}} \frac{du}{dx}; \\ \frac{d}{dx}(\arccos u) &= -\frac{1}{\sqrt{1-u^2}} \frac{du}{dx}; \\ \frac{d}{dx}(\arctan u) &= \frac{1}{1+u^2} \frac{du}{dx}; \\ \frac{d}{dx}(\operatorname{arcsec} u) &= \frac{1}{u\sqrt{u^2-1}} \frac{du}{dx}. \end{aligned}$$

As usually we could invert these formulas for taking antiderivatives:

Theorem 7.7.5

$$\begin{aligned}\int \frac{1}{\sqrt{a^2 - u^2}} du &= \arcsin \frac{u}{a} + C = -\arccos \frac{u}{a} + C; \\ \int \frac{1}{a^2 + u^2} du &= \frac{1}{a} \arctan \frac{u}{a} + C; \\ \int \frac{1}{u\sqrt{u^2 - a^2}} du &= \frac{1}{a} \operatorname{arcsec} \frac{u}{a} + C.\end{aligned}$$

And following formulas could be verified by differentiation:

Theorem 7.7.6

$$\begin{aligned}\int \arcsin u \, du &= u \arcsin u + \sqrt{1 - u^2} + C; \\ \int \arccos u \, du &= u \arccos u - \sqrt{1 - u^2} + C; \\ \int \arctan u \, du &= u \arctan u - \frac{1}{2} \ln(1 + u^2) + C; \\ \int \operatorname{arcsec} u \, du &= u \operatorname{arcsec} u - \ln \left| u\sqrt{u^2 - 1} \right| + C.\end{aligned}$$

7.8 Hyperbolic Functions

The following functions arise in many areas of mathematics and applications.

Definition 7.8.1

$$\begin{aligned}\text{hyperbolic sine function : } \sinh x &= \frac{e^x - e^{-x}}{2}; \\ \text{hyperbolic cosine function : } \cosh x &= \frac{e^x + e^{-x}}{2}; \\ \text{hyperbolic tangent function : } \tanh x &= \frac{e^x - e^{-x}}{e^x + e^{-x}}; \\ \text{hyperbolic cotangent function : } \coth x &= \frac{e^x + e^{-x}}{e^x - e^{-x}}.\end{aligned}$$

There are a lot of identities involving hyperbolic functions which are similar to the trigonometric ones. We will mention only few most important of them

Theorem 7.8.2

$$\begin{aligned}\cosh^2 x - \sinh^2 x &= 1; \\ 1 - \tanh^2 x &= (\cosh x)^{-2}; \\ \coth^2 x - 1 &= (\sinh x)^{-2}.\end{aligned}$$

From formula $(e^x)' = e^x$ easily follows the following formulas of differentiation:

Theorem 7.8.3

$$\begin{aligned}\frac{d}{dx}(\sinh x) &= \cosh x; \\ \frac{d}{dx}(\cosh x) &= \sinh x; \\ \frac{d}{dx}(\tanh x) &= (\cosh x)^{-2}; \\ \frac{d}{dx}(\coth x) &= -(\sinh x)^{-2}.\end{aligned}$$

Exercise 7.8.4 Find derivative of functions

$$f(x) = \frac{1 + \cosh x}{1 + \sinh x}; \quad f(x) = \ln |\tanh x|.$$

We again could rewrite these formulas for indefinite integral case:

Theorem 7.8.5

$$\begin{aligned}\int \sinh u \, du &= \cosh u + C; \\ \int \cosh u \, du &= \sinh u + C; \\ \int (\cosh u)^{-2} \, du &= \tanh u + C; \\ \int (\sinh u)^{-2} \, du &= \coth u + C.\end{aligned}$$

Exercise 7.8.6 Evaluate integrals

$$\int \frac{\sinh \sqrt{x}}{\sqrt{x}} \, dx; \quad \int \frac{1}{\cosh^2 3x} \, dx.$$

7.9 Indeterminate Forms and l'Hospital's Rule

In this section we describe a general tool which simplifies evaluation of limits.

Theorem 7.9.1 (Cauchy's Formula) *If f and g are continuous on $[a, b]$ and differentiable on (a, b) and if $g'(x) \neq 0$ for all $x \in (a, b)$, then there is a number $w \in (a, b)$ such that*

$$\frac{f(a) - f(b)}{g(a) - g(b)} = \frac{f'(w)}{g'(w)}.$$

PROOF. The proof follows from the application of Rolle's Theorem to the function

$$h(x) = [f(b) - f(a)]g(x) - [g(b) - g(a)]f(x).$$

□

Theorem 7.9.2 (l'Hospital's Rule) *Suppose that f and g are differentiable on an open interval (a, b) containing c , except possibly at c itself. If $f(x)/g(x)$ has the indeterminate form $0/0$ or ∞/∞ then*

$$\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \lim_{x \rightarrow c} \frac{f'(x)}{g'(x)},$$

provided

$$\lim_{x \rightarrow c} \frac{f'(x)}{g'(x)} \text{ exists or } \lim_{x \rightarrow c} \frac{f'(x)}{g'(x)} = \infty.$$

Exercise 7.9.3 Find the following limits

$$\begin{array}{ll} \lim_{x \rightarrow 1} \frac{x^3 - 3x + 2}{x^2 - 2x - 1}; & \lim_{x \rightarrow 0} \frac{\sin x - x}{\tan x - x}; \\ \lim_{x \rightarrow 0} \frac{x + 1 - e^x}{x^2}; & \lim_{x \rightarrow 0^+} \frac{\ln \sin x}{\ln \sin 2x}; \\ \lim_{x \rightarrow 0} \frac{e^x - e^{-x} - 2 \sin x}{x \sin x}; & \lim_{x \rightarrow \infty} \frac{x \ln x}{x + \ln x}. \end{array}$$

There are more indeterminate forms which could be transformed to the case $0/0$ or ∞/∞ :

(i). $0 \cdot \infty$: write $f(x)g(x)$ as

$$\frac{f(x)}{1/g(x)} \quad \text{or} \quad \frac{g(x)}{1/f(x)}.$$

(ii). 0^0 , 1^∞ , ∞^0 : instead of $f(x)^{g(x)}$ look for the limit L of $g(x) \ln f(x)$.
Then $f(x)^{g(x)} = e^L$.

(iii). $\infty - \infty$: try to pass to a quotient or a product.

Exercise 7.9.4 Find limits if exist.

$$\begin{array}{ll} \lim_{x \rightarrow 0^+} (e^x - 1)^x; & \lim_{x \rightarrow \infty} x^{1/x}; \\ \lim_{x \rightarrow -3^-} \left(\frac{x}{x^2 + 2x - 3} - \frac{4}{x + 3} \right) & \lim_{x \rightarrow \infty} \left(\frac{x^2}{x - 1} - \frac{x^2}{x + 1} \right); \\ \lim_{x \rightarrow 0} (\cot^2 x - \csc^2 x); & \lim_{x \rightarrow 0^+} (1 + 3x)^{\csc x}. \end{array}$$

Chapter 8

Techniques of Integration

We will study more advanced technique of integration.

8.1 Integration by Parts

Among different formulae of differentiation there is one which was not converted to the formulae of integration yet. This is derivative of a product of two functions. We will use it as follows:

Theorem 8.1.1 *If $u = f(x)$ and $v = g(x)$ and if f' and g' are continuous, then*

$$\int u dv = uv - \int v du.$$

Exercise 8.1.2 Evaluate integrals.

$$\begin{aligned} \int x e^{-x} dx; & \quad \int x \sec x \tan x dx; \\ \int x \csc^2 3x dx; & \quad \int x^2 \sin 4x dx; \\ \int e^x \cos x dx; & \quad \int \sin \ln x dx; \\ \int \cos \sqrt{x} dx; & \quad \int \sin^n x dx.. \end{aligned}$$

8.2 Trigonometric Integrals

To evaluate $\int \sin^m x \cos^n x dx$ we use the following procedure:

- (i). **If m is an odd integer:** use the change of variable $u = \cos x$ and express $\sin^2 x = 1 - \cos^2 x$.
- (ii). **If n is an odd integer:** use the change of variable $u = \sin x$ and express $\cos^2 x = 1 - \sin^2 x$.
- (iii). **If m and n are even:** Use half-angle formulas for $\sin^2 x$ and $\cos^2 x$ to reduce the exponents by one-half.

Exercise 8.2.1 Evaluate integrals.

$$\int \sin^3 x \cos^2 x \, dx; \quad \int \sin^4 x \cos^2 x \, dx.$$

To evaluate $\int \tan^m x \sec^n x \, dx$ we use the following procedure:

- (i). **If m is an odd integer:** use the change of variable $u = \sec x$ and express $\tan^2 x = \sec^2 x - 1$.
- (ii). **If n is an even integer:** use the change of variable $u = \tan x$ and express $\sec^2 x = 1 + \tan^2 x$.
- (iii). **If m is an even and n is odd numbers:** There is no a standard method, try the integration by parts.

Exercise 8.2.2 Evaluate integrals.

$$\int \cot^4 x \, dx; \quad \int \sin 4x \cos 3x \, dx.$$

8.3 Trigonometric Substitution

The following trigonometric substitution applicable if integral contains one of the following expression cases

Expression	Substitution
$\sqrt{a^2 - x^2}$	$x = a \sin \theta;$
$\sqrt{a^2 + x^2}$	$x = a \tan \theta;$
$\sqrt{x^2 - a^2}$	$x = a \sec \theta.$

Exercise 8.3.1 Evaluate integrals

$$\int \frac{1}{x^3\sqrt{x^2-25}}dx; \quad \int \frac{1}{x^2\sqrt{x^2+9}}dx;$$

$$\int \frac{1}{(16-x^2)^{5/2}}dx; \quad \int \frac{(4+x^2)^2}{x^3}dx.$$

8.4 Integrals of Rational Functions

To integrate a rational function $f(x)/g(x)$ we accomplish the following steps:

- (i). If the degree of $f(x)$ is not lower than the degree of $g(x)$, use long division to obtain the proper form.
- (ii). Express $g(x)$ as a product of linear factors $ax+b$ or irreducible quadratic factors cx^2+dx+e , and collect repeated factors so that $g(x)$ is a product of *different* factors of the form $(ax+b)^n$ or $(cx^2+dx+e)^m$ for a nonnegative n .
- (iii). Find real coefficients A_{ij} , B_{ij} , C_{ij} , D_{ij} such that

$$\frac{f(x)}{g(x)} = \sum_{k=1}^n \left(\frac{A_{1k}}{a_kx+b_k} + \frac{A_{n_kk}}{(a_kx+b_k)^2} + \cdots + \frac{A_{n_kk}}{(a_kx+b_k)^n} \right)$$

$$+ \sum_{k=1}^n \left(\frac{C_{1k}x+D_{1k}}{c_kx^2+d_kx+e_k} + \frac{C_{2k}x+D_{2k}}{(c_kx^2+d_kx+e_k)^2} + \cdots + \frac{C_{1k}x+D_{1k}}{(c_kx^2+d_kx+e_k)^n} \right).$$

Exercise 8.4.1 Evaluate integrals:

$$\int \frac{11x+2}{2x^2-5x-3} dx \quad \int \frac{4x}{(x^2+1)^3} dx$$

$$\int \frac{x^4+2x^2+3}{x^3-4x} dx$$

8.5 Quadratic Expressions and Miscellaneous Substitutions

There are a lot of different substitutions which could be useful in particular cases. Particularly, if the integrand is a rational expression in $\sin x$, $\cos x$, the following substitution will produce a rational expression in u :

$$\sin x = \frac{2u}{1+u^2}, \quad \cos x = \frac{1-u^2}{1+u^2}, \quad dx = \frac{2}{1+u^2}du,$$

where $u = \tan \frac{x}{2}$ for $-\pi < x < \pi$.

Exercise 8.5.1 Evaluate integrals

$$\int \frac{1}{\sqrt{7+6x-x^2}} dx; \quad \int \frac{1}{(x^2-6x+34)^{3/2}} dx;$$

$$\int \frac{1}{x(\ln^2 x + 3 \ln x + 2)} dx.$$

8.6 Improper Integrals

We could extend the notion of integral for the following *integrals with infinite limits* or *improper integral*

Definition 8.6.1 (i). If f is continuous on $[a, \infty)$, then

$$\int_a^\infty f(x) dx = \lim_{t \rightarrow \infty} \int_a^t f(x) dx,$$

provided the limit exists.

(ii). If f is continuous on $(-\infty, a]$, then

$$\int_{-\infty}^a f(x) dx = \lim_{t \rightarrow -\infty} \int_t^a f(x) dx,$$

provided the limit exists.

(iii). Let f be continuous for every x . If a is any real number, then

$$\int_{-\infty}^\infty f(x) dx = \int_{-\infty}^a f(x) dx + \int_a^\infty f(x) dx,$$

provided both of the improper integrals on the right converge.

Exercise 8.6.2 Determine if improper integrals converge and find the its value if so.

$$\int_0^\infty x e^{-x} dx; \quad \int_{-\infty}^\infty \cos^2 x dx;$$

$$\int_1^\infty \frac{x}{(1+x^2)^2} dx.$$

Definition 8.6.3 (i). If f is discontinuous on $[a, b)$ and discontinuous at b , then

$$\int_a^b f(x) dx = \lim_{t \rightarrow b^-} \int_a^t f(x) dx,$$

provided the limit exists.

(ii). If f is discontinuous on $(a, b]$ and discontinuous at a , then

$$\int_a^b f(x) dx = \lim_{t \rightarrow a^+} \int_t^b f(x) dx,$$

provided the limit exists.

(iii). If f has a discontinuity at c in the open interval (a, b) but is continuous elsewhere on $[a, b]$, then

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx,$$

provided both of the improper integrals on the right converge.

Exercise 8.6.4 Determine if improper integrals converge and find the its value if so.

$$\begin{aligned} \int_0^4 \frac{1}{(4-x)^{2/3}} dx; & \quad \int_1^2 \frac{x}{x^2-1} dx; \\ \int_0^{\pi/2} \tan x dx; & \quad \int_{1/e}^e \frac{1}{x(\ln x)^2} dx. \end{aligned}$$

Chapter 9

Infinite Series

9.1 Sequences

Definition 9.1.1 A *sequence* is a function f whose domain is the set of positive integers.

Example 9.1.2 (i). Sequence of even numbers: $a_1 = 2, a_2 = 4, a_3 = 6, \dots$

(ii). Sequence of prime numbers: $a_1 = 2, a_2 = 3, a_3 = 5, \dots$

Definition 9.1.3 A sequence $\{a_n\}$ has the *limit* L , or *converges to* L , denoted by

$$\lim_{n \rightarrow \infty} a_n = L, \quad \text{or} \quad a_n \rightarrow L \text{ when } n \rightarrow \infty.$$

if for every $\epsilon > 0$ there exists a positive number N such that

$$|a_n - L| < \epsilon \quad \text{whenever} \quad n > N.$$

If such a number L does not exist, the sequence *has no limit*, or *diverges*.

Definition 9.1.4 The notation

$$\lim_{n \rightarrow \infty} a_n = \infty, \quad \text{or} \quad a_n \rightarrow \infty \text{ or } n \rightarrow \infty.$$

means that for every positive real number P there exists a positive number N such that $a_n > P$ whenever $n > N$.

Theorem 9.1.5 Let $\{a_n\}$ be a sequence, let $f(n) = a_n$, and suppose that $f(x)$ exists for all real numbers $x > 1$.

- (i). If $\lim_{x \rightarrow \infty} f(x) = L$, then $\lim_{n \rightarrow \infty} a_n = L$.
- (ii). If $\lim_{x \rightarrow \infty} f(x) = \infty$, then $\lim_{n \rightarrow \infty} a_n = \infty$.

Theorem 9.1.6

$$\begin{aligned} \lim_{n \rightarrow \infty} r^n &= 0 & \text{if } |r| < 1 \\ \lim_{n \rightarrow \infty} r^n &= \infty & \text{if } |r| > 1 \end{aligned}$$

Exercise 9.1.7 Check if the sequences are convergent

$$\left\{ \frac{n^2}{\ln n + 1} \right\}; \quad \left\{ \frac{\cos n}{n} \right\}; \quad \left\{ \frac{e^n}{n^4} \right\};$$

Theorem 9.1.8 (The Squeeze Rule, or theorem about two policemen)

Suppose that (a_n) and (b_n) are two sequences which tend to the same limit l as $n \rightarrow \infty$. Suppose that (c_n) is another sequence such that there exists $n_0 \in \mathbb{N}$ such that $a_n \leq c_n \leq b_n$ for each $n \geq n_0$. Then $c_n \rightarrow l$ as $n \rightarrow \infty$.

Exercise 9.1.9 Show that $c_n = \sin(n^2)/n$ converge to 0.

9.2 Convergent or Divergent Series

Definition 9.2.1 An *infinite series* (or *series*) is an expression of the form

$$\sum_1^{\infty} a_n = a_1 + a_2 + a_3 + \dots$$

Here a_n is *n*th *term* of the series.

Definition 9.2.2 (i). The *k*th *partial sum* of the series is $S_k = \sum_{n=1}^k a_n$.

(ii). The *sequence of partial sums* of the series is S_1, S_2, S_3, \dots

Definition 9.2.3 A series is *convergent* or *divergent* iff the sequence of partial sums is correspondingly convergent or divergent. If limit of partial sum exists then it is the *sum* of series. A divergent series has no sum.

Example 9.2.4 (i). Series $\sum \frac{1}{n(n+1)}$ is convergent with sum 1.

(ii). Series $\sum (-1)^k$ is divergent.

(iii). The *harmonic series* $\sum \frac{1}{n}$ is divergent.

- (iv). The *geometric series* $\sum ar^n$ is convergent with sum $\frac{a}{1-r}$ if $|r| < 1$ and divergent otherwise.

Theorem 9.2.5 *If a series $\sum a_n$ is convergent, then $\lim_{x \rightarrow \infty} a_n = 0$.*

Exercise 9.2.6 Determine whether the series converges or diverges

$$\begin{aligned} & \sum (\sqrt{2})^{n-1}; & \sum (\sqrt{3})^{1-n}; \\ & \sum \frac{-1}{(n+1)(n+2)}; & \sum (n+3)^{-1}. \end{aligned}$$

9.3 Positive-Term Series

We will investigate first *positive-term series*—that is, series $\sum a_n$ such that $a_n > 0$ for all n —and will use these result for series of general type.

Theorem 9.3.1 *If $\sum a_n$ is a positive-term series and if there exists a number M such that*

$$S_n = a_1 + a_2 + \dots + a_n < M$$

for every n , then the series converges and has a sum $S \leq M$. If no such M exists, the series diverges.

Theorem 9.3.2 *If $\sum a_n$ is a series, let $f(n) = a_n$ and let f be the function obtained by replacing n with x . If f is positive-valued, continuous, and decreasing for every real number $x \geq 1$, then the series $\sum a_n$*

- (i). *converges if $\int_1^\infty f(x) dx$ converges;*
- (ii). *diverges if $\int_1^\infty f(x) dx$ diverges.*

Definition 9.3.3 A *p-series*, or a *hyperharmonic series*, is a series of the form

$$\sum_{n=1}^{\infty} \frac{1}{n^p} = 1 + \frac{1}{2^p} + \frac{1}{3^p} + \dots + \frac{1}{n^p} + \dots,$$

where p is a positive real number.

Theorem 9.3.4 *The p-series $\sum \frac{1}{n^p}$*

- (i). *converges if $p > 1$;*
- (ii). *diverges if $p \leq 1$.*

PROOF. The proof is a direct application of the Theorem 9.3.2. \square

Theorem 9.3.5 (Basic Comparison Tests) Let $\sum a_n$ and $\sum b_n$ be positive-term series.

- (i). If $\sum b_n$ converges and $a_n \leq b_n$ for every positive integer n , then $\sum a_n$ converges.
- (ii). If $\sum b_n$ diverges and $a_n \geq b_n$ for every positive integer n , then $\sum a_n$ diverges.

Theorem 9.3.6 (Limit Comparison Test) Let $\sum a_n$ and $\sum b_n$ be positive-term series. If there is a positive number c such that

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = c > 0,$$

then either both series converge or both series diverge.

Exercise 9.3.7 Determine convergency

$$\begin{array}{ll} \sum \frac{\ln n}{n}; & \sum \frac{1}{1 + 16n^2}; \\ \sum \sin n^4 e^{-n^5}; & \sum \frac{1}{n^n}; \\ \sum \frac{3n + 5}{n2^n}; & \sum \frac{n^2}{n^3 + 1}; \\ \sum \tan \frac{1}{n}; & \sum \frac{\sin n + 2^n}{n + 5^n}. \end{array}$$

9.4 The Ratio and Root Tests

The following two test of divergency are very important. Yet there several cases then they are not inconclusive (see the third clause).

Theorem 9.4.1 (Ratio Test) Let $\sum a_n$ be a positive-term series, and suppose that

$$\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = L.$$

- (i). If $L < 1$, the series convergent.
- (ii). If $L > 1$ the series divergent.

(iii). If $L = 1$, apply a different test; the series may be convergent or divergent..

Exercise 9.4.2 Determine convergency

$$\sum \frac{100^n}{n!}; \quad \sum \frac{3n}{\sqrt{n^3 + 1}}.$$

Theorem 9.4.3 (Root Test) Let $\sum a_n$ be a positive-term series, and suppose that

$$\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = L.$$

(i). If $L < 1$, the series convergent.

(ii). If $L > 1$ the series divergent.

(iii). If $L = 1$, apply a different test; the series may be convergent or divergent..

Exercise 9.4.4 Determine convergency

$$\begin{aligned} \sum \frac{2^n}{n^2}; & \quad \sum \left(\frac{n}{\ln n}\right)^n; \\ \sum \frac{n!}{n^n}; & \quad \sum \frac{1}{(\ln n)^n}. \end{aligned}$$

9.5 Alternating Series and Absolute Convergence

The simplest but still important case of non positive-term series are *alternating series*, in which the terms are alternately positive and negative..

Theorem 9.5.1 (Alternating Series Test) The alternating series

$$\sum_{n=1}^{\infty} (-1)^{n-1} a_n = a_1 - a_2 + a_3 - a_4 + \cdots + (-1)^{n-1} a_n + \cdots$$

is convergent if the following two conditions are satisfied:

(i). $a_n \geq a_{k+1} \geq 0$ for every k ;

(ii). $\lim a_n = 0$.

Theorem 9.5.2 Let $\sum (-1)^{n-1} a_n$ be an alternating series that satisfies conditions (i) and (ii) of the alternating series test. If S is the sum of the series and S_n is a partial sum, then

$$|S - S_n| \leq a_{n+1};$$

that is, the error involved in approximating S by S_n is less than or equal to a_{n+1} .

Definition 9.5.3 A series $\sum a_n$ is *absolutely convergent* if the series

$$\sum |a_n| = |a_1| + |a_2| + \cdots + |a_n| + \cdots$$

is convergent.

Definition 9.5.4 A series $\sum a_n$ is *conditionally convergent* if $\sum a_n$ is convergent and $\sum |a_n|$ is divergent.

Theorem 9.5.5 If a series $\sum a_n$ is absolutely convergent, then $\sum a_n$ is convergent.

Exercise 9.5.6 Determine convergency

$$\begin{array}{ll} \sum \frac{(-1)^n 2}{n^2 + n}; & \sum (-1)^n \frac{\sqrt[3]{n}}{n+1}; \\ \sum (-1)^n \frac{\arctan n}{n^2}; & \sum \frac{1}{n} \sin \frac{(2n-1)\pi}{2}. \end{array}$$

Appendix A

Algebra

We briefly recall some basic notion and results from algebra.

A.1 Numbers

The following set of *numbers* will be used in the course

- (i). \mathbb{N} —*natural* numbers: 1, 2, 3, ...
- (ii). \mathbb{Z} —*integer* numbers: ..., -2, -1, 0, 1, 2, ...
- (iii). \mathbb{Q} —*rational* numbers of the form $\frac{n}{m}$, $n \in \mathbb{Z}$, $m \in \mathbb{N}$.
- (iv). \mathbb{R} —*real* numbers, e.g, 0, 1, -3.12, $\sqrt{2}$, π , e .

Binary operations between numbers are +, -, ·, /.

Exercise A.1.1 Which set of numbers is closed with respect to the above four operations?

A.2 Polynomial. Factorization of Polynomials

A *polynomial* $p(x)$ (in a variable x) is a function on real line defined by an expression of the form:

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0. \quad (\text{A.2.1})$$

Here a_i are fixed real numbers, $a_n \neq 0$ and n is the degree of polynomial $p(x)$.

According to the Main Theorem of algebra every polynomial $p(x)$ could be represented as a product of linear binomials and quadratic terms as follows:

$$p(x) = (b_1x + c_1) \cdots (b_jx + c_j)(d_1x^2 + f_1x + g_1) \cdots (d_kx^2 + f_kx^2 + g + k),$$

moreover $n = 2k + j$, where n is the degree of $p(x)$.

Exercise A.2.1 Decompose to products:

(i). $p(x) = x^{16} - 1$.

(ii). $p(x) = x^4 + 16y^4$.

A.3 Binomial Formula

Binomial formula of Newton:

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} x^k y^{n-k}, \quad \text{where} \quad \binom{n}{k} = \frac{n!}{k!(n-k)!}.$$

Here $n! = 1 \cdot 2 \cdots n$. These coefficients can be determined from the *Pascal triangle*.

Exercise* A.3.1 Prove the following properties of the binomial coefficients

$$\begin{aligned} \binom{n+1}{k} &= \binom{n}{k} + \binom{n}{k+1}. \\ \binom{n}{k+1} &= \frac{n-k}{k+1} \binom{n}{k}. \\ \sum_{k=0}^n \binom{n}{k} &= 2^n. \\ \sum_{k=0}^n (-1)^k \binom{n}{k} &= 0. \end{aligned}$$

A.4 Real Axis

We will be mainly interested in real numbers \mathbb{R} which could be represented by the *coordinate line* (or *real axis*). This gives *one-to-one* correspondence between sets of real numbers and points of the real line.

A.5 Absolute Value

The *absolute value* $|a|$ (or *modulus*) of a real number a is defined as follows

$$|a| = \begin{cases} a & \text{if } a \geq 0, \\ -a & \text{if } a < 0. \end{cases}$$

It has the following properties (for $b > 0$)

- (i). $|a| < b$ iff¹ $-b < a < b$.
- (ii). $|a| > b$ iff either $a > b$ or $a < -b$.
- (iii). $|a| = b$ iff $a = b$ or $a = -b$.

Exercise A.5.1 Prove the following properties of absolute value:

- (i). $|a + b| \leq |a| + |b|$.
- (ii). $|ab| = |a| |b|$.

An *equation* is an equality that involves variables, e.g. $x^3 + 5x^2 - x + 10 = 0$. A *solution* of an equation (or *root* of an equation) is a number b that produces a true statement after substitution $x = b$ into equation. Equation could be solved by either *analytic* or *computational* means.

A.6 Inequalities

Order relations between numbers are given by $>$, $<$, \leq , \geq , $=$.

They have the following properties:

- (i). If $a > b$ and $b > c$, then $a > c$ (*transitivity property*).
- (ii). If $a > b$, then $a \pm c > b \pm c$.
- (iii). If $a > b$ and $c > 0$, then $ac > bc$.
- (iv). If $a > b$ and $c < 0$, then $ac < bc$.

An *inequality* is a statement involves variables and at least one of symbols $>$, $<$, \leq , \geq , e.g. $x^3 > 2x^2 - 5x + 1$. *Solution* of an inequality is similar for the case of equations (see Section A.5). They are often given by unions of intervals.

¹The notation *iff* is used for an abbreviation of *if and only if*.

Intervals on real line are the following sets:

$$\begin{aligned}[a, b] &= \{x \in \mathbb{R} \mid a \leq x \leq b\}; \\(a, b) &= \{x \in \mathbb{R} \mid a < x < b\}; \\(a, b] &= \{x \in \mathbb{R} \mid a < x \leq b\}; \\[a, b) &= \{x \in \mathbb{R} \mid a \leq x < b\}.\end{aligned}$$

Particularly a can be $-\infty$ and $b = \infty$.

Appendix B

Function and Their Graph

B.1 Rectangular (Cartesian) Coordinates

Considering real axis in Section A.4 we introduce one-to one correspondence between real numbers and points of a line. This connection between numbers and geometric objects may be extended for other objects as well.

A *rectangular coordinate system* (or *Cartesian coordinates*) is an assignment of *ordered pairs* (a, b) to points in a plane, see [1, Fig. 6, p. 10].

REMARK B.1.1 It is also possible to introduce Cartesian coordinates in our three-dimensional world by means of triples of real numbers (x, y, z) . This construction could be extended to arbitrary number of dimensions.

Theorem B.1.2 *The distance between two points $P_1 = (x_1, y_1)$ and $P_2 = (x_2, y_2)$ is*

$$d(P_1, P_2) = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}. \quad (\text{B.1.1})$$

This theorem is a direct consequence of Pythagorean theorem.

Theorem B.1.3 *The midpoint M of segment P_1P_2 is*

$$M(P_1P_2) = M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right). \quad (\text{B.1.2})$$

PROOF. The theorem follows from two observations:

- (i). $d(P_1, M) = d(M, P_2)$;
- (ii). $d(P_1, M) + d(M, P_2) = d(P_1, P_2)$.

□

Exercise B.1.4 Prove the above two observation (Hint: use the distance formula (B.1.1)).

B.2 Graph of an Equation

An *equation in x and y* is an equality such as

$$2x + 3y = 5, \quad y = x^2 + 3x - 6, \quad y^x + \sin xy = 8.$$

A *solution* is an ordered pair (a, b) that produced a true statement when $x = a$ and $y = b$. The *graph of the equation* consists of all points (a, b) in a plane that corresponds to the solutions.

B.3 Line Equations

The general *equation* of a line in a plane is given by the formula

$$ax + by + c = 0. \tag{B.3.1}$$

This equation connect different geometric objects:

(i). *Slope m* :

$$m = \frac{y_2 - y_1}{x_2 - x_1}. \tag{B.3.2}$$

(ii). *Point-Slope form* $y - y_1 = m(x - x_1)$.

(iii). *Slope-Intercept form* $y = mx + b$ or $y = m(x - c)$.

Special lines

(i). Vertical: m undefined; horizontal: $m = 0$.

(ii). Parallel: $m_1 = m_2$.

(iii). Perpendicular $m_1 m_2 = -1$.

Exercise B.3.1 Prove the above geometric properties.

B.4 Symmetries and Shifts

We will say that a graph of an equation possesses a *symmetry* if there is a transformation of a plane such that it maps the graph to itself.

Example B.4.1 There several examples of elementary symmetries:

- (i). y -axis: substitution $x \rightarrow (-x)$, e.g.¹ equation $y = |x|$.
- (ii). x -axis: substitution $y \rightarrow (-y)$, e.g. $|y| = x$.
- (iii). Central symmetry: substitution both $x \rightarrow (-x)$ and $y \rightarrow (-y)$, e.g. $y = x$ or $|y| = |x|$.
- (iv). x -shifts: substitution $x \rightarrow (x + a)$ for $a \neq 0$, e.g. $y = \{x\}$ with $a = 1$. Here $\{x\}$ denotes the *fractional part* of x , i.e.² it is defined by two conditions: $0 \leq \{x\} < 1$ and $x - \{x\} \in \mathbb{Z}$.
- (v). y -shifts: substitution $y \rightarrow (y + b)$ for $b \neq 0$, e.g. $\{y\} = x$ with $b = 1$.
- (vi). General shifts: substitution both $x \rightarrow (x + a)$ and $y \rightarrow (y + b)$ for $a \neq 0$, $b \neq 0$, e.g. $y = [x]$ with $a = b = 1$. Here $[x]$ denotes the *entire part* of x , i.e. $[x] \in \mathbb{Z}$ and $[x] \leq x < [x] + 1$.

- Exercise* B.4.2** (i). Is there an equation with y -axis symmetry B.4.1.(i) and x -axis symmetry B.4.1.(ii) but without central symmetry B.4.1.(iii)?
- (ii). Is there an equation with x -shift symmetry B.4.1.(iv) with some $a \neq 0$ and y -shift symmetry B.4.1.(v) with some $b \neq 0$ but without general shift symmetry B.4.1.(vi)?

Symmetries are important because they allow us to reconstruct a whole picture from its parts.

B.5 Definition of a Function. Domain and Range

The main object of calculus is *function*. We recall basic notations and definitions.

Definition B.5.1 A *function* f from a set D to a set E is a correspondence that assigns to each element x of the set D exactly one element y of the set E .

The element y of E is the *value* of f at x , notation— $f(x)$. The set D is the *domain* of the function f , E —*codomain* of f . The *range* of f is the subset of codomain E consisting of all possible function values $f(x)$ for x in D . Here x is *independent variable* and y is *dependent variable*.

¹The abbreviation *e.g.* denotes *for example*.

²The abbreviation *i.e.* denotes *namely*.

B.6 One-to-One Functions. Periodic Functions

We say that f is *one-to-one* function if $f(x) \neq f(y)$ whenever $x \neq y$. For numerically defined functions like $y = \sqrt{x-2}$ the domain is assumed to be all x that f is *is defined at* x , or $f(x)$ exists.

The *graph of the a function* f with domain D is the graph of the equation $y = f(x)$ for x in D . The x -intercept of the graph are solutions of the equation $f(x) = 0$ and called *zeros*.

The following transformation are useful for sketch of graphs:

- (i). *horizontal shift*: $y = f(x)$ to $y = f(x - a)$;
- (ii). *vertical shift*: $y = f(x)$ to $y = f(x) + b$;
- (iii). *horizontal stretch/compression*: $y = f(x)$ to $y = f(cx)$;
- (iv). *vertical stretch/compression*: $y = f(x)$ to $y = cf(x)$;
- (v). *horizontal reflections*: $y = f(x)$ to $y = -f(x)$;
- (vi). *vertical reflections*: $y = f(x)$ to $y = f(-x)$;

B.7 Increasing and Decreasing Functions. Odd and Even Functions

A function $f(x)$ is *increasing* if $f(x) > f(y)$ for all $x > y$. A function $f(x)$ is *decreasing* if $f(x) < f(y)$ for all $x > y$.

A function $f(x)$ is *even* if $f(-x) = f(x)$ and $f(x)$ is *odd* if $f(-x) = -f(x)$.

Exercise B.7.1 Which type of symmetries listed in Example (B.4.1) have to or may even and odd functions possess?

There are natural operation on functions

- (i). *sum*: $(f + g)(x) = f(x) + g(x)$.
- (ii). *difference*: $(f - g)(x) = f(x) - g(x)$.
- (iii). *product*: $(fg)(x) = f(x)g(x)$.
- (iv). *quotient*: $(f/g)(x) = f(x)/g(x)$.

For $p(x)$ be a polynomial (see (A.2.1)) $y = p(x)$ defines a *polynomial function*. If $p(x)$ and $q(x)$ are two polynomials (see (A.2.1)) then $y = p(x)/q(x)$ is a *rational function*. Function which are obtained from polynomials by four algebraic operations B.7.1.(i) and taking rational powers are *algebraic*. All other function (e.g $\sin x$, $\exp x$) are *trancendental*.

The *composite function* $f \circ g$ is defined by $(f \circ g)(x) = f(g(x))$. An *identity function* is a function h with the property that $h(x) = x$ If the composition of two functions f and g is an identity function, then the functions are *inverses* of each other.

Appendix C

Conic Section

C.1 Circle

The beautiful and important objects arise by intersection of planes and cones, i.e. *conic sections*.

The simplest conic section is circle.

Definition C.1.1 A *circle* with center (x_1, y_1) and radius r consists of points on the distance r from (x_1, y_1) .

By the distance formula (B.1.1) the circle is defined by an *equation*:

$$(x - x_1)^2 + (y - y_1)^2 = r^2. \quad (\text{C.1.1})$$

Circles are obtained as intersections of cones with planes orthogonal to their axes.

Exercise C.1.2 (i). Write an equation of a circle which is tangent to a circle $x^2 - 6x + y^2 + 4y - 12 = 0$ and has the origin $(3, 0)$.

(ii). Write an equation of a circle, which has a center at $(1, 2)$ and contains the center of the circle given by the equation $x^2 - 7x + y^2 + 8y - 17 = 0$.

(iii). Write equations of all circles with a given radius r which are tangent to both axes.

C.2 Parabola

Definition C.2.1 A *parabola* is the set of all points in a plane equidistant from a fixed point F (the *focus* of the parabola) and a fixed line l (the *directrix*) that lie in the plane.

The *axis* of the parabola is the line through F that is perpendicular to the directrix. The *vertex* of the parabola is the point V on the axis halfway from F to l .

A parabola with axis coinciding with y axis and the vertex at the origin with focus $F = (0, p)$ has an *equation*

$$x^2 = 4py. \quad (\text{C.2.1})$$

Exercise* C.2.2 Verify the above equation of a parabola. (Hint: use distance formula (B.1.1)).

Exercise C.2.3 List all symmetries of a parabola.

For a parabola with the vertex (h, k) and a horizontal directrix an equation takes the form

$$(x - h)^2 = 4p(y - k). \quad (\text{C.2.2})$$

In general any equation of the form $y = ax^2 + bx + c$ defines a parabola with horizontal directrix.

Exercise C.2.4 (i). Find the vertex, the focus, and the directrix of the parabolas:

- (a) $3y^2 = -5x$.
- (b) $x^2 = 3y$.
- (c) $y^2 + 14y + 4x + 45 = 0$.
- (d) $y = 8x^2 + 16x + 10$.

(ii). Find an equation of the parabola with properties:

- (a) vertex $V(-3, 4)$; directrix $y = 6$.
- (b) vertex $V(1, 1)$; focus $F(-2, 1)$.
- (c) focus $F(1, -3)$; directrix $y = 5$.

C.3 Ellipse

Definition C.3.1 An *ellipse* is the set of all points in a plane. the sum of whose distances from two fixed points F and F' (the *foci*) in the plane is constant. The midpoint of the segment FF' is the *center* of the ellipse.

Let $F(-c, 0)$ and $F'(c, 0)$, $2a$ be the constant sum of distances, and $b = (a^2 - c^2)^{1/2}$. Then the ellipse has an *equation*

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \quad (\text{C.3.1})$$

Exercise* C.3.2 Verify the above equation of the ellipse. (Hint: use distance formula (B.1.1).)

The ellipse intercepts x -axis in points $V(-a, 0)$ and $V'(a, 0)$ —*vertices* of the ellipse. The line segment VV' is the *major axis* of the ellipse. Similarly the ellipse intercepts y -axis in points $M(-b, 0)$ and $M'(b, 0)$ and the line segment MM' is the *minor axis* of the ellipse.

Exercise C.3.3 List all symmetries of an ellipse.

For the ellipse with center in a point (h, k) an equation is given as

$$\frac{(x - h)^2}{a^2} + \frac{(y - k)^2}{b^2} = 1. \quad (\text{C.3.2})$$

Definition C.3.4 The *eccentricity* e of an ellipse is

$$e = \frac{c}{a} = \frac{\sqrt{a^2 - b^2}}{a}$$

Exercise C.3.5 (i). Find the vertices and the foci of the ellipse

(a) $4x^2 + 2y^2 = 8$.

(b) $x^2/3 + 3y^2 = 9$

(ii). Find an equation for the ellipse with center at the origin and

(a) Vertices $V(\pm 9, 0)$; foci $F(\pm 6, 0)$.

(b) Foci $F(\pm 6, 0)$; minor axis of length 4.

(c) Eccentricity $3/4$; vertices $V(0, \pm 5)$.

C.4 Hyperbola

Definition C.4.1 A *hyperbola* is the set of all points in a plane, the difference of whose distances from two fixed points F and F' (the *foci*) in the plane is a positive constant. The midpoint of the segment FF' is the *center* of the hyperbola.

Let a hyperbola has foci $F(\pm c, 0)$, $2a$ denote the constant difference, and let $b^2 = c^2 - a^2$. Then the hyperbola has an *equation*

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1. \quad (\text{C.4.1})$$

Exercise* C.4.2 Verify the above equation of the hyperbola. (Hint: use distance formula (B.1.1)).

Points $V(a, 0)$ and $V'(-a, 0)$ of interception of the hyperbola with x -axis are *vertices* and the line segment VV' is the *transverse axis* of the hyperbola. Points $W(0, b)$ and $W'(0, -b)$ span *conjugate axis* of the hyperbola. This two segments intercept in the *center* of the hyperbola.

Exercise C.4.3 Find all symmetries of a hyperbola.

If a graph approaches a line as the absolute value of x gets increasingly large, then the line is called an *asymptote* for the graph. It could be shown that lines $y = (b/a)x$ and $y = -(b/a)x$ are asymptotes for the hyperbola.

For the hyperbola with the center in a point (h, k) an equation is given as

$$\frac{(x - h)^2}{a^2} - \frac{(y - k)^2}{b^2} = 1. \quad (\text{C.4.2})$$

Exercise C.4.4 (i). Find vertices and foci of the hyperbola, sketch its graph.

- (a) $x^2/49 - y^2/16 = 1$.
- (b) $y^2 - 4x^2 - 12y - 16x + 16 = 0$.
- (c) $9y^2 - x^2 - 36y + 12x - 36 = 0$.

(ii). Find an equation for the hyperbola that has its center at the origin and satisfies to the given conditions

- (a) foci $F(0, \pm 4)$; vertices $V(0, \pm 1)$.

- (b) foci $F(0, \pm 5)$; conjugate axis of length 4.
- (c) vertices $V(\pm 3, 0)$; asymptotes $y = \pm 2x$.

C.5 Conclusion

The graph of every quadratic equation $Ax^2 + Cy^2 + Dx + Ey + F = 0$ is one of conic section

- (i). Circle;
- (ii). Ellipse;
- (iii). Parabola;
- (iv). Hyperbola;

or a degenerated case

- (i). A point;
- (ii). Two crossed lines;
- (iii). Two parallel lines;
- (iv). One line;
- (v). The empty set.

Appendix D

Trigonometric Functions

An *angle* is determined by two rays having the same initial point O (*vertex*). Angles are measured either by *degree measure* 1° or *radian measure*. The complete counterclockwise revolution is 360° or 2π radians.

We consider six *trigonometric functions*

Name	Notation	Expression	Name	Notation	Expression
<i>sine</i>	sin	y/r	<i>cosecant</i>	csc	r/y
<i>cosine</i>	cos	x/r	<i>secant</i>	sec	r/x
<i>tangent</i>	tan	y/x	<i>cotangent</i>	cot	x/y

There are a lot of useful identities between trigonometric functions:

(i). *Reciprocal and Ratio Identities*:

(a) $\csc \phi = (\sin \phi)^{-1}$, $\sec \phi = (\cos \phi)^{-1}$;

(b) $\tan \phi = \sin \phi / \cos \phi$, $\cot = \cos \phi / \sin \phi$;

(c) $\cot \phi = (\tan \phi)^{-1}$

(ii). *Pythagorean Identities*

(a) $\sin^2 \phi + \cos^2 \phi = 1$;

(b) $1 + \tan^2 \phi = \sec^2 \phi$;

(c) $1 + \cot^2 \phi = \csc^2 \phi$;

(iii). *Law of Sines and Law of Cosine*

(a) $\sin \alpha / a = \sin \beta / b = \sin \gamma / c = 2R$;

(b) $a^2 = b^2 + c^2 - 2bc \cos \alpha$;

(iv). Additional identities

(a) Cosine and secant are even functions; sine, tangent, cosecant, cotangent are odd functions;

(b) $\sin(\alpha \pm \beta) = \sin \alpha \cos \beta \pm \sin \beta \cos \alpha$;

(c) $\cos(\alpha \pm \beta) = \cos \alpha \cos \beta \mp \sin \alpha \sin \beta$;

Appendix E

Exponential and Logarithmic Functions

Definition E.0.1 The *exponential function* with a base a is defined by $f(x) = a^x$, where $a > 0$, $a \neq 1$, and x is any real number.

It is increasing if $a > 1$ and decreasing if $0 < a < 1$. It is also one-to-one function. This allow us to solve equations and inequalities.

Exercise E.0.2 Find solutions

(i). $5^{3x} = 5^{x^2-1}$;

(ii). $2^{|x-3|} > 2^2$;

(iii). $(0.5)^{x^2} > (0.5)^{5x-6}$.

There are *laws of exponents*:

(i). $a^u a^v = a^{u+v}$;

(ii). $a^u / a^v = a^{u-v}$;

(iii). $(a^u)^v = a^{uv}$;

(iv). $(ab)^u = a^u b^u$;

(v). $(a/b)^u = a^u / b^u$.

Definition E.0.3 If a is a positive real number other than 1, then the *logarithm* of x with base a is defined by $y = \log_a x$ if and only if $x = a^y$ for every $x > 0$ and every real number y .

Thus logarithm is inverse to exponential function. As consequences logarithm one-to-one function, for $a > 1$ it is an increasing function, for $a < 1$ it is decreasing.

Exercise E.0.4 Find solution

(i). $\log_2(x^2 - x) = \log_2 2$;

(ii). $\log_{0.5} |2x - 5| > \log_{0.5} 4$.

There are corresponding *laws of logarithms*

(i). $\log_a(uv) = \log_a u + \log_a v$;

(ii). $\log_a(u/v) = \log_a u - \log_a v$;

(iii). $\log_a(u^c) = c \log_a u$ for any real number c .

The *change-of-base formula for logarithms*: if $x > 0$ and if a and b are positive real numbers other than 1, then

$$\log_b x = \frac{\log_a x}{\log_a b}. \quad (\text{E.0.1})$$

Exercise E.0.5 Find solution: $\log_x(3x - 1) = 2$.

Bibliography

- [1] Earl Swokowski, Michael Olinick, and Dennis Pence. *Calculus*. PWS Publishing, Boston, 6-th edition, 1994.

Index

- A notation, 6
- O notation, 7
- o notation, 8
- o notation, 7
- p -series, 56

- absolute, 20
- absolute value, 62
- absolutely convergent, 59
- algebraic, 68
- alternating series, 58
- analytic, 62
- angle, 74
 - vertex of, 74
- antiderivative, 27
- arbitrary constant, 27
- arc length function, 37
- arccosine, 44
- arcsecant, 44
- arcsine, 44
- arctangent, 44
- asymptote, 72
- average value, 31
- average velocity, 14
- axis, 70

- Binary operations, 60
- binary operations, 60
- Binomial formula, 61
- binomial formula, 61

- Cartesian coordinates, 64
- center, 70, 72
- chain rule, 15, 17

- change-of-base formula for logarithms, 77
- circle, 69
 - equation of, 69
- codomain, 66
- composite function, 68
- computational, 62
- concave downward, 24
- concave upward, 24
- conditionally convergent, 59
- conic sections, 69
- conjugate axis, 72
- continuity, 11
- continuous, 11
- convergent, 55
- coordinate line, 61
- coordinate system
 - Cartesian, 64
 - rectangular, 64
- cosecant, 74
- cosine, 74
- cotangent, 74
- Critical numbers, 21
- critical numbers, 21
- cross section, 36

- decreasing, 20, 67
- definite integral, 30
- degree measure, 74
- dependent variable, 66
- derivative, 15
- discontinuity
 - jump, 11
- difference, 67

- differential, 19
- Differential equation, 28
- differentiation
 - implicit, 18
 - logarithmic, 40
- directrix, 69
- discontinuity, 11
 - infinite, 11
 - removable, 11
- discontinuous, 11
- distance between two points, 64
- divergent, 55
- domain, 66
- e.g., 66
- eccentricity, 71
- ellipse
 - vertices of, 71
- ellipse, 70
 - center of, 70
 - eccentricity of, 71
 - equation, 71
 - foci of, 70
 - major axis of, 71
 - minor axis of, 71
- entire part, 66
- equation, 62, 65, 69–72
 - in x and y , 65
 - differential, 28
 - graph of the, 65
 - of hyperbola, 72
 - of line, 65
 - of circle, 69
 - of parabola, 70
 - of tangent line, 14
 - solution, 62
 - analytic, 62
 - computational, 62
- equation in x and y , 65
- equation of tangent line, 14
- even, 67
- explicit function, 17
- exponential function, 76
- exponential function with base a , 42
- extrema, 20
- extreme values, 20
- foci, 70, 72
- focus, 69
- formula
 - Cauchy's, 47
 - change-of-base for logarithms, 77
- fractional part, 66
- function
 - value of, 66
- function, 66
 - algebraic, 68
 - codomain of, 66
 - composite, 68
 - continuous, 11
 - closed interval, 12
 - open interval, 12
 - decreasing, 67
 - difference of, 67
 - discontinuous, 11
 - domain of, 66
 - even, 67
 - explicit, 17
 - graph of, 67
 - implicit, 18
 - increasing, 67
 - integrable, 30
 - inverse, 38
 - inverses, 68
 - is defined at x , 67
 - odd, 67
 - one-to-one, 67
 - polynomial, 68
 - product of, 67
 - quotient of, 67

- range of, 66
- rational, 68
- smooth, 37
- sum of, 67
- zeros, 67
- functions
 - transcendental, 38, 68
- geometric series, 56
- global, 20
- graph
 - function, 67
- graph of the a function, 67
- graph of the equation, 65
- harmonic series, 55
- horizontal reflections, 67
- horizontal shift, 67
- horizontal stretch/compression, 67
- hyperbola, 72
 - center of, 72
 - conjugate axis of, 72
 - equation of, 72
 - foci of, 72
 - transverse axis of, 72
 - vertices, 72
- hyperbolic cosine function, 45
- hyperbolic cotangent function, 45
- hyperbolic sine function, 45
- hyperbolic tangent function, 45
- hyperharmonic series, 56
- i.e., 66
- identity function, 68
- iff, 62
- implicit differentiation, 18
- implicit function, 18
- improper integral, 52
- increasing, 20, 67
- indefinite integral, 27
- independent variable, 66
- inequality, 62
 - solution, 62
- Infinite, 11
- infinite series, 55
- initial conditions, 28
- instantaneous velocity, 14
- integer, 60
- integrable function, 30
- integral
 - definite, 30
 - indefinite, 27
- integrals with with infinite limits, 52
- interval, 12
- Intervals, 62
- intervals, 63
- inverse cosine function, 44
- inverse function, 38
- inverse secant function, 44
- inverse sine function, 44
- inverse tangent function, 44
- inverses, 68
- is defined at x , 67
- Jump, 11
- law
 - cosines, 74
 - exponents, 76
 - logarithm, 77
 - sines, 74
- Law of Cosine, 74
- Law of Sines, 74
- laws of exponents, 76
- laws of logarithms, 77
- limit, 8
 - of function, 8
 - one-sided, 10
- limit of Riemann sums, 30
- line
 - equation of, 65
 - point-slope form, 65

- slope-intercept form, 65
 - slope of, 65
 - tangent
 - slope, 14
- linear approximation, 18
- local maximum, 20
- local minimum, 20
- logarithm, 76
- logarithmic differentiation, 40
- logarithmic function with base a , 43
- major axis, 71
- maximum, 20
 - absolute, 20
 - global, 20
 - local, 20
- midpoint, 64
- minimum, 20
 - absolute, 20
 - global, 20
 - local, 20
- minor axis, 71
- modulus, 62
- natural, 60
- natural exponential function, 41
- natural logarithm function, 39
- numbers, 60
 - integer, 60
 - natural, 60
 - rational, 60
 - real, 60
- odd, 67
- one-sided, 10
- one-to-one, 67
- Order relations, 62
- order relations, 62
- parabola, 69
 - axis of, 70
 - directrix of, 69
 - equation of, 70
 - focus of, 69
 - vertex of, 70
- partial sum, 55
- Pascal triangle, 61
- point of inflection, 24
- Point-Slope form, 65
- polynomial, 60
- polynomial function, 68
- positive-term series, 56
- product, 67
- Pythagorean Identities, 74
- quotient, 67
- radian measure, 74
- range, 66
- rational, 60
- rational function, 68
- real, 60
- real axis, 61
- Reciprocal and Ratio Identities, 74
- rectangular coordinate system, 64
- related rates, 18
- Removable, 11
- Riemann sum, 30
- root, 62
- rule
 - l'Hospital's, 47
 - squeeze , 55
- secant, 74
- sequence, 54
- sequence of partial sums, 55
- series, 55
 - absolutely convergent, 59
 - alternating, 58
 - conditionally convergent, 59
 - convergent, 55
 - divergent, 55
 - hyperharmonic, 56

- positive-term, 56
- sum of, 55
- term of, 55
- sine, 74
- Slope, 65
- slope
 - of tangent line, 14
- slope of a line, 65
- Slope-Intercept form, 65
- smooth, 37
- Solution, 62
- solution, 62
- squeeze rule, 55
- sum, 55, 67
- summation notation, 29
- symmetry, 65
- tangent, 74
- tangent line
 - equation, 14
- term, 55
- test
 - alternating series, 58
 - Basic Comparison, 57
 - first derivative, 23
 - Integral, 56
 - Limit Comparison, 57
 - Ratio, 57
 - Root, 58
- test value, 23
- theorem
 - about two policemen, 55
 - intermediate value, 13
 - Lagrange's, 22
 - mean value, 22
 - Rolle's, 22
 - sandwich, 10
- trancendental, 68
- transcendental, 38
- transitivity property, 62
- transverse axis, 72
- trigonometric
 - Pythagorean identities, 74
 - reciprocal and ratio identities, 74
 - trigonometric functions, 74
- trigonometric functions, 74
- two policemen
 - theorem, 55
- value, 66
- variable
 - dependent, 66
 - independent, 66
- velocity
 - average, 14
 - instantaneous, 14
- vertex, 70, 74
- vertical reflections, 67
- vertical shift, 67
- vertical stretch/compression, 67
- vertices, 71, 72
- zeros, 67