

FROM TRIANGULATED CATEGORIES TO MODULE CATEGORIES VIA LOCALISATION

ASLAK BAKKE BUAN AND ROBERT J. MARSH

ABSTRACT. We show that the category of finite-dimensional modules over the endomorphism algebra of a rigid object in a Hom-finite triangulated category is equivalent to the Gabriel-Zisman localisation of the category with respect to a certain class of maps. This generalises the 2-Calabi-Yau tilting theorem of Keller-Reiten, in which the module category is obtained as a factor category, to the rigid case.

INTRODUCTION

Localisation is an important tool in category theory. At a fundamental level, it is the introduction of formal inverses for a class of morphisms in the category known as Gabriel-Zisman localisation [10, Chap. 1]. This category always exists provided there are no set-theoretic obstructions. Morphisms in the new category can be regarded as compositions of the original morphisms and the formal inverses that were added (up to a certain equivalence relation). If the class of morphisms to be inverted satisfies certain axioms then the new morphisms can be described via a calculus of fractions: in particular, each new morphism can be written as the composition of one of the original morphisms and a formal inverse, often represented as a diagram of morphisms known as a roof; see [18, Sect. 3]. This is, for example, the case when the bounded derived category of a module category is formed from the homotopy category of complexes (see e.g. [11, III.2-4]). In this case the localisation of a triangulated category gives us a new triangulated category, which is often the case in applications of localisation to triangulated categories; see [18].

Here, we present an interesting case where localisation produces instead an abelian category, in fact the module category over the (opposite) endomorphism algebra of a rigid object in a triangulated category satisfying some mild assumptions. Furthermore, the class of morphisms inverted does not satisfy the axioms mentioned above.

We consider the tilting theory of cluster categories and, more generally, Hom-finite Calabi-Yau triangulated categories, which has recently been widely investigated.

Date: 22 February 2011.

2010 Mathematics Subject Classification. Primary 18E30, 18E35, 16G20; Secondary 13F60.

Key words and phrases. Triangulated categories, 2-Calabi-Yau categories, localisation, module categories, rigid objects, cluster-tilting objects, approximations.

This work was supported by the Engineering and Physical Sciences Research Council [grant number EP/G007497/1] and by the NFR [FRINAT grant number 196600].

The study of such categories was originally motivated by their links to cluster algebras, and indeed there has been a considerable amount of activity and a lot of results in this direction; see [15, 20] for recent surveys.

However, the study of such categories has also contributed to new developments in the theory of finite dimensional (and more generally, non-commutative) algebras. One of the central theorems in this direction is the following:

2-Calabi-Yau tilting theorem (Keller-Reiten [16, Prop. 2.1]).

Let \mathcal{C} be a triangulated Hom-finite Krull-Schmidt 2-Calabi-Yau category over an algebraically-closed field k , and let T be a cluster-tilting object in \mathcal{C} . Then the category $\mathcal{C}/\Sigma T$ is equivalent to the category $\text{mod End}_{\mathcal{C}}(T)^{op}$ of finite dimensional $\text{End}_{\mathcal{C}}(T)^{op}$ -modules.

Here Σ denotes the suspension functor of \mathcal{C} . Recall that a *cluster-tilting object* in \mathcal{C} is an object satisfying $\text{Ext}_{\mathcal{C}}^1(T, X) = 0$ for an object X of \mathcal{C} if and only if X is in the additive closure of T . Here, for objects C_1 and C_2 in \mathcal{C} , $\text{Ext}_{\mathcal{C}}^1(C_1, C_2)$ is shorthand for $\text{Hom}_{\mathcal{C}}(C_1, \Sigma C_2)$, following the usual convention.

The category $\mathcal{C}/\Sigma T$ has the same objects as \mathcal{C} , with maps given by maps in \mathcal{C} modulo maps factoring through $\text{add } \Sigma T$ (the additive closure of ΣT). This generalises a corresponding result of [8, Thm. 2.2] in the case of cluster categories.

It turns out that the assumption that \mathcal{C} is 2-Calabi-Yau is not required for this theorem. This is a result of Koenig-Zhu [17, Cor. 4.4]. See also [13, Prop. 6.2].

Our aim in this paper is to use localisation to generalise this result to the case where T is only assumed to be a rigid object, i.e. an object satisfying $\text{Ext}_{\mathcal{C}}^1(T, T) = 0$. We consider a triangulated Hom-finite Krull-Schmidt triangulated category \mathcal{C} , over a field k . In addition, we assume that \mathcal{C} is skeletally small (as a way of ensuring that there are no set-theoretic obstructions to the localisations we use). Note that this condition is satisfied for cluster categories. We assume that T is a rigid object in \mathcal{C} . We show that with these assumptions, $\text{mod End}_{\mathcal{C}}(T)^{op}$ can be obtained as the Gabriel-Zisman localisation of \mathcal{C} with respect to a suitable class of maps in \mathcal{C} .

More precisely, let \mathcal{X}_T denote the full subcategory of \mathcal{C} whose objects are the objects X of \mathcal{C} having no non-zero maps from T . Consider the class of maps $\tilde{\mathcal{S}}$ in \mathcal{C} consisting of the maps $f: X \rightarrow Y$ such that when f is completed to a triangle

$$\Sigma^{-1}Z \xrightarrow{h} X \xrightarrow{f} Y \xrightarrow{g} Z$$

both g and h factor through \mathcal{X}_T . Note that $\tilde{\mathcal{S}}$ is well-defined (see Lemma 2.4).

Let $\mathcal{C}_{\tilde{\mathcal{S}}}$ be the Gabriel-Zisman localisation [10, Chapter 1] of \mathcal{C} with respect to $\tilde{\mathcal{S}}$. This is defined by formally inverting all maps in $\tilde{\mathcal{S}}$. Let $L_{\tilde{\mathcal{S}}}: \mathcal{C} \rightarrow \mathcal{C}_{\tilde{\mathcal{S}}}$ denote the localisation functor. We prove that the functor $H = \text{Hom}_{\mathcal{C}}(T, -): \mathcal{C} \rightarrow \text{mod End}_{\mathcal{C}}(T)^{op}$ inverts the maps in $\tilde{\mathcal{S}}$. Therefore, there is a uniquely defined functor $G: \mathcal{C}_{\tilde{\mathcal{S}}} \rightarrow \text{mod End}_{\mathcal{C}}(T)^{op}$, such that $H = GL_{\tilde{\mathcal{S}}}$. Our main theorem is:

Theorem. *Let \mathcal{C} be a skeletally small Hom-finite Krull-Schmidt triangulated category with rigid object T . Let $\tilde{\mathcal{S}}$ be the class of maps defined above. Then the induced functor $G: \mathcal{C}_{\tilde{\mathcal{S}}} \rightarrow \text{mod End}_{\mathcal{C}}(T)^{op}$ is an equivalence.*

We remark that there is another recent approach to the construction of abelian categories from triangulated categories (as subquotients) by Nakaoka [19] (we discuss this in Section 6).

The paper is organised as follows. In Section 1 we recall some results that we need to use, including a triangulated version of Wakamatsu's Lemma. In Section 2, we study the Hom-functor associated with a rigid object T . A key result in this section is that a map in \mathcal{C} is inverted by $\text{Hom}_{\mathcal{C}}(T, -)$ if and only if it lies in $\tilde{\mathcal{S}}$. In Section 3, we study the main properties of the Gabriel-Zisman localisation of \mathcal{C} at $\tilde{\mathcal{S}}$. In Section 4 we prove our main result, and in Section 5 we describe its relationship to the result of Iyama-Yoshino mentioned above. In Section 6 we give a short discussion of the relationship to the work of Nakaoka [19]. In Section 7 we give some examples.

We would like to thank Yann Palu for his simpler proofs of Lemmas 3.6 and 5.3 and his comments on an earlier version of the paper. We would also like to thank Bill Crawley-Boevey and Henning Krause for useful discussions and the referee for helpful comments. Aslak Buan would like to thank the Algebra, Geometry and Integrable Systems Group and the School of Mathematics at the University of Leeds for their kind hospitality. Robert Marsh would like to thank the Department of Mathematical Sciences at the NTNU, Trondheim, for their kind hospitality.

1. PRELIMINARIES

Let \mathcal{C} be a triangulated Hom-finite Krull-Schmidt category over field k . Let Σ denote the suspension functor of \mathcal{C} .

Let \mathcal{X} be a full subcategory of \mathcal{C} . Then a *right \mathcal{X} -approximation* of C in \mathcal{C} is a map $X \rightarrow C$, with X in \mathcal{X} , such that for all objects Y in \mathcal{X} , the sequence $\text{Hom}_{\mathcal{C}}(Y, X) \rightarrow \text{Hom}_{\mathcal{C}}(Y, C) \rightarrow 0$ is exact. A map $X \xrightarrow{f} C$ is called *right minimal* if, for every $X \xrightarrow{g} X$ such that $fg = f$, we have that g is an isomorphism. A map is called a *minimal right \mathcal{X} -approximation* of C if it is right minimal and it is a right \mathcal{X} -approximation of C . Dually, we have the concepts of *left \mathcal{X} -approximations* and *left minimal* maps.

The full subcategory \mathcal{X} is called *functorially finite* if, for every object C in \mathcal{C} , there exists a right \mathcal{X} -approximation ending in C and a left \mathcal{X} -approximation starting in C .

The following is well-known and straightforward to check.

Lemma 1.1. *Let \mathcal{X} be a full additive subcategory of \mathcal{C} , and C an object of \mathcal{C} .*

- (a) *If there is a right \mathcal{X} -approximation of C then there is a minimal right \mathcal{X} -approximation of C , unique up to isomorphism.*
- (b) *If $g: X \rightarrow C$ is a minimal right \mathcal{X} -approximation of C , then each right approximation is, up to isomorphism, of the form $f \amalg 0: X \amalg X' \rightarrow C$.*

A full subcategory \mathcal{X} of \mathcal{C} is called *extension-closed* if, for each triangle $X' \rightarrow Y \rightarrow X'' \rightarrow \Sigma X'$ in \mathcal{C} with X', X'' in \mathcal{X} , also Y is in \mathcal{X} . Let $\mathcal{X}^{\perp} = \{Y \in \mathcal{C} \mid \text{Ext}_{\mathcal{C}}^1(X, Y) = 0 \text{ for all } X \in \mathcal{X}\}$, and dually let ${}^{\perp}\mathcal{X} = \{Y \in \mathcal{C} \mid \text{Ext}_{\mathcal{C}}^1(Y, X) = 0 \text{ for all } X \in \mathcal{X}\}$.

The next lemma is well-known and is a triangulated version of *Wakamatsu's Lemma*; see e.g. [13, Section 2]; see also [14, Lemma 2.1].

Lemma 1.2. *Let \mathcal{X} be an extension-closed subcategory of a triangulated category \mathcal{C} .*

- (a) *Let $X \rightarrow C$ be a right \mathcal{X} -approximation of C and $\Sigma^{-1}C \rightarrow Y \rightarrow X \rightarrow C$ a completion to a triangle. Then Y is in \mathcal{X}^\perp , and the map $\Sigma^{-1}C \rightarrow Y$ is a left \mathcal{X}^\perp -approximation of $\Sigma^{-1}C$.*
- (b) *Let $C \rightarrow X$ be a left \mathcal{X} -approximation of C and $\Sigma^{-1}Z \rightarrow C \rightarrow X \rightarrow Z \rightarrow \Sigma C$ a completion to a triangle. Then Z is in ${}^\perp\mathcal{X}$, and the map $Z \rightarrow \Sigma C$ is a right ${}^\perp\mathcal{X}$ -approximation of ΣC .*

We shall also often use the fact that in the situation of Lemma 1.2(b), the map $\Sigma^{-1}Z \rightarrow C$ is a right $\Sigma^{-1}({}^\perp\mathcal{X})$ -approximation of C .

2. HOM-FUNCTORS ASSOCIATED TO RIGID OBJECTS

Let T be a rigid object in \mathcal{C} . Let T^\perp denote $(\text{add } T)^\perp$, where $\text{add } T$ denotes the additive closure of T ; we define ${}^\perp T$ similarly. Note that $\Sigma T^\perp = \mathcal{X}_T$, as defined in the introduction. It is easy to see that both $\text{add } T$ and T^\perp are extension closed. In this section we will study these perpendicular categories. We will go on to characterise the maps sent to zero by the functor $\text{Hom}_{\mathcal{C}}(T, -)$ and use this to characterise the maps inverted by this functor.

Using Lemma 1.2, one obtains the following well-known fact (see [13, Prop. 2.3]).

Lemma 2.1. *T^\perp is covariantly finite, and ${}^\perp T$ is contravariantly finite.*

The following lemma is a triangulated variation of the Auslander-Reiten correspondence [1, Prop. 1.10]. This lemma is well-known, but we include the short proof for convenience.

Lemma 2.2. ${}^\perp(T^\perp) = \text{add } T = ({}^\perp T)^\perp$.

Proof. We prove the first equality; the proof of the second is dual. One inclusion is obvious. For the other, let X be in ${}^\perp(T^\perp)$, and let $T_0 \rightarrow X$ be a right $\text{add } T$ -approximation of X . Complete it to a triangle $Z \rightarrow T_0 \rightarrow X \rightarrow \Sigma Z$. Then Z is in T^\perp by Lemma 1.2, hence by the assumption on X we have that $\text{Hom}_{\mathcal{C}}(X, \Sigma T) = 0$, and the triangle splits. Therefore X is in $\text{add } T$. \square

Consider now the functor $F = \text{Hom}_{\mathcal{C}}(T, -): \mathcal{C} \rightarrow \text{mod } \text{End}_{\mathcal{C}}(T)^{op}$. We first determine which maps are killed and which maps are inverted by this functor.

Lemma 2.3. *Let $f: X \rightarrow Y$ be a map in \mathcal{C} . Then $\text{Hom}_{\mathcal{C}}(T, f) = 0$ if and only if f factors through ΣT^\perp .*

Proof. It is clear that $\text{Hom}_{\mathcal{C}}(T, f) = 0$ if f factors through ΣT^\perp .

Now assume $\text{Hom}_{\mathcal{C}}(T, f) = 0$. Consider the right $\text{add } T$ -approximation $T_0 \rightarrow X$. Complete to a triangle

$$U \rightarrow T_0 \rightarrow X \rightarrow \Sigma U,$$

where by Lemma 1.2 we have that U is in T^\perp .

Now we have a diagram

$$\begin{array}{ccccc} U & \longrightarrow & T_0 & \longrightarrow & X & \longrightarrow & \Sigma U \\ & & & & \downarrow f & & \\ & & & & Y & & \end{array}$$

where the composition $T_0 \rightarrow X \xrightarrow{f} Y$ vanishes by assumption. Hence the map f factors through ΣU , which is in ΣT^\perp . This completes the proof. \square

Let $\tilde{\mathcal{S}}$ be the class of maps $f: X \rightarrow Y$ such that when f is completed to a triangle

$$\Sigma^{-1}Z \xrightarrow{h} X \xrightarrow{f} Y \xrightarrow{g} Z$$

both g and h factor through ΣT^\perp .

Lemma 2.4. *The class $\tilde{\mathcal{S}}$ is well-defined.*

Proof. Let $f: X \rightarrow Y$ be a map in \mathcal{C} and let

$$\Sigma^{-1}Z \xrightarrow{h} X \xrightarrow{f} Y \xrightarrow{g} Z$$

and

$$\Sigma^{-1}Z' \xrightarrow{h'} X \xrightarrow{f} Y \xrightarrow{g'} Z$$

be completions of f to a triangle in \mathcal{C} . Since \mathcal{C} is a triangulated category there is a morphism of triangles

$$\begin{array}{ccccccc} \Sigma^{-1}Z & \xrightarrow{h} & X & \xrightarrow{f} & Y & \xrightarrow{g} & Z \\ \downarrow \Sigma^{-1}u & & \parallel & & \parallel & & \downarrow u \\ \Sigma^{-1}Z' & \xrightarrow{h'} & X & \xrightarrow{f} & Y & \xrightarrow{g'} & Z' \end{array}$$

(see e.g. axiom (TR3) in [12, 1.1]). By [12, Prop. 1.2(c)], u (and therefore also $\Sigma^{-1}u$) is an isomorphism. It follows that g factors through \mathcal{X}_T if and only if g' factors through \mathcal{X}_T , and similarly for h and h' . \square

We have the following characterisation of $\tilde{\mathcal{S}}$:

Lemma 2.5. *A map $f: X \rightarrow Y$ belongs to $\tilde{\mathcal{S}}$ if and only if $\mathrm{Hom}_{\mathcal{C}}(T, f)$ is an isomorphism.*

Proof. This follows directly by considering the long exact sequence obtained from applying $\mathrm{Hom}_{\mathcal{C}}(T, -)$ to the triangle $\Sigma^{-1}Z \rightarrow X \rightarrow Y \rightarrow Z$, in combination with Lemma 2.3. \square

Note that the same proof shows:

Lemma 2.6. *Let $f: X \rightarrow Y$ and the completed triangle be as above. Then:*

- (a) *The map h factors through ΣT^\perp if and only if $\mathrm{Hom}_{\mathcal{C}}(T, f)$ is a monomorphism.*
- (b) *The map g factors through ΣT^\perp if and only if $\mathrm{Hom}_{\mathcal{C}}(T, f)$ is an epimorphism.*

(Compare with [17, Thm. 2.3]).

3. LOCALISATION

In this section, we consider the Gabriel-Zisman localisation of \mathcal{C} at the class $\tilde{\mathcal{S}}$ of maps defined above. It turns out, that in order to study this localisation, it is helpful to consider localisation at a smaller better-behaved class of maps \mathcal{S} contained in $\tilde{\mathcal{S}}$ which has the property that the corresponding localisation functor also inverts $\tilde{\mathcal{S}}$. We then investigate some of the properties of localisation at \mathcal{S} which we shall need to prove the main result in the next section.

For a class \mathcal{M} of maps in \mathcal{C} , the *Gabriel-Zisman localisation* [10, Chapter 1] $\mathcal{C}_{\mathcal{M}}$ can be defined as follows. Let the objects in $\mathcal{C}_{\mathcal{M}}$ be the same as the objects in \mathcal{C} . The maps in $\mathcal{C}_{\mathcal{M}}$ are defined as follows (following [18, Section 2.2]). For each element m in \mathcal{M} , introduce an element x_m . This is the formal inverse of m . Then construct an oriented graph \mathcal{G} as follows. The vertices of \mathcal{G} are the objects of \mathcal{C} and the arrows are the maps in \mathcal{C} together with the elements x_m for each m in \mathcal{M} . Here the orientation of the arrow corresponding to a map from X to Y is $X \rightarrow Y$, while for an element m in \mathcal{M} , the edge x_m has the same vertices as m , but the opposite orientation. Then the maps in $\mathcal{C}_{\mathcal{M}}$ from X to Y are equivalence classes of paths in \mathcal{G} starting at X and ending at Y . The equivalence relation is defined as follows. Consider the relation given by the following:

- two consecutive arrows can be replaced by their composition;
- for m in \mathcal{M} , a composition $X \xrightarrow{m} Y \xrightarrow{x_m} X$ or a composition $X \xrightarrow{x_m} Y \xrightarrow{m} X$ can be replaced by $X \xrightarrow{1_X} X$

and close under reflexivity, symmetry and transitivity to obtain an equivalence relation.

Note that we have assumed \mathcal{C} to be skeletally small, in order to avoid set-theoretic problems in forming localisations.

There is a canonical functor $L_{\mathcal{M}}: \mathcal{C} \rightarrow \mathcal{C}_{\mathcal{M}}$ with the property that each functor starting in \mathcal{C} which inverts the maps in \mathcal{M} factors uniquely through $L_{\mathcal{M}}$.

In our situation, by Lemma 2.5, this means that there is a unique functor $G: \mathcal{C}_{\tilde{\mathcal{S}}} \rightarrow \text{mod End}_{\mathcal{C}}(T)^{op}$, making the diagram

$$\begin{array}{ccc}
 \mathcal{C} & \xrightarrow{H=\text{Hom}_{\mathcal{C}}(T,-)} & \text{mod End}_{\mathcal{C}}(T)^{op} \\
 & \searrow L_{\tilde{\mathcal{S}}} & \nearrow G \\
 & & \mathcal{C}_{\tilde{\mathcal{S}}}
 \end{array}$$

commute, and our main theorem is that this functor is actually an equivalence. However, in order to prove this, it will be convenient to consider a subclass \mathcal{S} of $\tilde{\mathcal{S}}$ consisting of maps $f: X \rightarrow Y$ such that when f is completed to a triangle

$$\Sigma^{-1}Z \xrightarrow{h} X \xrightarrow{f} Y \xrightarrow{g} Z,$$

g factors through ΣT^{\perp} , while $\Sigma^{-1}Z$ is in ΣT^{\perp} . Note that this class is also well-defined, using a similar argument to that used for $\tilde{\mathcal{S}}$ in Lemma 2.4 and noting that \mathcal{X}_T is closed under isomorphism. Since \mathcal{S} is contained in $\tilde{\mathcal{S}}$, it is clear that $\text{Hom}_{\mathcal{C}}(T, f)$

is an isomorphism for any map $f: X \rightarrow Y$ in \mathcal{S} , and we get the following diagram:

$$(1) \quad \begin{array}{ccc} \mathcal{C} & \xrightarrow{H=\text{Hom}_{\mathcal{C}}(T,-)} & \text{mod End}_{\mathcal{C}}(T)^{op} \\ & \searrow L_{\mathcal{S}} \quad \nearrow F & \\ & \mathcal{C}_{\mathcal{S}} & \\ & \searrow L_{\tilde{\mathcal{S}}} \quad \downarrow J \quad \nearrow G & \\ & \mathcal{C}_{\tilde{\mathcal{S}}} & \end{array}$$

We claim that the diagram commutes. By the universal property of localisation, we have $FL_{\mathcal{S}} = H$ and $JL_{\mathcal{S}} = L_{\tilde{\mathcal{S}}}$. Using in addition that $GL_{\tilde{\mathcal{S}}} = H$, it follows that $GJL_{\mathcal{S}} = FL_{\mathcal{S}}$. By the universal property of $L_{\mathcal{S}}$, we have $GJ = F$.

Our strategy is to first prove that the functor $F: \mathcal{C}_{\mathcal{S}} \rightarrow \text{mod End}_{\mathcal{C}}(T)^{op}$ is an equivalence. Then, using that both localisation functors invert exactly the maps in $\tilde{\mathcal{S}}$, the functor $\mathcal{C}_{\mathcal{S}} \rightarrow \mathcal{C}_{\tilde{\mathcal{S}}}$ is an equivalence, and our result follows.

Remark 3.1. *The classes of maps \mathcal{S} and $\tilde{\mathcal{S}}$ do not satisfy all of the axioms for admitting a calculus of left fractions or a calculus of right fractions. It is easy to see that the axiom LF3 (and its counterpart RF3), in the notation of [18, Section 3.1], are not in general satisfied. Note that this implies that the corresponding localisation functors do not admit a right or left adjoint; see [18, Section 2.3].*

We consider the full subcategory $\mathcal{C}(T)$ of \mathcal{C} consisting of objects X in \mathcal{C} such that there exists a triangle $T_1 \rightarrow T_0 \rightarrow X \rightarrow \Sigma T_1$ in \mathcal{C} , with T_0, T_1 in $\text{add } T$ (see [13, Prop. 6.2], [16, Section 5.1]). The following characterises the objects in $\mathcal{C}(T)$.

Lemma 3.2. *For an object X in \mathcal{C} , the following are equivalent.*

- (a) X is in $\mathcal{C}(T)$.
- (b) If, in the triangle $U \rightarrow T_0 \xrightarrow{f} X \rightarrow \Sigma U$, the map f is a right $\text{add } T$ -approximation, then U is also in $\text{add } T$.
- (c) If, in the triangle $U \rightarrow T_0 \xrightarrow{f} X \rightarrow \Sigma U$, the map f is a minimal right $\text{add } T$ -approximation, then U is also in $\text{add } T$.

Proof. Assume there is a triangle $T_1 \rightarrow T_0 \rightarrow X \rightarrow \Sigma T_1$ in \mathcal{C} , with T_0, T_1 in $\text{add } T$. Then, using the fact that $\text{Hom}_{\mathcal{C}}(T, -)$ is a homological functor and that T is rigid, it is clear that $T_0 \rightarrow X$ is a right $\text{add } T$ -approximation. The statement now follows from combining this with Lemma 1.1. \square

The importance of $\mathcal{C}(T)$ here is due to the following lemma.

Lemma 3.3. *Let Y be an object in \mathcal{C} . Then there exists a map $f: X \rightarrow Y$ in \mathcal{S} where X is an object in $\mathcal{C}(T)$.*

Proof. Let $T_0 \xrightarrow{u} Y$ be a minimal right $\text{add } T$ -approximation of Y and complete to a triangle

$$Z \rightarrow T_0 \xrightarrow{u} Y \rightarrow \Sigma Z.$$

Let $T_1 \rightarrow Z$ be the minimal right add T -approximation of Z , and complete to a triangle

$$\Sigma^{-1}U \rightarrow T_1 \rightarrow Z \rightarrow U.$$

We have that both Z and $\Sigma^{-1}U$ belong to T^\perp , by Lemma 1.2.

Consider now the following diagram, obtained by applying the octahedral axiom to the composition $T_1 \rightarrow Z \rightarrow T_0$.

$$\begin{array}{ccccccc} \Sigma^{-1}U & \longrightarrow & T_1 & \xrightarrow{w} & Z & \longrightarrow & U \\ \downarrow & & \parallel & & \downarrow v & & \downarrow \\ \Sigma^{-1}X & \longrightarrow & T_1 & \xrightarrow{vw} & T_0 & \longrightarrow & X \\ & & & & \downarrow u & & \downarrow s \\ & & & & Y & \xlongequal{\quad} & Y \\ & & & & \downarrow & & \downarrow t \\ & & & & \Sigma Z & \longrightarrow & \Sigma U \end{array}$$

The map $X \xrightarrow{s} Y$ belongs to \mathcal{S} , since U is in ΣT^\perp and the map $Y \xrightarrow{t} \Sigma U$ factors through ΣZ , which belongs to ΣT^\perp . \square

We consider now the category $\mathcal{C}_\mathcal{S}$ obtained by localising \mathcal{C} with respect to \mathcal{S} . For a map f in \mathcal{C} , we denote its image in $\mathcal{C}_\mathcal{S}$ by $\underline{f} = L_\mathcal{S}(f)$. We provide a series of lemmas needed for the proof of our main theorem.

First note that we have the following direct consequence of Lemma 3.3.

Corollary 3.4. *Let Y be an object of $\mathcal{C}_\mathcal{S}$. Then there is an object X of $\mathcal{C}(T)$ such that $X \simeq Y$ in $\mathcal{C}_\mathcal{S}$.*

Note that a priori we do not know if $\mathcal{C}_\mathcal{S}$ is additive. This will follow from our main result. We will need some elementary properties of $\mathcal{C}_\mathcal{S}$ and the localisation functor $L_\mathcal{S}: \mathcal{C} \rightarrow \mathcal{C}_\mathcal{S}$.

- Lemma 3.5.**
- (a) *For U in ΣT^\perp , the zero map $u_0: U \rightarrow 0$ belongs to \mathcal{S} . Furthermore, the inverse of $\underline{u_0}$ in $\mathcal{C}_\mathcal{S}$ is $\underline{u^0}$, where $u^0: 0 \rightarrow U$ is the zero map.*
 - (b) *For U in ΣT^\perp , and any object X in \mathcal{C} , the projection $\pi_X: X \amalg U \rightarrow X$ belongs to \mathcal{S} . Furthermore, the inverse of $\underline{\pi_X}$ in $\mathcal{C}_\mathcal{S}$ is $\underline{\iota_X}$, where $\iota_X: X \rightarrow X \amalg U$ is the inclusion map.*
 - (c) *Let $u: X \rightarrow Y$ be a map in \mathcal{C} factoring through ΣT^\perp . Then $\underline{u} = \underline{0}$ in $\mathcal{C}_\mathcal{S}$.*
 - (d) *Let $u, v: X \rightarrow Y$ be maps in \mathcal{C} , such that v factors through ΣT^\perp . Then $\underline{u+v} = \underline{u}$ in $\mathcal{C}_\mathcal{S}$.*

Proof. Part (a) is straightforward. For (b) consider the triangle

$$U \rightarrow U \amalg X \xrightarrow{\pi_X} X \xrightarrow{0} \Sigma U.$$

Since U is in ΣT^\perp , we have that π_X belongs to \mathcal{S} . Note that as $\pi_X \iota_X = 1_X$ in \mathcal{C} , we also have that $\underline{\pi_X \iota_X} = \underline{1_X}$ in $\mathcal{C}_\mathcal{S}$. This proves the second claim. Part (c) is a

direct consequence of (a). To prove (d), assume v factors as $X \xrightarrow{f} U \xrightarrow{g} Y$, with U in ΣT^\perp . Then, using (b), we obtain

$$\begin{aligned} \underline{u + v} &= \underline{u + gf} = \underline{(g \ 1_Y)} \underline{\begin{pmatrix} f \\ u \end{pmatrix}} = \underline{(g \ 1_Y)} \underline{\begin{pmatrix} 1_X & 0 \\ 0 & u \end{pmatrix}} \underline{\begin{pmatrix} f \\ 1_X \end{pmatrix}} = \\ &= \underline{(g \ 1_Y)} \underline{\iota_Y \pi_Y} \underline{\begin{pmatrix} 1_U & 0 \\ 0 & u \end{pmatrix}} \underline{\iota_X \pi_X} \underline{\begin{pmatrix} f \\ 1_X \end{pmatrix}} = \underline{1_Y} \underline{(0 \ 1_Y)} \underline{\begin{pmatrix} 1_U & 0 \\ 0 & u \end{pmatrix}} \underline{\begin{pmatrix} 0 \\ 1_X \end{pmatrix}} \underline{1_X} = \underline{u}, \end{aligned}$$

where (c) is used for the second-to-last equality. \square

The next lemma will be helpful in simplifying the description of maps in \mathcal{C}_S . We would like to thank Yann Palu for this simplification of an earlier version of the proof of the lemma.

Lemma 3.6. *Let U be an object in $\mathcal{C}(T)$, let $u: U \rightarrow Y$ be a map in \mathcal{C} and let $s: X \rightarrow Y$ be a map in \mathcal{S} . Then u factors through s .*

Proof. Since U lies in $\mathcal{C}(T)$, there is a triangle:

$$T_1 \rightarrow T_0 \rightarrow U \rightarrow \Sigma T_1$$

with T_0, T_1 in $\text{add} T$. Complete s to a triangle $\Sigma^{-1}Z \xrightarrow{h} X \xrightarrow{s} Y \xrightarrow{g} Z$ where $\Sigma^{-1}Z$ lies in ΣT^\perp and g factors through ΣT^\perp . We thus have the diagram:

$$\begin{array}{ccccccc} T_1 & \longrightarrow & T_0 & \xrightarrow{f} & U & \longrightarrow & \Sigma T_1 \\ \downarrow \text{dotted} & & \downarrow \text{dotted} & & \downarrow u & & \downarrow \text{dotted} \\ \Sigma^{-1}Z & \xrightarrow{h} & X & \xrightarrow{s} & Y & \xrightarrow{g} & Z \end{array}$$

Since T_0 lies in $\text{add} T$ and g factors through ΣT^\perp , $guf = 0$ and there are maps $T_1 \rightarrow \Sigma^{-1}Z$ and $T_0 \rightarrow X$ giving a map of triangles as indicated in the diagram. Since T_1 lies in $\text{add} T$ and Z lies in $\Sigma^2 T^\perp$, the map v vanishes, so $gu = 0$ and u factors through s as required. \square

As a consequence of this we obtain the following.

Proposition 3.7. *Let U, V be objects in \mathcal{C} with U in $\mathcal{C}(T)$. Then $\text{Hom}_{\mathcal{C}}(U, V) \rightarrow \text{Hom}_{\mathcal{C}_S}(U, V)$ is surjective.*

Proof. A map in \mathcal{C}_S is a composition of maps in \mathcal{C} and formal inverses of maps in \mathcal{S} . Assume we have a composition $U \xrightarrow{u} Y \xrightarrow{s^{-1}} X$ for a map $X \xrightarrow{s} Y$ in \mathcal{S} . Then, in \mathcal{C} we have the commutative diagram

$$\begin{array}{ccc} & U & \\ h \swarrow & \downarrow u & \\ X & \xrightarrow{s} & Y \end{array}$$

where the map $h: U \rightarrow X$ such that $u = sh$ exists by Lemma 3.6.

Now, $\underline{s^{-1}u} = \underline{s^{-1}sh} = \underline{h}$.

By this it is clear that any map in \mathcal{C}_S from the object U of $\mathcal{C}(T)$ can be obtained as the image of a map in \mathcal{C} . \square

4. MAIN RESULT

As before, let T be a rigid object in the triangulated Hom-finite Krull-Schmidt category \mathcal{C} , let \mathcal{S} be as described in Section 3 and let $\tilde{\mathcal{S}}$ be as defined in Section 2. In this section we prove our main result, i.e. that if \mathcal{C} is skeletally small, the localisation of \mathcal{C} at the class $\tilde{\mathcal{S}}$ is equivalent to $\text{mod End}_{\mathcal{C}}(T)$.

We first consider the canonical functor $F: \mathcal{C}_{\mathcal{S}} \rightarrow \text{mod End}_{\mathcal{C}}(T)^{op}$, as described in Section 3, and prove that it is an equivalence.

Proposition 4.1. *The functor $F: \mathcal{C}_{\mathcal{S}} \rightarrow \text{mod End}_{\mathcal{C}}(T)^{op}$ is an equivalence.*

We recall the following result:

Theorem 4.2 (Iyama-Yoshino). *Let T be a rigid object in a Hom-finite Krull-Schmidt triangulated category \mathcal{C} . Then the functor $\text{Hom}_{\mathcal{C}}(T, -)$ induces an equivalence*

$$\mathcal{C}(T)/\Sigma T \rightarrow \text{mod End}_{\mathcal{C}}(T)^{op}.$$

This was proved in [13, Prop. 6.2]. The case where \mathcal{C} is d -Calabi-Yau was proved in [16, Section 5.1]. In view of Lemma 2.3, we will need the following variation of Theorem 4.2; the proof is fairly similar. For an additive category \mathcal{C} and additive subcategory \mathcal{C}' , we will use \mathcal{C}/\mathcal{C}' to denote the quotient of \mathcal{C} by the ideal of maps which factor through \mathcal{C}' ; it is also an additive category. For an object C in \mathcal{C} , we let \mathcal{C}/C denote $\mathcal{C}/\text{add } C$.

Lemma 4.3. *The functor $H = \text{Hom}_{\mathcal{C}}(T, -): \mathcal{C} \rightarrow \text{mod End}_{\mathcal{C}}(T)^{op}$ is dense. Its restriction H' to $\mathcal{C}(T)$ is full and induces an equivalence $\mathcal{C}(T)/\Sigma T^{\perp} \rightarrow \text{mod End}_{\mathcal{C}}(T)^{op}$.*

Proof. Let M be a finite dimensional $\text{End}_{\mathcal{C}}(T)^{op}$ -module and choose a minimal projective presentation

$$P_1 \rightarrow P_0 \rightarrow M \rightarrow 0.$$

Since $\text{Hom}_{\mathcal{C}}(T, -)$ induces an equivalence from $\text{add } T$ to the finite dimensional projective $\text{End}_{\mathcal{C}}(T)^{op}$ -modules, the map $P_1 \rightarrow P_0$ is the image of a map $T_1 \rightarrow T_0$ in $\text{add } T$. Complete to a triangle

$$T_1 \rightarrow T_0 \rightarrow X \rightarrow \Sigma T_1.$$

Since $\text{Hom}_{\mathcal{C}}(T, \Sigma T_1) = 0$, we get an exact sequence

$$\text{Hom}_{\mathcal{C}}(T, T_1) \rightarrow \text{Hom}_{\mathcal{C}}(T, T_0) \rightarrow \text{Hom}_{\mathcal{C}}(T, X) \rightarrow 0.$$

Hence, it is clear that $M \simeq \text{Hom}_{\mathcal{C}}(T, X)$ and thus that H is dense.

Now, let X, Y be objects in $\mathcal{C}(T)$, and consider a map $\alpha: \text{Hom}_{\mathcal{C}}(T, X) \rightarrow \text{Hom}_{\mathcal{C}}(T, Y)$. Then there are triangles $T_1 \rightarrow T_0 \rightarrow X \rightarrow \Sigma T_1$ and $U_1 \rightarrow U_0 \rightarrow Y \rightarrow \Sigma U_1$ whose images under $\text{Hom}_{\mathcal{C}}(T, -)$ give minimal projective presentations for $\text{Hom}_{\mathcal{C}}(T, X)$ and $\text{Hom}_{\mathcal{C}}(T, Y)$, respectively.

Hence there are vertical maps such that the following diagram commutes:

$$\begin{array}{ccccccc} \text{Hom}_{\mathcal{C}}(T, T_1) & \longrightarrow & \text{Hom}_{\mathcal{C}}(T, T_0) & \longrightarrow & \text{Hom}_{\mathcal{C}}(T, X) & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow \alpha & & \\ \text{Hom}_{\mathcal{C}}(T, U_1) & \longrightarrow & \text{Hom}_{\mathcal{C}}(T, U_0) & \longrightarrow & \text{Hom}_{\mathcal{C}}(T, Y) & \longrightarrow & 0 \end{array}$$

Lifting the left hand square of this diagram to add T , we obtain a diagram

$$\begin{array}{ccccccc} T_1 & \longrightarrow & T_0 & \longrightarrow & X & \longrightarrow & \Sigma T_1 \\ \downarrow & & \downarrow & & \vdots & & \downarrow \\ U_1 & \longrightarrow & U_0 & \longrightarrow & Y & \longrightarrow & \Sigma U_1 \end{array}$$

The induced map $X \rightarrow Y$ is mapped to α and hence the functor H' is full. The statement now follows from Lemma 2.3. \square

Proof of Proposition 4.1. First note that for an object X in \mathcal{C}_S we have that $F(X) = H(X) = \text{Hom}_{\mathcal{C}}(T, X)$. If $f: X \rightarrow Y$ is a map in \mathcal{C} , we have that $F(\underline{f}) = \text{Hom}_{\mathcal{C}}(T, f)$.

We have that F is dense by Lemma 4.3 and the fact that $H = FL_S$. We now show that that F is full and faithful. For this let X, Y be objects in \mathcal{C}_S . By Lemma 3.3, there are objects X', Y' in $\mathcal{C}(T)$, and maps $u: X' \rightarrow X$ and $v: Y' \rightarrow Y$ in \mathcal{S} , such that $\underline{u}: X' \rightarrow X$ and $\underline{v}: Y' \rightarrow Y$ are isomorphisms. Let $\beta = F(\underline{u})$, and let $\gamma = F(\underline{v})$.

F is full:

Let $\alpha: FX \rightarrow FY$ be an arbitrary map in $\text{mod End}_{\mathcal{C}}(T)^{op}$, and let $\alpha' = \gamma^{-1}\alpha\beta$, i.e. we have the commutative diagram

$$\begin{array}{ccc} FX' & \xrightarrow{\beta} & FX \\ \downarrow \alpha' & & \downarrow \alpha \\ FY' & \xrightarrow{\gamma} & FY \end{array}$$

By Lemma 4.3 H' is dense, so, since $FL_S = H$ and X' lies in $\mathcal{C}(T)$, there is a map $f': X' \rightarrow Y'$ in \mathcal{C} such that $FL_S(f') = \alpha'$. Hence $F(\underline{f}') = \alpha'$.

Setting $f = \underline{v}\underline{f}'\underline{u}^{-1}: X \rightarrow Y$ in \mathcal{C}_S , we obtain $F(f) = F(\underline{v})F(\underline{f}')F(\underline{u})^{-1} = \gamma\alpha'\beta^{-1} = \gamma\gamma^{-1}\alpha\beta\beta^{-1} = \alpha$ as required.

F is faithful:

Let X, Y be objects of \mathcal{C}_S , and assume we have maps $f, g: X \rightarrow Y$ in \mathcal{C}_S such that $F(f) = F(g)$. By Proposition 3.7, there are $f', g': X' \rightarrow Y'$ in \mathcal{C} such that $\underline{f}' = \underline{v}^{-1}f\underline{u}$ and $\underline{g}' = \underline{v}^{-1}g\underline{u}$.

Then

$$\begin{aligned} F(\underline{f}') &= F(\underline{v}^{-1}f\underline{u}) = F(\underline{v})^{-1}F(f)F(\underline{u}) = \\ & F(\underline{v})^{-1}F(g)F(\underline{u}) = F(\underline{v}^{-1}g\underline{u}) = F(\underline{g}'). \end{aligned}$$

Hence we have that $H(\underline{f}') = H(\underline{g}')$, and by Lemma 4.3, this implies that $\underline{g}' - \underline{f}'$ factors through ΣT^\perp in \mathcal{C} . By Lemma 3.5 we have that

$$\underline{f}' = \underline{f}' + (\underline{g}' - \underline{f}') = \underline{g}'$$

in \mathcal{C}_S , and hence $f = \underline{v}\underline{f}'\underline{u}^{-1} = \underline{v}\underline{g}'\underline{u}^{-1} = g$ in \mathcal{C}_S as required.

Hence F is dense, full and faithful, and we are done. \square

By Lemma 2.5, we have that $\text{Hom}_{\mathcal{C}}(T, f)$ is invertible if and only if f belongs to $\tilde{\mathcal{S}}$. Since $FL_S = H$ (see Diagram (1)) and F is an equivalence (by Proposition 4.1), it follows that L_S inverts the maps in $\tilde{\mathcal{S}}$. Hence there is an induced functor $I: C_{\tilde{\mathcal{S}}} \rightarrow C_S$

such that $IL_{\tilde{\mathcal{S}}} = L_{\mathcal{S}}$. Since $L_{\tilde{\mathcal{S}}}$ inverts the maps in \mathcal{S} there is an induced functor $J: C_{\mathcal{S}} \rightarrow C_{\tilde{\mathcal{S}}}$ such that $JL_{\mathcal{S}} = L_{\tilde{\mathcal{S}}}$. It follows that $IJJ_{\mathcal{S}} = L_{\mathcal{S}}$. It follows from the universal property of $L_{\mathcal{S}}$ that IJ is equal to the identity functor on $C_{\mathcal{S}}$. Similarly JI is equal to the identity functor on $C_{\tilde{\mathcal{S}}}$. Hence the induced functors I and J between $C_{\mathcal{S}}$ and $C_{\tilde{\mathcal{S}}}$ are isomorphisms. Our main theorem follows:

Theorem 4.4. *Let \mathcal{C} be a skeletally small Hom-finite Krull-Schmidt triangulated category with rigid object T . Let $\tilde{\mathcal{S}}$ be the class of maps defined above. Then the induced functor $G: C_{\tilde{\mathcal{S}}} \rightarrow \text{mod End}_{\mathcal{C}}(T)^{op}$ is an equivalence.*

5. THE CLUSTER-TILTING OBJECT CASE

In this section we compare our results with recent work showing how to obtain module categories of (opposite) endomorphism algebras of rigid objects in triangulated categories from the categories themselves.

We call a rigid object T in \mathcal{C} a *cluster-tilting object* if $\text{Ext}_{\mathcal{C}}^1(T, X) = 0$ only if X is in $\text{add } T$. Note that by [17, Lemma 3.2] (or, alternatively, Lemma 2.2), it also follows that $\text{Ext}_{\mathcal{C}}^1(X, T) = 0$ only if X is in $\text{add } T$, when T is a cluster-tilting object (see also [17, Defn. 3.1]), so the cluster-tilting objects studied here coincide with the maximal 1-orthogonal objects studied in [13, 17]. The motivation for our work was the following theorem, which generalises previous work dealing with Calabi-Yau triangulated categories [16, Prop. 2.1] or cluster categories [8, Thm. 2.2] (note that part (a) was recalled as Theorem 4.2 above but we repeat it here for comparison). Also note that we state results here only as they apply to the object case (rather than for a rigid subcategory).

Theorem 5.1. *Let T be a rigid object in a Hom-finite Krull-Schmidt triangulated category \mathcal{C} .*

(a) [Iyama-Yoshino] *The functor $\text{Hom}_{\mathcal{C}}(T, -)$ induces an equivalence*

$$\mathcal{C}(T)/\Sigma T \rightarrow \text{mod End}_{\mathcal{C}}(T)^{op}.$$

(b) [Koenig-Zhu] *If T is a cluster-tilting object then there is an equivalence*

$$\mathcal{C}/\text{add } \Sigma T \rightarrow \text{mod End}_{\mathcal{C}}(T)^{op}.$$

As mentioned above, part (a) was proved in [13, Prop. 6.2] (the d -Calabi Yau case in [16, Section 5.1]). Part (b) was proved in [17, Cor. 4.4].

Note that to prove (b) from (a), it is sufficient to realise that $\mathcal{C}(T) = \mathcal{C}$ in the case where T is cluster-tilting. This is well-known; we repeat the easy proof here for convenience.

Lemma 5.2. *If T is a cluster-tilting object in \mathcal{C} , then $\mathcal{C}(T) = \mathcal{C}$.*

Proof. Let C be in \mathcal{C} and consider the triangle

$$U \rightarrow T_0 \rightarrow C \rightarrow \Sigma U,$$

obtained by completing a right $\text{add } T$ -approximation $T_0 \rightarrow C$. Then U lies in $T^{\perp} = \text{add } T$ by Lemma 1.2. \square

In this section, we point out how Theorem 5.1 relates to our main theorem.

We first give a description of the maps with domain in $\mathcal{C}(T)$ which lie in the kernel of $\text{Hom}_{\mathcal{C}}(T, -)$. We would like to thank Yann Palu for his short proof of this lemma which replaces the longer version in an earlier version of the paper.

Lemma 5.3. *Let X, Y be objects in \mathcal{C} , with X in $\mathcal{C}(T)$. Suppose that $f: X \rightarrow Y$ factors through ΣT^\perp . Then f factors through $\text{add } \Sigma T$.*

Proof. Since X lies in $\mathcal{C}(T)$, there is a triangle:

$$T_1 \rightarrow T_0 \xrightarrow{g} X \xrightarrow{h} \Sigma T_1,$$

with T_0, T_1 in $\text{add } T$. Since $f: X \rightarrow Y$ factors through ΣT^\perp and T_0 lies in $\text{add } T$, $fg = 0$, so f factors through h and hence through $\text{add } \Sigma T$, since T_1 lies in $\text{add } T$. \square

Combining Lemmas 5.3 and 4.3, we obtain part (a) of Theorem 5.1. Part (b) follows from the observation that $\mathcal{C}(T) = \mathcal{C}$, when T is a cluster-tilting object.

Summarising, we have the following:

Theorem 5.4. *There are equivalences of categories*

$$\mathcal{C}(T)/\Sigma T \rightarrow \mathcal{C}(T)/\Sigma T^\perp \rightarrow \text{mod } \text{End}_{\mathcal{C}}(T)^{op} \rightarrow \mathcal{C}_{\tilde{\mathcal{S}}}$$

Proof. The first equivalence follows from Lemma 5.3, the second is Lemma 4.3, while the third is our main result, Theorem 4.4. \square

Assume T is a cluster-tilting object in \mathcal{C} , so that we have $\mathcal{C} = \mathcal{C}(T)$. Let us finish by pointing out that the equivalence obtained by the composition $\mathcal{C}/\Sigma T^\perp \rightarrow \text{mod } \text{End}_{\mathcal{C}}(T)^{op} \rightarrow \mathcal{C}_{\tilde{\mathcal{S}}}$ in the above theorem, has a natural interpretation.

Let $Q: \mathcal{C} \rightarrow \mathcal{C}/\Sigma T^\perp$ be the canonical quotient functor, and $H_0: \mathcal{C}/\Sigma T^\perp \rightarrow \text{mod } \text{End}_{\mathcal{C}}(T)^{op}$ the induced functor. We then have $H_0Q = H = GL_{\tilde{\mathcal{S}}}$. It is clear that Q maps ΣT^\perp to 0, and is universal among additive functors with respect to this property. Moreover, by Lemma 2.5 in combination with the fact that H_0 is an equivalence, it is clear that Q inverts the maps in $\tilde{\mathcal{S}}$.

On the other hand, $L_{\tilde{\mathcal{S}}}$ is universal with respect to inverting $\tilde{\mathcal{S}}$, and it is clear that $L_{\tilde{\mathcal{S}}}$ maps ΣT^\perp to 0. Moreover, $L_{\tilde{\mathcal{S}}}$ is additive with respect to the additive structure on $\mathcal{C}_{\tilde{\mathcal{S}}}$ induced by the equivalence G .

Combining these two facts, we obtain canonical functors $U: \mathcal{C}/\Sigma T^\perp \rightarrow \mathcal{C}_{\tilde{\mathcal{S}}}$, and $V: \mathcal{C}_{\tilde{\mathcal{S}}} \rightarrow \mathcal{C}/\Sigma T^\perp$. Arguing as for diagram (1), and using that Q is full, we obtain a commutative diagram

$$\begin{array}{ccc}
\mathcal{C} & \xrightarrow{H = \text{Hom}_{\mathcal{C}}(T, -)} & \text{mod End}_{\mathcal{C}}(T)^{op} \\
& \searrow Q & \nearrow H_0 \\
& \mathcal{C}/\Sigma T^{\perp} & \\
& \swarrow L_{\bar{\mathcal{S}}} & \searrow G \\
& \mathcal{C}_{\bar{\mathcal{S}}} & \\
& \uparrow V & \downarrow U
\end{array}$$

By universality, it follows that U is an isomorphism and that V is the inverse of U .

6. COTORSION PAIRS

Nakaoka [19] considers the notion of a *cotorsion pair* in a triangulated category. In this section we compare the results obtained here to those of Nakaoka.

According to [19, 2.3] a cotorsion pair can be defined as a pair $(\mathcal{U}, \mathcal{V})$ of full additive subcategories satisfying

- (a) $U^{\perp} = \mathcal{V}$;
- (b) ${}^{\perp}\mathcal{V} = \mathcal{U}$;
- (c) For any object C , there is a (not necessarily unique) triangle:

$$U \rightarrow C \rightarrow \Sigma V \rightarrow \Sigma U,$$

with $U \in \mathcal{U}$ and $V \in \mathcal{V}$.

Nakaoka points out that $(\mathcal{U}, \mathcal{V})$ is a cotorsion pair if and only if $(\mathcal{U}, \Sigma\mathcal{V})$ is a torsion theory in the sense of [13, 2.2]. This is not the same as a torsion theory in the sense of [5, Defn. 2.1], since there is no assumption of closure under the suspension functor. It follows from Wakamatsu's Lemma (see Lemma 1.2) and Lemma 2.2 that, in our context, $(\text{add } T, T^{\perp})$ is a torsion pair.

Nakaoka proves the following theorem:

Theorem 6.1. *Let \mathcal{C} be a triangulated category and $(\mathcal{U}, \mathcal{V})$ a cotorsion pair in \mathcal{C} . Let $\mathcal{W} = \mathcal{U} \cap \mathcal{V}$. Let \mathcal{C}^+ be the full subcategory of \mathcal{C} consisting of objects C such that there is a distinguished triangle*

$$W \rightarrow C \rightarrow \Sigma^{-1}V \rightarrow \Sigma W$$

in \mathcal{C} with $W \in \mathcal{W}$ and $V \in \mathcal{V}$. Let \mathcal{C}^- be the full subcategory of \mathcal{C} consisting of objects C such that there is a distinguished triangle

$$\Sigma^{-1}U \rightarrow C \rightarrow W \rightarrow U$$

in \mathcal{C} with $U \in \mathcal{U}$ and $W \in \mathcal{W}$. Then $\mathcal{C}^+ \cap \mathcal{C}^-$ contains $\mathcal{U} \cap \mathcal{V}$ and

$$\mathcal{H} := \frac{\mathcal{C}^+ \cap \mathcal{C}^-}{\mathcal{U} \cap \mathcal{V}}$$

is abelian.

If $(\mathcal{U}, \mathcal{V}) = (\text{add } T, T^\perp)$, where T is a rigid object in \mathcal{C} then it is easy to check that $\mathcal{C}^+ = \mathcal{C}$ and $\mathcal{C}^- = \Sigma^{-1}\mathcal{C}(T)$. Thus, in this case, Nakaoka’s result produces the subfactor abelian category $\text{mod } \text{End}_{\mathcal{C}}(T)^{op}$ as in the Theorem 5.1(a) of Iyama-Yoshino, whereas in our approach we produce this category via localisation.

Nakaoka already points out that the special case where $\mathcal{U} = \mathcal{V}$ is a cluster-tilting object (or, more generally, a cluster-tilting subcategory) recovers a result of Koenig-Zhu [17, 3.3] (see Theorem 5.1(b)).

It is thus a natural question to ask whether the subfactor abelian category \mathcal{H} in Nakaoka’s theorem can be obtained from the triangulated category via localisation. However, the methods of this paper do not apply in this situation since an appropriate generalisation of Lemma 4.3 does not hold in general for a cotorsion pair.

In a sequel to this article [6], we provide an alternative approach to showing that $\mathcal{C}_{\tilde{\mathcal{S}}}$ is abelian, by first considering the factor category $\mathcal{C}/\mathcal{X}_T$. We prove that this category is preabelian, and that the family of regular maps \mathcal{R} admits a calculus of left/right fractions, and moreover that we can recover $\text{mod } \Gamma$ by localising with respect to \mathcal{R} . We note that although this approach is more closely related to the work of Koenig-Zhu [17] and Nakaoka [19], it still does not apply in the more general cotorsion theory set-up considered by Nakaoka.

7. EXAMPLES

In this section we give two examples to illustrate the main result. We refer to [2] for background on finite dimensional algebras and their representation theory. The first example includes a map which lies in $\tilde{\mathcal{S}}$ but not in \mathcal{S} . In the second example, \mathcal{C} does not contain any cluster-tilting objects; it is also interesting to see how a module category of finite representation type arises from the localisation of a cluster category with infinitely many indecomposable objects. We also see that it is possible that the image of a indecomposable module under localisation at $\tilde{\mathcal{S}}$ can be decomposable.

Example 7.1. Let Q be the quiver:

$$1 \longrightarrow 2 \longrightarrow 3 \longrightarrow 4.$$

The indecomposable modules over kQ are determined by their support on the vertices of Q . We denote by M_{ij} the indecomposable with support $\{i, i + 1, \dots, j\}$. Let T be the rigid module $T_1 \amalg T_2 \amalg T_3$, where $T_1 = M_{44}$, $T_2 = M_{14}$ and $T_3 = M_{11}$. Regarded as an object in the cluster category \mathcal{C} corresponding to kQ , T is rigid, and it is easy to check that $\text{End}_{\mathcal{C}}(T)^{op}$ is isomorphic to the algebra Λ given by the following quiver with a single relation:


$$1 \xleftarrow{\dots\dots\dots} 2 \xleftarrow{\quad} 3.$$

The indecomposable Λ -modules are the simple modules S_1, S_2, S_3 and the modules S_2 and S_3 with dimension vectors $(1, 1, 0)$ and $(0, 1, 1)$ respectively. If M is any object in \mathcal{C} then

$$G(L_{\tilde{\mathcal{S}}}(M)) = H(M) = \text{Hom}_{\mathcal{C}}(T, M).$$

This enables us to compute the $\text{End}_{\mathcal{C}}(T)^{op}$ -module $G(M)$ corresponding to each indecomposable object M in \mathcal{C} .

The images of the indecomposable objects of \mathcal{C} under $\text{Hom}_{\mathcal{C}}(T, -)$ and the images of the irreducible maps between them are given in their corresponding positions in the Auslander-Reiten quiver of \mathcal{C} below (drawn with the images of the indecomposable projective modules on the left hand side, repeated on the right hand side).



Thus we see that $\text{Hom}_{\mathcal{C}}(T, -)$ is dense, as predicted by Lemma 4.3. We note that an indecomposable object in \mathcal{C} can be sent to a decomposable object by $\text{Hom}_{\mathcal{C}}(T, -)$. We also remark that, in this example, all of the irreducible maps in $\text{mod } \text{End}_{\mathcal{C}}(T)^{op}$ are of the form $\text{Hom}_{\mathcal{C}}(T, u)$ where u is irreducible in \mathcal{C} .

The right minimal almost split map $s: M_{44} \amalg \Sigma M_{24} \rightarrow M_{34}$ can be completed to the triangle


$$\Sigma M_{34} \rightarrow M_{44} \amalg \Sigma M_{24} \rightarrow M_{34} \rightarrow M_{13}.$$

Since ΣM_{34} lies in ΣT^{\perp} and the map $M_{34} \rightarrow M_{13}$ factors through $M_{23} \in \Sigma T^{\perp}$, it follows that $s \in \mathcal{S}$ and thus is inverted by the localisation functor $L_{\tilde{\mathcal{S}}}$. Under the functor $H = \text{Hom}_{\mathcal{C}}(T, -)$, the map s is mapped to an isomorphism


$$\underline{s}: S_1 \amalg S_3 \rightarrow S_1 \amalg S_3.$$

We note that the left minimal almost split map $t: M_{34} \rightarrow M_{33} \amalg M_{24}$ lies in $\tilde{\mathcal{S}}$ but not in \mathcal{S} . It is also inverted by $L_{\tilde{\mathcal{S}}}$.

Example 7.2. In this example we assume that k is algebraically closed. We consider the path algebra of the following quiver Q of type \tilde{A}_2 :



The simple module S_2 is a rigid module at the mouth of a tube \mathcal{T} of rank 2 in the module category of this quiver, giving rise to a rigid object T in the cluster category \mathcal{C} . Then $\Gamma = \text{End}_{\mathcal{C}}(T)^{op}$ is given by a quiver with a single vertex and a loop whose square is zero. The indecomposable Γ -modules are the uniserial projective module, P , and the simple module S . Note that $\text{Hom}_{\mathcal{C}}(T, -)$ kills all of the objects in the homogeneous tubes. The images of the other indecomposable objects in \mathcal{C} and the images of the irreducible maps between them under $\text{Hom}_{\mathcal{C}}(T, -)$ are given in their corresponding positions in the Auslander-Reiten quiver below.



We note that the cluster category $\mathcal{C}_{\mathcal{T}}$ of a tube \mathcal{T} (defined using [7, Section 3]) is a full subcategory of the cluster category of $\text{mod } kQ$ and has been studied in [3, 4, 9, 21]. It does not contain any cluster-tilting objects. Theorem 4.4 applies to the case of a maximal rigid object T in $\mathcal{C}_{\mathcal{T}}$. It follows that the module categories over the algebras studied in [21], i.e. the endomorphism algebras of maximal rigid objects in $\mathcal{C}_{\mathcal{T}}$, can be obtained as localisations of $\mathcal{C}_{\mathcal{T}}$. The above example gives rise to the case of a rank two tube, with T a maximal rigid object in $\mathcal{C}_{\mathcal{T}}$. We see that, as in [21, Thm. 4.9], $\text{Hom}_{\mathcal{C}_{\mathcal{T}}}(T, M)$ can be decomposable even if M is indecomposable. It follows from Theorem 4.4 that the same is true for $L_{\tilde{\mathcal{S}}}(M)$.

REFERENCES

- [1] Auslander, Maurice; Reiten, Idun *Applications of contravariantly finite subcategories*, Adv. Math. 86 (1991), no. 1, 111–152.
- [2] Auslander, Maurice; Reiten, Idun; Smalø, Sverre O. *Representation theory of Artin algebras*, Cambridge Studies in Advanced Mathematics, 36. Cambridge University Press, Cambridge, 1997.
- [3] Barot, Michael; Kussin, Dirk; Lenzing, Helmut *The Grothendieck group of a cluster category*, J. Pure Appl. Algebra 212(1) (2008), 33–46.
- [4] Barot, Michael; Kussin, Dirk; Lenzing, Helmut *The cluster category of a canonical algebra*, Trans. Amer. Math. Soc. 362 (2010), 4313–4330.
- [5] Beligiannis, Apostolos; Reiten, Idun *Homological and homotopical aspects of torsion theories*, Mem. Amer. Math. Soc. 188 (2007), no. 883.
- [6] Buan, Aslak Bakke; Marsh, Robert J. *From triangulated categories to module categories via localisation II: calculus of fractions*, preprint arXiv:1102.4597 [math.CT], 2011.
- [7] Buan, Aslak Bakke; Marsh, Robert J.; Reineke, Markus; Reiten, Idun; Todorov, Gordana *Tilting theory and cluster combinatorics*, Adv. Math. **204** (2006), 572–618.
- [8] Buan, Aslak Bakke; Marsh, Robert J.; Reiten, Idun *Cluster-tilted algebras*, Trans. Amer. Math. Soc. 359 (2007), no. 1, 323–332.
- [9] Buan, Aslak Bakke; Marsh, Robert J.; Vatne, Dagfinn Furnes *Cluster structures from 2-Calabi-Yau categories with loops*, Math. Z. **265**(4) (2010), 951–970.
- [10] Gabriel, P.; Zisman, M. *Calculus of fractions and homotopy theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 35 Springer-Verlag New York, Inc., New York 1967.
- [11] Gelfand, Sergei I.; Manin, Yuri I *Methods of homological algebra*, Second edition. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2003.

- [12] Happel, Dieter *Triangulated categories in the representation theory of finite-dimensional algebras*, London Mathematical Society Lecture Note Series, 119. Cambridge University Press, Cambridge, 1988.
- [13] Iyama, Osamu; Yoshino, Yuji *Mutation in triangulated categories and rigid Cohen-Macaulay modules*, *Invent. Math.* 172 (2008), no. 1, 117–168.
- [14] Jørgensen, Peter *Auslander-Reiten triangles in subcategories*, *J. K-Theory* 3 (2009), no. 3, 583–601.
- [15] Keller, Bernhard *Cluster algebras, quiver representations and triangulated categories*, in *Triangulated Categories* (edited by Holm, T; Jørgensen, P.; Rouquier, R.), London Math. Soc. Lecture Note Ser., vol. 375, Cambridge University Press, Cambridge, 2010.
- [16] Keller, Bernhard; Reiten, Idun *Cluster-tilted algebras are Gorenstein and stably Calabi-Yau*, *Adv. Math.* 211 (2007), no. 1, 123–151.
- [17] Koenig, Steffen; Zhu, Bin *From triangulated categories to abelian categories: cluster tilting in a general framework*, *Math. Z.* 258 (2008), no. 1, 143–160.
- [18] Krause, Henning *Localization theory for triangulated categories*, in *Triangulated Categories* (edited by Holm, T; Jørgensen, P.; Rouquier, R.), London Math. Soc. Lecture Note Ser., vol. 375, Cambridge University Press, Cambridge, 2010.
- [19] Nakaoka, Hiroyuki *General heart construction on a triangulated category (I): unifying t-structures and cluster tilting subcategories*, *Appl. Categ. Structures*, Online First 2010, DOI: 10.1007/s10485-010-9223-2.
- [20] Reiten, Idun *Cluster categories*, Preprint arXiv:1012.4949v1 [math.RT], 2010, to appear in *Proceedings of ICM2010*.
- [21] Vatne, Dagfinn Furnes *Endomorphism rings of maximal rigid objects in cluster tubes*, Preprint arXiv:0905.1796v1 [math.RT], 2009.

INSTITUTT FOR MATEMATISKE FAG, NORGES TEKNISK-NATURVITENSKAPELIGE UNIVERSITET, N-7491 TRONDHEIM, NORWAY

E-mail address: `aslakb@math.ntnu.no`

SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS, LS2 9JT, UK

E-mail address: `marsh@maths.leeds.ac.uk`