

# REFLECTION GROUP PRESENTATIONS ARISING FROM CLUSTER ALGEBRAS

MICHAEL BAROT AND ROBERT J. MARSH

ABSTRACT. We give a presentation of a finite crystallographic reflection group in terms of an arbitrary seed in the corresponding cluster algebra of finite type and interpret the presentation in terms of companion bases in the associated root system.

## INTRODUCTION

The article [7] proved that cluster algebras of finite type can be classified by the Dynkin diagrams. A matrix (the *Cartan counterpart*) is associated to each seed in such a cluster algebra, and the cluster algebra associated to a Dynkin diagram  $\Delta$  is characterised by the fact that it contains seeds whose Cartan counterparts coincide with the Cartan matrix of  $\Delta$  (up to simultaneous permutation of the rows and columns).

It is well known that the Dynkin diagrams also classify the finite crystallographic reflection groups (see e.g. [8]). Our main result is a presentation of the reflection group associated to a Dynkin diagram in terms of an arbitrary seed in the corresponding cluster algebra. The presentation is given in terms of an edge-weighted oriented graph, known as a *diagram*, which is associated by Fomin and Zelevinsky to the seed.

If the Cartan counterpart of the seed is the Cartan matrix, then our presentation reduces to the usual Coxeter presentation of the reflection group. However, in general, the diagram of a seed contains cycles, and the presentation includes extra relations arising from the chordless cycles in the diagram (i.e. induced subgraphs which are cycles).

---

*Date:* 10 December 2011.

*2010 Mathematics Subject Classification.* Primary: 13F60, 20F55, 51F15; Secondary: 16G20.

*Key words and phrases.* Reflection group, Weyl group, finite type, Dynkin diagram, cluster algebra, companion basis, quasi-Cartan companion, mutation, diagram, presentation, cycle, Coxeter graph.

This work was supported by DGAPA, Universidad Nacional Autónoma de México, the Engineering and Physical Sciences Research Council [grant number EP/G007497/1] and the Institute for Mathematical Research (FIM, Forschungsinstitut für Mathematik) at the ETH, Zürich.

More precisely, if  $\Gamma$  is the diagram associated to a seed in a cluster algebra of finite type, we define, for vertices  $i, j$  of  $\Gamma$ ,

$$m_{ij} = \begin{cases} 2 & \text{if } i \text{ and } j \text{ are not connected;} \\ 3 & \text{if } i \text{ and } j \text{ are connected by an edge of weight 1;} \\ 4 & \text{if } i \text{ and } j \text{ are connected by an edge of weight 2;} \\ 6 & \text{if } i \text{ and } j \text{ are connected by an edge of weight 3.} \end{cases}$$

Let  $W(\Gamma)$  to be the group with generators  $s_i$ ,  $i$  a vertex of  $\Gamma$ , subject to the following relations (where  $e$  denotes the identity element):

- (1)  $s_i^2 = e$  for all  $i$ ;
- (2)  $(s_i s_j)^{m_{ij}} = e$  for all  $i \neq j$ ;
- (3) For any chordless cycle  $C$  in  $\Gamma$ :

$$i_0 \xrightarrow{w_1} i_1 \xrightarrow{w_2} \cdots \rightarrow i_{d-1} \xrightarrow{w_0} i_0,$$

where either all of the weights are 1, or  $w_0 = 2$ , we have:

$$(s_{i_0} s_{i_1} \cdots s_{i_{d-2}} s_{i_{d-1}} s_{i_{d-2}} \cdots s_{i_1})^2 = e.$$

**Theorem A** *Let  $\Gamma$  be the diagram associated to a seed in a cluster algebra of finite type. Then  $W(\Gamma)$  is isomorphic to the corresponding reflection group.*

This work was mainly motivated by the notions of quasi-Cartan companion [1] and companion basis [10, Defn. 4.1] (see also [9]). A companion basis for a skew-symmetric matrix  $B$  arising in a seed of the cluster algebra is a set  $S$  of roots in the corresponding root system which form a basis for the root lattice and for which the matrix  $A$  whose entries are  $(\alpha, \beta^\vee)$  for  $\alpha, \beta \in S$  is a positive quasi-Cartan companion matrix for  $B$  in the sense of [1]. This means that the diagonal entries are equal to 2, that the values of the off-diagonal entries in  $A$  coincide with the corresponding entries of  $B$  up to sign, and that the symmetrisation of the matrix is positive definite. Quasi-Cartan companions are used in [1] to give criteria for whether a cluster algebra is of finite type and companion bases were used in [9, 10] to describe the dimension vectors of indecomposable modules over cluster-tilted algebras in type  $A$  (dimension vectors of indecomposable modules over cluster-tilted algebras in the general concealed case were also independently calculated later in [11]).

The reflections corresponding to the elements of a companion basis associated to a seed in a finite type cluster algebra generate the corresponding reflection group. We show that the generators in a presentation of the reflection group as considered here can be regarded as this set of reflections. This perspective also allows us to show that, in the simply-laced case, the presentations obtained here coincide with root basis presentations of crystallographic reflection groups (amongst others) which were obtained in [5] using signed graphs.

We were also motivated by the article [2], which gives a presentation of a complex semisimple Lie algebra of simply-laced Dynkin type  $\Delta$  in terms of any positive definite unit form of type  $\Delta$  (i.e. equivalent to the unit form associated to  $\Delta$ ). This presentation also contains relations arising from chordless cycles, in this case arising in the bigraph describing the unit form.

The article is organised as follows. In Section 1, we recall some of the theory of cluster algebras of finite type from [7]. In Section 2, we collect together some of the properties of the seeds of such cluster algebras. In particular, we consider the diagrams associated to such seeds in [7] and the way in which they mutate, as well as the oriented cycles appearing in them. In Section 3 we associate a group to an arbitrary seed in the cluster algebra, using generators and relations. In Section 4 we find an efficient subset of the relations sufficient to define the group (this is the set of relations given above). In Section 5 we show that the group defined is invariant (up to isomorphism) under mutation of the seed. Since the group associated to a seed whose Cartan counterpart is a Cartan matrix is the reflection group corresponding to the cluster algebra, it follows that the group associated to any seed is isomorphic to this reflection group. In Section 6 we explain the connections with quasi-Cartan companion matrices and companion bases and with the extended Coxeter groups associated to signed graphs in [5].

## 1. CLUSTER ALGEBRAS OF FINITE TYPE

We first recall the definition of a (skew-symmetrisable) *cluster algebra* without coefficients, as defined by Fomin-Zelevinsky [6].

Let  $\mathbb{F} = \mathbb{Q}(u_1, u_2, \dots, u_n)$  be the field of rational functions in  $n$  indeterminates over  $\mathbb{Q}$ . Recall that a square matrix  $B$  is said to be *skew-symmetrisable* if there is a diagonal matrix  $D$  of the same size with positive diagonal entries such that  $DB$  is skew-symmetric. We consider pairs  $(\mathbf{x}, B)$ , where  $\mathbf{x} = \{x_1, x_2, \dots, x_n\}$  is a free generating set of  $\mathbb{F}$  and  $B$  is an  $n \times n$  skew-symmetrisable matrix.

Given  $k \in [1, n] = \{1, 2, \dots, n\}$ , a new such pair  $\mu_k(\mathbf{x}, B) = (\mathbf{x}', B')$ , the *mutation* of  $(\mathbf{x}, B)$  at  $k$ , is defined as follows. We have:

$$B'_{ij} = \begin{cases} -B_{ij} & \text{if } i = k \text{ or } j = k; \\ B_{ij} + \frac{|B_{ik}|B_{kj} + B_{ik}|B_{kj}|}{2} & \text{otherwise.} \end{cases}$$

Define an element  $x'_k \in \mathbb{F}$  by the *exchange relation*

$$x'_k x_k = \prod_{B_{ik} > 0} x_i^{B_{ik}} + \prod_{B_{ik} < 0} x_i^{-B_{ik}},$$

and set  $\mathbf{x}' = \{x_1, x_2, \dots, x_{k-1}, x'_k, x_{k+1}, \dots, x_n\}$ . We write  $\mu_k(B) = B'$ .

Now we fix a pair  $(\mathbf{x}, B)$ , the *initial seed*. We shall call any pair that can be obtained from  $(\mathbf{x}, B)$  by a sequence of mutations a *seed*. Two seeds are regarded as the same if there is a permutation of  $[1, n]$  which takes the matrix in the first seed to the second (when regarded as a simultaneous permutation of the rows and columns) and the free generating set in the first to the free generating set in the second. Note that mutating twice at  $k$  sends a seed to itself.

Each tuple  $\mathbf{x} = \{x_1, \dots, x_n\}$  such that there exists a seed  $(\mathbf{x}, B)$  is called a *cluster* and its entries  $x_i$  are called *cluster variables*. The cluster algebra  $\mathcal{A}(\mathbf{x}, B)$  is the  $\mathbb{Q}$ -subalgebra of  $\mathbb{F}$  generated by the set of all cluster variables. It does not depend (up to strong isomorphism, i.e. isomorphism of cluster algebras) on the choice of free generating set in the initial seed so we shall write  $\mathcal{A}(B)$  for  $\mathcal{A}(\mathbf{x}, B)$ .

A cluster algebra is said to be of *finite type* if it has finitely many seeds. Given a skew-symmetrisable matrix  $B$ , its *Cartan counterpart*  $A(B)$  is a matrix of the same size defined by

$$A_{ij} = \begin{cases} 2 & \text{if } i = j; \\ -|B_{ij}| & \text{otherwise.} \end{cases}$$

We recall a special case of a result of Fomin-Zelevinsky. Note that the original result deals with more general cluster algebras (with coefficients) and allows more general matrices (sign-skew-symmetric matrices) in the seeds.

**Theorem 1.1.** [7, 1.4] *There is a canonical bijection between the strong isomorphism classes of skew-symmetrisable cluster algebras of finite type and the Cartan matrices of finite type. A cluster algebra corresponds to a given Cartan matrix  $A$  provided it contains a seed for which the Cartan counterpart of the matrix is  $A$ .*

A skew-symmetrisable matrix  $B$  is said to be *2-finite* provided  $|B_{ij}B_{ji}| \leq 3$  for all  $i, j \in [1, n]$ . Note that this means the opposite pair of entries  $(B_{ij}, B_{ji})$  in  $B$  must be one of  $(1, 1)$ ,  $(1, 2)$ ,  $(2, 1)$ ,  $(1, 3)$  or  $(3, 1)$ . Then Fomin-Zelevinsky also show that:

**Theorem 1.2.** [7, 1.8] *A cluster algebra  $\mathcal{A}$  is of finite type if and only if all the matrices appearing in the seeds of  $\mathcal{A}$  are 2-finite.*

As observed by Fomin-Zelevinsky [7, §9].

**Corollary 1.3.** *Let  $B$  be a 2-finite matrix. Then any square submatrix of  $B$ , obtained by taking a subset of the rows and the corresponding subset of the columns, is again 2-finite. Thus, passing to such a submatrix preserves the fact that the corresponding cluster algebra is of finite type.*

Fomin-Zelevinsky [7, 7.3] define the *diagram*  $\Gamma(B)$  of a skew-symmetrisable matrix  $B$  to be the weighted oriented graph with vertices  $[1, n]$ . There is an arrow from  $i$  to  $j$  if and only if  $B_{ij} > 0$ , weighted with the positive integer  $|B_{ij}B_{ji}|$ . It is useful to regard the situation of having no arrow between two vertices as an arrow of weight zero. By [7, 7.5], all 3-cycles in the underlying unoriented graph of  $\Gamma(B)$  are oriented cyclically. In the 2-finite case, the mutation rule for diagrams is as follows:

**Proposition 1.4.** [7, Lemma 8.5] *Let  $B$  be a 2-finite skew-symmetrisable matrix. Then  $\Gamma(\mu_k(B))$  is uniquely determined by  $\Gamma(B)$  as follows:*

- (a) *Reverse the orientations of all edges in  $\Gamma(B)$  incident with  $k$  (leaving the weights unchanged).*
- (b) *For any path in  $\Gamma(B)$  of form  $i \rightarrow k \rightarrow j$ , let  $c$  be the weight on the edge  $j \rightarrow i$  (possibly zero). Let  $c'$  be determined by*

$$\sqrt{c} + \sqrt{c'} = \sqrt{ab}.$$

*Then  $\Gamma(B)$  changes as in Figure 1.*

We call  $\Gamma(\mu_k(B))$  the *mutation* of  $\Gamma$  at  $B$  and use the notation  $\mu_k(\Gamma)$ .

We say that a skew-symmetrisable matrix  $B$  (and the corresponding diagram) is of *finite type* if the corresponding cluster algebra is of finite type. Hence, by Corollary 1.3, we have:



FIGURE 1. Mutation of a diagram.

**Corollary 1.5.** [7, §9] *Let  $\Gamma$  be a diagram of finite type. Then any induced subdiagram of  $\Gamma$  is again of finite type.*

## 2. DIAGRAMS OF FINITE TYPE

In this section, we collect together some properties of diagrams of finite type. We call an induced subgraph of an unoriented graph which is a cycle a *chordless cycle*. We first recall:

**Proposition 2.1.** [7, Prop. 9.7] *Let  $\Gamma$  be a diagram of finite type. Then a chordless cycle in the underlying unoriented graph of  $\Gamma$  is always cyclically oriented in  $\Gamma$ . In addition, the unoriented graph underlying the cycle must be one of those in Figure 2.*

We will refer to such cycles as chordless cycles in  $\Gamma$ .




FIGURE 2. The chordless cycles in a diagram of finite type.




FIGURE 3. 3-vertex connected subdiagrams of diagrams of finite type.

**Lemma 2.2.** *Let  $\Gamma$  be a diagram of finite type and let  $\Xi$  be an induced subdiagram with three vertices which is connected. Then the unoriented graph underlying  $\Xi$  must be one of those in Figure 3.*

*Proof.* This follows from Proposition 2.1, together with the fact that a diagram of form  $1 \rightarrow 2 \rightarrow 3$  (with positive weights) cannot have a weight greater than 2 or two weights equal to 2, by [7, Prop. 9.3] and its proof.  $\square$

**Corollary 2.3.** *Let  $\Gamma$  be a graph of finite type and suppose that are 3 distinct vertices  $i, j, k$  of  $\Gamma$  with both  $i$  and  $j$  connected to  $k$ . Then the effect of the mutation of  $\Gamma$  at  $k$  on the induced subdiagram must be as in Figure 4 (starting on one side or the other), up to switching  $i$  and  $j$ .*

*Proof.* This follows from Lemma 2.2 and Proposition 1.4.  $\square$




FIGURE 4. Local picture of mutation of a diagram of finite type.

We say that a vertex in a diagram  $\Gamma$  is *connected* to another if there is an edge between them. The next result describes the way vertices in  $\Gamma$  can be connected to a chordless cycle.

**Lemma 2.4.** *Suppose that a vertex  $k$  is connected to at least one vertex in a chordless cycle in a diagram  $\Gamma$  of finite type. Then  $k$  is connected to at most two vertices of the cycle. If it is connected to two vertices, they must be adjacent in the cycle.*

*Proof.* Consider a chordless cycle  $C$ :

$$i_0 \rightarrow i_1 \rightarrow \cdots \rightarrow i_{r-1} \rightarrow i_0$$

in  $\Gamma$ , and suppose that  $k$  is connected to  $i_{a_1}, i_{a_2}, \dots, i_{a_t}$  for some  $t \leq r$ , where  $0 \leq a_1 < a_2 < \cdots < a_t \leq r - 1$ . Suppose first that  $t > 2$ . Since, by Proposition 2.1, the chordless cycles (in the underlying unoriented graph)

$$i_{a_1} \text{ --- } i_{a_1+1} \text{ --- } \cdots \quad i_{a_2} \text{ --- } k \text{ --- } i_{a_1}$$

and

$$i_{a_2} \text{ --- } i_{a_2+1} \text{ --- } \cdots \quad i_{a_3} \text{ --- } k \text{ --- } i_{a_2}$$

are cyclically oriented in  $\Gamma$ , we must have

$$i_{a_2} \rightarrow k \rightarrow i_{a_1}$$

from the first and

$$i_{a_3} \rightarrow k \rightarrow i_{a_2}$$

from the second, giving a contradiction.

If  $r = 2$ , consider the cycle

$$i_{a_2} \text{ --- } i_{a_2+1} \text{ --- } \cdots \quad i_{a_1} \text{ --- } k \text{ --- } i_{a_2}$$

(where the subscripts are written mod  $r$ ). If it is a chordless cycle, it must be oriented, giving

$$i_{a_1} \rightarrow k \rightarrow i_{a_2},$$

again giving a contradiction. Hence this last cycle is not a chordless cycle, which implies that  $i_{a_1}$  and  $i_{a_2}$  are adjacent in  $C$  as required.  $\square$

**Lemma 2.5.** *Let  $\Gamma$  be a diagram of finite type and  $\Gamma' = \mu_k(\Gamma)$  the mutation of  $\Gamma$  at vertex  $k$ . In Figures 5 and 6 we list various types of induced subdiagrams in  $\Gamma$  (on the left) and corresponding cycles  $C'$  in  $\Gamma'$  (on the right) arising from the mutation of  $\Gamma$  at  $k$ . The diagrams are drawn so that  $C'$  is always a clockwise cycle, and in case (i),  $C'$  has at least 3 vertices, while in case (j),  $C'$  has at least 4 vertices. Then every chordless cycle in  $\Gamma'$  arises in such a way.*





FIGURE 5. Induced subdiagrams of  $\Gamma$  and the corresponding chordless cycles in  $\Gamma' = \mu_k(\Gamma)$ , part 1.

*Proof.* Let  $C'$  be a chordless cycle in  $\Gamma'$ . We consider the different possibilities for the vertex  $k$  in relation to  $C'$ . If  $k$  is a vertex of  $C'$ , then it follows from Proposition 2.1 that  $C'$  arises as in case (a)-(f) or (j). If  $k$  does not lie in  $C'$  and is not connected to any vertex of  $C'$  then  $C'$  arises as in case (k). If  $k$  does not lie in  $C'$  and is connected



- (k)  $C$  is an oriented cycle in  $\Gamma$  not connected to  $k$ . Then  $C'$  is the corresponding cycle in  $\Gamma'$ .
- (l)  $C$  is an oriented cycle in  $\Gamma$  with exactly one vertex connected to  $k$  (via an edge of unspecified weight). Then  $C'$  is the corresponding cycle in  $\Gamma'$ .

FIGURE 6. Induced subdiagrams of  $\Gamma$  and the corresponding chordless cycles in  $\Gamma' = \mu_k(\Gamma)$ , part 2.

to exactly one vertex of  $C'$ , then  $C'$  arises as in case (l). Note that the edge linking  $k$  to  $C'$  must have weight 1 or 2 (since a diagram with two edges with weights 3 and  $a = 1, 2$  or 3 is not of finite type; see the proof of [7, Prop. 9.3]).

By Lemma 2.4, the only remaining case is where  $k$  does not lie in  $C'$  and  $k$  is connected to two adjacent vertices of  $C'$ . If the 3-cycle  $C_k$  containing these 3 vertices in  $\Gamma'$  has all of its weights equal to 1, then  $C'$  also has this property, since the vertices in  $C_k \cup C'$  form a chordless cycle in  $\Gamma$  containing two consecutive edges of weight 1. Then  $C'$  is as in case (i).

If  $C_k$  has weights 1, 2, 2 then it is not possible for the edge of weight 1 to be shared with  $C'$ , since then in  $\Gamma$ , the vertices of  $C'$  form a chordless cycle which is not oriented. If the common edge has weight 2 and  $C'$  has more than 3 vertices, then the vertices of  $C_k \cup C'$  form a chordless cycle in  $\Gamma$  with at least 5 vertices and at least one edge of weight 2, contradicting Proposition 2.1. The only other possibility is when  $C'$  has exactly 3 vertices, and then, by Proposition 2.1, it must arise as in (g) or (h). The result is proved.  $\square$

### 3. THE GROUP OF A DIAGRAM ARISING IN A CLUSTER ALGEBRA FINITE TYPE

In this section we define a group (by generators and relations) for any diagram  $\Gamma$  of finite type. We shall see later that it is isomorphic to the reflection group of the same Dynkin type as the cluster algebra in which  $\Gamma$  arises.

Let  $\Gamma$  be a diagram of finite type. For vertices  $i, j$  of  $\Gamma$  define

$$m_{ij} = \begin{cases} 2 & \text{if } i \text{ and } j \text{ are not connected;} \\ 3 & \text{if } i \text{ and } j \text{ are connected by an edge of weight 1;} \\ 4 & \text{if } i \text{ and } j \text{ are connected by an edge of weight 2;} \\ 6 & \text{if } i \text{ and } j \text{ are connected by an edge of weight 3.} \end{cases}$$

Then we define  $W_\Gamma$  to be the group with generators  $s_i$ ,  $i = 1, 2, \dots, n$ , subject to the following relations:

- (R1)  $s_i^2 = e$  for all  $i$ ,
- (R2)  $(s_i s_j)^{m_{ij}} = e$  for all  $i \neq j$ ,

where  $e$  denotes the identity element of  $W_\Gamma$ .

Note that, in the presence of (R1), relation (R2) is symmetric in that  $(s_j s_i)^{m_{ji}} = e$  follows from  $(s_i s_j)^{m_{ij}} = e$ , since  $m_{ij}$  is symmetric in  $i$  and  $j$ . We shall usually use this fact without comment.

For the remaining relations, we suppose that  $C$ , given by:

$$i_0 \rightarrow i_1 \rightarrow \dots \rightarrow i_{d-1} \rightarrow i_0$$

is a chordless cycle in  $\Gamma$ . For  $a = 0, 1, 2, \dots, d-1$ , define (with subscripts modulo  $d$ ):

$$r(i_a, i_{a+1}) = s_{i_a} s_{i_{a+1}} \cdots s_{i_{a+d-1}} s_{i_{a+d-2}} \cdots s_{i_{a+1}}$$

and

$$r(i_a, i_{a-1}) = s_{i_a} s_{i_{a-1}} \cdots s_{i_{a-(d-1)}} s_{i_{a-(d-2)}} \cdots s_{i_{a-1}}$$

- (R3)(a) If all of the weights in the edges of  $C$  are equal to 1, then we set  $r(i_a, i_{a+1})^2 = e$  and  $r(i_a, i_{a-1})^2 = e$  for  $a = 0, 1, \dots, d-1$ .
- (R3)(b) If  $C$  has some edges of weight 2, then set  $r(i_a, i_{a+1})^k = e$  where  $k = 4 - w_a$  and  $w_a$  is the weight of the edge between  $i_a$  and  $i_{a+1}$ . Similarly, we set  $r(i_a, i_{a-1})^k = e$  where  $k = 4 - w'_a$  and  $w'_a$  is the weight of the edge between  $i_a$  and  $i_{a-1}$ .

Note that if we think of  $r(i_a, i_{a+1})$  (respectively,  $r(i_a, i_{a-1})$ ) as a path in  $\Gamma$ , then in (R3)(b),  $w_a$  (respectively,  $w'_a$ ) is the weight of the only edge in  $C$  not used in the path.

**Remark 3.1.** We note that if  $\Gamma$  is the diagram associated to a matrix whose Cartan counterpart is the Cartan matrix of a Dynkin diagram  $\Delta$ , then  $W_\Gamma$  is the reflection group of type  $\Delta$ . In this case, the relations (R3)(a) and (R3)(b) do not occur, since  $\Gamma$  is a tree.

**Remark 3.2.** We also note that, as in Coxeter groups, if  $i \neq j$  then

$$s_i s_j s_i \cdots = s_j s_i s_j \cdots,$$

where there are  $m_{ij}$  terms on each side of the equation. In particular, if there is no edge in  $\Gamma$  between  $i$  and  $j$ , then  $s_i s_j = s_j s_i$ , and if there is an edge in  $\Gamma$  between  $i$  and  $j$  with weight 1, then  $s_i s_j s_i = s_j s_i s_j$ . We shall use these relations liberally.

**Remark 3.3.** We shall see later (using the companion basis perspective) that, in the simply-laced case, the group  $W_\Gamma$  coincides with a group defined in [5, Defn. 6.2]. See the end of Section 6. In this context, the elements  $r(i_a, i_{a+1})^2$  are referred to as cut elements.

**Example 3.4.** We consider the example in Figure 7, which is the diagram of a seed of type  $D_4$ .




FIGURE 7. A diagram of type  $D_4$ .

The relations in this case are:

- (R1)  $s_i^2 = e$  for  $i = 1, 2, 3, 4$ ;
- (R2)  $(s_1 s_2)^3 = (s_2 s_3)^3 = (s_3 s_4)^3 = (s_4 s_1)^3 = e$  and  $(s_1 s_3)^2 = (s_2 s_4)^2 = e$ ;
- (R3) For each element  $r(i, j)$  of the following list we have  $r(i, j)^2 = e$ .

$$r(1, 2) = s_1 s_2 s_3 s_4 s_3 s_2;$$

$$r(2, 3) = s_2 s_3 s_4 s_1 s_4 s_3;$$

$$r(3, 4) = s_3 s_4 s_1 s_2 s_1 s_4;$$

$$r(4, 1) = s_4 s_1 s_2 s_3 s_2 s_1;$$

$$r(2, 1) = s_2 s_1 s_4 s_3 s_4 s_1;$$

$$r(3, 2) = s_3 s_2 s_1 s_4 s_1 s_2;$$

$$r(4, 3) = s_4 s_3 s_2 s_1 s_2 s_3;$$

$$r(1, 4) = s_1 s_4 s_3 s_2 s_3 s_4.$$

We shall see in the next section that the relations in (R3) can be reduced (in fact, it is enough to choose just one of them).

#### 4. CYCLIC SYMMETRY

It turns out that, in the presence of the relations (R1) and (R2) in the above, many of the relations in (R3)(a) and (b) are redundant. In the sequel, we shall usually use (R1) without comment.

**Lemma 4.1.** Let  $\Gamma$  be a diagram of finite type containing a chordless cycle  $C$ :

$$i_0 \rightarrow i_1 \rightarrow \cdots \rightarrow i_{d-1} \rightarrow i_0$$

with the weights of all of its edges equal to 1. Let  $W$  be the group with generators  $s_1, s_2, \dots, s_n$  subject to relations (R1), (R2) and  $r(i_a, i_{a+1})^2 = e$  or  $r(i_a, i_{a-1})^2 = e$  for a fixed value of  $a$ . Then all of the relations in (R3)(a) hold for  $C$ .

*Proof.* Note first that it is enough to consider the case  $a = 0$ . Furthermore, we consider only the first case; the second case is similar. For simplicity of notation, we renumber the vertices of  $C$  so that  $i_j = j$  for  $j = 0, 1, \dots, a-1$ . Thus we assume that  $r(0, 1)^2 = e$  in  $W$ .

We have, using (R2):

$$\begin{aligned}
r(d-1, 0) &= s_{d-1}s_0s_1 \cdots s_{d-3}s_{d-2}s_{d-3} \cdots s_1s_0 \\
&= s_0s_0s_{d-1}s_0s_1 \cdots s_{d-3}s_{d-2}s_{d-3} \cdots s_1s_{d-1}s_{d-1}s_0 \\
&= s_0s_{d-1}s_0s_{d-1}s_1 \cdots s_{d-3}s_{d-2}s_{d-3} \cdots s_1s_{d-1}s_{d-1}s_0 \\
&= s_0s_{d-1}(s_0s_1 \cdots s_{d-3}s_{d-1}s_{d-2}s_{d-1}s_{d-3} \cdots s_1)s_{d-1}s_0 \\
&= s_0s_{d-1}(s_0s_1 \cdots s_{d-3}s_{d-2}s_{d-1}s_{d-2}s_{d-3} \cdots s_1)s_{d-1}s_0 \\
&= s_0s_{d-1}r(0, 1)s_{d-1}s_0.
\end{aligned}$$

It follows that  $r(d-1, 0)^2 = e$ . Repeating this argument inductively, we see that  $r(a, a+1)^2 = e$  for all  $a$ . Furthermore,

$$\begin{aligned}
r(d-1, d-2) &= s_{d-1}s_{d-2} \cdots s_1s_0s_1 \cdots s_{d-3}s_{d-2} \\
&= s_{d-1}s_{d-2} \cdots s_1(s_0s_1 \cdots s_{d-3}s_{d-2}s_{d-1}s_{d-2} \cdots s_1)s_1s_2 \cdots s_{d-1} \\
&= s_{d-1}s_{d-2} \cdots s_1r(0, 1)s_1s_2 \cdots s_{d-1}.
\end{aligned}$$

So  $r(d-1, d-2)^2 = e$ . A similar argument to the above then shows that  $r(a, a-1)^2 = e$  for all  $a$ .  $\square$

**Lemma 4.2.** *Let  $\Gamma$  be a diagram of finite type containing a 3-cycle  $C$  as in Figure 8 (where we number the vertices 1, 2, 3 for convenience). Let  $W$  be the group with generators  $s_1, s_2, \dots, s_n$  subject to relations (R1) and (R2) arising from the diagram  $\Gamma$ . Then the following are equivalent:*

- (a)  $r(1, 2)^2 = e$ ;
- (b)  $r(3, 2)^2 = e$ ;
- (c)  $r(3, 1)^2 = e$ ;
- (d)  $r(2, 1)^2 = e$ .

Furthermore, if one of the above holds, then the following both hold:

- (e)  $r(1, 3)^3 = e$ ;
- (f)  $r(2, 3)^3 = e$ .




FIGURE 8. A 3-cycle (see Lemma 4.2).

*Proof.* Note that  $r(1, 2) = s_1s_2s_3s_2$ ,  $r(2, 1) = s_2s_1s_3s_1$ ,  $r(1, 3) = s_1s_3s_2s_3$ ,  $r(3, 1) = s_3s_1s_2s_1$ ,  $r(2, 3) = s_2s_3s_1s_3$  and  $r(3, 2) = s_3s_2s_1s_2$ .

(a) implies (b): This follows from:

$$r(3, 2) = (s_3s_2)r(1, 2)(s_2s_3).$$

(b) implies (c): Using (R2) we have:

$$r(3, 1) = s_3s_1s_2s_1 = s_3s_2s_1s_2 = r(3, 2).$$

(c) implies (d): This follows from:

$$r(2, 1) = (s_1s_3)r(3, 1)(s_3s_1).$$

(d) implies (a): Using (R2) we have:

$$(s_1s_2)r(1, 2)(s_2s_1) = s_1s_2s_1s_2s_3s_1 = s_1s_1s_2s_1s_3s_1 = s_2s_1s_3s_1 = r(2, 1)$$

which implies the result.

Now assume that one of (a)–(d) holds (and thus that all of them hold). To prove (e), we know that  $r(3, 1)^2 = e$  by (c), so

$$s_3s_1s_2s_1s_3s_1s_2s_1 = e.$$

Hence,

$$(1) \quad s_3s_1s_2s_1s_3 = s_1s_2s_1.$$

Thus, noting that  $s_3s_1s_3 = s_1s_3s_1s_3s_1$  by (R2),

$$\begin{aligned} s_3r(1, 3)^3s_3 &= s_3s_1s_3s_2s_3s_1s_3s_2s_3s_1s_3s_2 \\ &= s_1s_3s_1s_3s_1s_2s_1s_3s_1s_3s_1s_2s_1s_3s_1s_3s_1s_2. \end{aligned}$$

Hence, using equation (1) and (R2), we have:

$$\begin{aligned} s_3s_1s_3r(1, 3)^3s_3s_1s_3 &= s_1(s_3s_1s_2s_1s_3)s_1(s_3s_1s_2s_1s_3)s_1(s_3s_1s_2s_1s_3) \\ &= s_1(s_1s_2s_1)s_1(s_1s_2s_1)s_1(s_1s_2s_1) \\ &= (s_2s_1)^3 = e. \end{aligned}$$

The proof of (f) (given (a)–(d)) is the same except that the roles of  $s_1$  and  $s_2$  are exchanged.  $\square$

Note that it is not claimed that (e) and (f) are equivalent to (a),(b),(c) and (d) (we do not know if this holds, but it seems to be unlikely).

**Example 4.3.** *Lemma 4.2 implies that in Example 3.4, we can replace the relations in (R3) with any fixed one, e.g.*

$$r(1, 2)^2 = (s_1s_2s_3s_4s_3s_2)^2 = e.$$

**Lemma 4.4.** *Let  $\Gamma$  be a diagram of finite type containing a chordless 4-cycle  $C$  as in Figure 9 (where we number the vertices 1, 2, 3, 4 for convenience). Let  $W$  be the group with generators  $s_1, s_2, \dots, s_n$  subject to relations (R1) and (R2) arising from the diagram  $\Gamma$ . Then the following are equivalent:*

- (a)  $r(1, 2)^2 = e$ ;
- (b)  $r(2, 1)^2 = e$ ;

- (c)  $r(3, 4)^2 = e$ ;
- (d)  $r(4, 3)^2 = e$ .

Furthermore, if one of the above holds, then the following all hold:

- (e)  $r(2, 3)^3 = e$ ;
- (f)  $r(1, 4)^3 = e$ ;
- (g)  $r(4, 1)^3 = e$ .
- (h)  $r(3, 2)^3 = e$ ;




FIGURE 9. A chordless 4-cycle (see Lemma 4.4).

*Proof.* Note first that  $r(1, 2) = s_1 s_2 s_3 s_4 s_3 s_2$ ,  $r(2, 1) = s_2 s_1 s_4 s_3 s_4 s_1$ ,  $r(2, 3) = s_2 s_3 s_4 s_1 s_4 s_3$ ,  $r(3, 2) = s_3 s_2 s_1 s_4 s_1 s_2$ ,  $r(3, 4) = s_3 s_4 s_1 s_2 s_1 s_4$ ,  $r(4, 3) = s_4 s_3 s_2 s_1 s_2 s_3$ ,  $r(1, 4) = s_1 s_4 s_3 s_2 s_3 s_4$  and  $r(4, 1) = s_4 s_1 s_2 s_3 s_2 s_1$ .

(a) implies (b): Using (R2), we have:

$$\begin{aligned}
 r(2, 1) &= s_2 s_1 s_4 s_3 s_4 s_1 \\
 &= s_1 (s_1 s_2 s_1) (s_4 s_3 s_4) s_1 \\
 &= s_1 (s_2 s_1 s_2) (s_3 s_4 s_3) s_2 s_2 s_1 \\
 &= (s_1 s_2) (s_1 s_2 s_3 s_4 s_3 s_2) (s_2 s_1) \\
 &= (s_1 s_2) r(1, 2) (s_2 s_1).
 \end{aligned}$$

(b) implies (c): We have:

$$\begin{aligned}
 r(3, 4) &= s_3 s_4 s_1 s_2 s_1 s_4 \\
 &= s_3 s_4 s_1 (s_2 s_1 s_4 s_3 s_4 s_1) s_1 s_4 s_3 \\
 &= s_3 s_4 s_1 r(2, 1) s_1 s_4 s_3.
 \end{aligned}$$

The arguments for (c) implies (d) and (d) implies (a) are similar to those for (a) implies (b) and (b) implies (c), respectively, using the symmetry of the 4-cycle. Suppose that one of (a)–(d) holds. Then (a) holds, so

$$(s_1 s_2 s_3 s_4 s_3 s_2)^2 = e.$$

Hence, using (R2), we have:

$$(s_1 s_2 s_4 s_3 s_4 s_2)^2 = e.$$

Thus, noting that  $s_2$  and  $s_4$  commute, we have:

$$(s_1 s_4 s_2 s_3 s_2 s_4)^2 = e.$$

So we get  $(s_4 s_1 s_4 s_2 s_3 s_2)^2 = e$  and hence

$$(2) \quad s_4 s_1 s_4 s_2 s_3 s_2 = s_2 s_3 s_2 s_4 s_1 s_4.$$

Therefore, using first  $s_3s_2s_3 = s_2s_3s_2s_3s_2$  and then equation (2), we get

$$\begin{aligned} s_3r(2,3)s_3 &= s_3s_2s_3s_4s_1s_4 \\ &= s_2s_3(s_2s_3s_2s_4s_1s_4) \\ &= s_2s_3(s_4s_1s_4s_2s_3s_2) \\ &= s_2s_3s_4(s_1s_2)s_4s_3s_2, \end{aligned}$$

where in the last equation we used that  $s_2$  and  $s_4$  commute. Consequently

$$\begin{aligned} (s_3r(2,3)s_3)^3 &= s_2s_3s_4(s_1s_2)^3s_4s_3s_2 \\ &= e, \end{aligned}$$

since  $(s_1s_2)^3 = e$ . This shows that  $r(2,3)^3 = e$ .

By symmetrical arguments, (b) implies (f), (c) implies (g) and (d) implies (h) (noting that the orientation of the edges in the 4-cycle does not play a role in the argument).  $\square$

**Corollary 4.5.** *Let  $\Gamma$  be a diagram of finite type. Then, in the presence of relations (R1) and (R2), the relations (3) considered in the introduction imply the relations (R3) considered above. In particular,  $W_\Gamma$  is isomorphic to the group  $W(\Gamma)$  defined in the introduction.*

## 5. INVARIANCE OF THE GROUP UNDER MUTATION

In this section we prove the main result, that the group of a diagram of finite type depends only on the Dynkin type, up to isomorphism. We will prove this by showing that the group of the diagram is invariant (up to isomorphism) under mutation.

We fix a diagram  $\Gamma$  of finite type and a vertex  $k$  of  $\Gamma$ . Then we have the mutation  $\Gamma' = \mu_k(\Gamma)$ .

For a vertex  $i$  of  $\Gamma$ , we define an element  $t_i \in W_\Gamma$  as follows:

$$t_i = \begin{cases} s_k s_i s_k & \text{if there is an arrow } i \rightarrow k \text{ in } \Gamma \text{ (possibly weighted);} \\ s_i & \text{otherwise.} \end{cases}$$

Our aim is to show that the elements  $t_i$  satisfy the relations (R1)-(R3) defining  $W_{\Gamma'}$ . We denote the values of  $m_{ij}$  for  $\Gamma'$  by  $m'_{ij}$ . Note that (R1) is clear, since  $(s_k s_i s_k)^2 = s_k s_i^2 s_k = e$ , so it remains to check (R2) and (R3). We will use the previous section to reduce the number of checks that have to be done.

We first deal with some easy cases:

**Lemma 5.1.** *Let  $i, j$  be distinct vertices of  $\Gamma$ .*

- (a) *If  $i = k$  or  $j = k$  then  $(t_i t_j)^{m'_{ij}} = e$ .*
- (b) *If at most one of  $i, j$  is connected to  $k$  in  $\Gamma$  (or, equivalently, in  $\Gamma'$ ), then  $(t_i t_j)^{m'_{ij}} = e$ .*

*Proof.* For (a), suppose first that  $i = k$ . Note that  $m'_{ij} = m_{ij}$ . The only non-trivial case is thus when there is an arrow from  $j$  to  $k$ , so that  $t_j = s_k s_j s_k$ . But then  $t_i t_j = s_k s_k s_j s_k = s_j s_k$  and the result follows. The case when  $j = k$  is similar. For (b), suppose first that  $i$  is connected to  $k$ . The only non-trivial case is when the

edge between  $i$  and  $k$  is oriented towards  $k$ , so that  $t_i = s_k s_i s_k$ . Then  $t_i t_j = s_k s_i s_k s_j$  and the result follows from the fact that  $s_k s_j = s_j s_k$  (since  $j$  is not connected to  $k$ ). The case when  $j$  is connected to  $k$  is similar.  $\square$


**Proposition 5.2.** *The elements  $t_i$ , for  $i$  a vertex of  $\Gamma$ , satisfy the relations (R1)–(R3) defining  $W_{\Gamma'}$ .*

*Proof.* After Lemma 5.1, the remaining relations to check are the relations from (R2), that  $(t_i t_j)^{m'_{ij}} = e$  when both  $i$  and  $j$  are connected to  $k$ , and the relations (R3) corresponding to the chordless cycles in  $\Gamma'$ . For the former, we need to check that, in going (i) from left to right or (ii) from right to left in Corollary 2.3 (in each case, from  $\Gamma$  to  $\Gamma'$ ) we have that  $(t_i t_j)^{m'_{ij}} = e$ . (Note that it is enough to check this for the given labelling of  $i, j$ , since the relation for the pair  $(t_j, t_i)$  follows from the relation for  $(t_i, t_j)$ .) At the same time, we will check that the cycle relations in (R3) hold if any cycles are produced; we shall denote by  $r'(i, j)$ , etc., the elements in the cycle relations for  $W_{\Gamma'}$ . Note that by Lemmas 4.2 and 4.4, we only have to check one of the relations in the first part of each of those lemmas (i.e. in which the left hand side is a square).

- (a)(i) We have  $(t_i t_j)^2 = (s_k s_i s_k s_k s_j s_k)^2 = s_k (s_i s_j)^2 s_k = e$ .
- (a)(ii) We have  $(t_i t_j)^2 = (s_i s_j)^2 = e$ .
- (b)(i) We have:  $t_i t_j = s_k s_i s_k s_j = s_i s_k s_i s_j = s_i s_k s_j s_i$  and therefore  $s_i (t_i t_j)^3 s_i = (s_k s_j)^3 = e$ . And  $r'(j, k)^2 = (t_j t_k t_i t_k)^2 = (s_j s_k s_k s_i s_k s_k)^2 = (s_j s_i)^2 = e$ .
- (b)(ii) We have  $(t_i t_j)^2 = (s_i s_k s_j s_k)^2 = r(i, k)^2 = e$ .
- (c)(i) We have  $(t_i t_j)^2 = (s_k s_i s_k s_k s_j s_k)^2 = s_k (s_i s_j)^2 s_k = e$ .
- (c)(ii) We have  $(t_i t_j)^2 = (s_i s_j)^2 = e$ .
- (d)(i) We have:  $s_k (t_i t_j)^4 s_k = s_k (s_k s_i s_k s_j)^4 s_k = (s_i s_k s_j s_k)^4 = (s_i s_j s_k s_j)^4$  and therefore  $s_j s_k (t_i t_j)^4 s_k s_j = (s_j s_i s_j s_k)^4 = (s_i s_k)^4 = e$ . And  $r'(j, k) = (t_j t_k t_i t_k)^2 = (s_j s_k s_k s_i s_k s_k)^2 = (s_j s_i)^2 = e$ .
- (d)(ii) We have:  $(t_i t_j)^2 = (s_i s_k s_j s_k)^2 = r(i, k)^2 = e$ .
- (e)(i) We have:  $(t_i t_j)^4 = (s_k s_i s_k s_j)^4 = (s_i s_k s_i s_j)^4$  and therefore  $s_i (t_i t_j)^4 s_i = (s_k s_i s_j s_i)^4 = (s_k s_j)^4 = e$ . And  $r'(i, k) = t_i t_k t_j t_k = s_k s_i s_k s_k s_j s_k = s_k s_i s_j s_k$  and therefore  $s_k r'(i, k)^2 s_k = (s_i s_j)^2 = e$ .
- (e)(ii) We have  $(t_i t_j)^2 = (s_i s_k s_j s_k)^2 = r(i, k)^2 = e$ .
- (f)(i) We have:  $s_j (t_i t_j)^3 s_j = s_j (s_k s_i s_k s_j)^3 s_j = (s_j s_k s_i s_k)^3 = r(j, k)^3 = e$ . And  $s_k r'(i, j)^2 s_k = s_k (t_i t_j t_k t_j)^2 s_k = s_k (s_k s_i s_k s_j s_k s_j)^2 s_k = (s_i s_k s_j s_k s_j s_k)^2 = (s_i s_j s_k s_j)^2 = r(i, j)^2 = e$ .
- (f)(ii) The argument is the same as for (f)(i) by symmetry.


Finally, we need to check that the elements  $t_i$  satisfy the cycle relations (R3) for all the chordless cycles in  $\Gamma'$ . Note that Lemma 2.5 describes how the chordless cycles in  $\Gamma'$  arise from induced subdiagrams of  $\Gamma$ . In each case, we need to check that the corresponding cycle relations hold (using Lemmas 4.1, 4.2 and 4.4 to reduce the work). Note that, in the above, we have already checked the cases (a)–(d) in Lemma 2.5. We now check the remaining cases. We choose a labelling of the vertices in each case.

- (e) We label the vertices as follows.




We have  $r'(1, 2) = t_1 t_2 t_3 t_4 t_3 t_2 = s_1 s_1 s_2 s_1 s_3 s_4 s_3 s_1 s_2 s_1$  and therefore  $s_1 s_2 s_1 r'(1, 2) s_1 s_2 s_1 = (s_1 s_2 s_1 s_2 s_1) s_3 s_4 s_3 = s_2 s_3 s_4 s_3 = r(2, 3)$ . Hence  $r(2, 3)^2 = e$  implies  $r'(1, 2)^2 = e$ .

(f) We label the vertices as follows.




It follows from the commutativity of  $s_2$  and  $s_4$  that  $r'(1, 2) = t_1 t_2 t_3 t_4 t_3 t_2 = s_1 (s_2 s_2) s_3 (s_2 s_4 s_2) s_3 (s_2 s_2) = s_1 s_3 s_4 s_3 = r(1, 3)$ , hence  $r'(1, 2)^2 = r(1, 3)^2 = e$ .

(g) We label the vertices as follows.




We have  $r'(4, 3) = t_4 t_3 t_2 t_3 = s_1 s_4 s_1 s_3 s_2 s_3$ . Using that  $s_1$  and  $s_3$  commute we get  $s_1 r'(4, 3) s_1 = s_4 s_1 s_3 s_2 s_3 s_1 = s_4 s_3 (s_1 s_2 s_1) s_3 = s_4 s_3 s_2 s_1 s_2 s_3 = r(4, 3)$  and hence  $r(4, 3)^2 = e$  implies that  $r'(4, 3)^2 = e$ .

(h) We use the following labelling of the vertices.



We have  $r'(3, 4)^2 = (t_3 t_4 t_1 t_4)^2 = (s_3 s_4 s_2 s_1 s_2 s_4)^2 = (s_3 s_4 s_1 s_2 s_1 s_4)^2 = r(3, 4)^2 = e$ .


(i) We use the following labelling of the vertices.



Note that  $t_h = s_k s_h s_k$ , while  $t_i = s_i$  for all  $i \neq h$ .

$$\begin{aligned} r'(1, 2)^2 &= (t_1 t_2 \cdots t_{h-1} t_h t_{h-1} \cdots t_2)^2 \\ &= (s_1 s_2 \cdots s_{h-1} s_k s_h s_k s_{h-1} \cdots s_2)^2 \\ &= (s_1 s_2 \cdots s_{h-1} s_h s_k s_h s_{h-1} \cdots s_2)^2 = r(1, 2)^2 = e. \end{aligned}$$

(j) We label the vertices as follows.




Here we have  $t_1 = s_k s_1 s_k$  and  $t_i = s_i$  for all  $i \neq 1$ .

$$\begin{aligned} r'(h, h-1)^2 &= (t_h t_{h-1} \cdots t_1 t_k t_1 \cdots t_{h-1})^2 \\ &= (s_h s_{h-1} \cdots s_2 s_k s_1 (s_k s_k) s_k s_1 s_k s_2 \cdots s_{h-1})^2 \\ &= (s_h s_{h-1} \cdots s_2 (s_k s_1 s_k s_1 s_k) s_2 \cdots s_{h-1})^2 \\ &= (s_h s_{h-1} \cdots s_2 s_1 s_2 \cdots s_{h-1})^2 = r(h, h-1)^2 = e. \end{aligned}$$

(k) This is trivial.

(l) Suppose that  $k$  is connected to exactly one vertex of  $C$  via an edge of unspecified weight. We label the vertices as follows.



If the edge between  $k$  and  $h$  points towards  $h$ , then  $t_h = s_h$  and the relations for  $C'$  are the same as the relations for  $C$ . If the edge points towards  $k$ , then  $t_h = s_k s_h s_k$  and we have:

$$\begin{aligned} r'(1, 2)^2 &= (t_1 t_2 \cdots t_h t_{h-1} \cdots t_2)^2 \\ &= (s_1 s_2 \cdots s_{h-1} s_k s_h s_k s_{h-1} \cdots s_2)^2 \\ &= s_k (s_1 s_2 \cdots s_{h-1} s_h s_{h-1} \cdots s_2)^2 s_k \\ &= s_k r(1, 2)^2 s_k = e. \end{aligned}$$

This finishes the proof.  $\square$

Note that, in the simply-laced case, some arguments of this kind have been used in the proof of [5, Thm. 6.10]. We explain more about the connection with [5] at the end of Section 6.

Recall that we are in the situation where  $\Gamma' = \mu_k(\Gamma)$ . We denote by  $s'_i$  the defining generators of  $W_{\Gamma'}$ .

For each vertex  $i$  of  $\Gamma$  define the element  $t'_i$  in  $W_{\Gamma'}$  as follows:

$$t'_i = \begin{cases} s'_k s'_i s'_k & \text{if there is an arrow } k \rightarrow i \text{ in } \Gamma' \text{ (of arbitrary weight);} \\ s'_i & \text{otherwise.} \end{cases}$$

**Proposition 5.3.** *The elements  $t'_i$ , for  $i$  a vertex of  $\Gamma'$ , satisfy the relations (R1)-(R3) defining  $W_{\Gamma}$ .*

*Proof.* This follows from Proposition 5.2, interchanging  $\Gamma$  and  $\Gamma'$  and using the fact that the definition of the group  $W_{\Gamma}$  is unchanged under reversing the orientation of all the arrows in  $\Gamma$ .  $\square$

We can now prove the following:

**Theorem 5.4.** (a) *Let  $\Gamma$  be a diagram of finite type and  $\Gamma' = \mu_k(\Gamma)$  the mutation of  $\Gamma$  at a vertex  $k$ . Then  $W_{\Gamma} \cong W_{\Gamma'}$ .*

(b) *Let  $\mathcal{A}$  be a cluster algebra of finite type. Then the groups  $W_{\Gamma}$  associated to the diagrams  $\Gamma$  arising from the seeds of  $\mathcal{A}$  are all isomorphic (to the reflection group associated to the Dynkin diagram associated to  $\mathcal{A}$ ).*

*Proof.* Denote the generators of  $W_{\Gamma'}$  (as defined in Section 3) by  $s'_i$ , for  $1 \leq i \leq n$ . By Proposition 5.2, there is a group homomorphism  $\varphi : W_{\Gamma'} \rightarrow W_{\Gamma}$  such that  $\varphi(s'_i) = t_i$  for all  $i$ . By Proposition 5.3 there is a group homomorphism  $\psi : W_{\Gamma} \rightarrow W_{\Gamma'}$  such that  $\psi(s_i) = t'_i$ . If there is no arrow  $i \rightarrow k$  in  $\Gamma$ , then there is also no arrow  $k \rightarrow i$  in  $\Gamma'$  and consequently

$$\psi(\varphi(s'_i)) = \psi(t_i) = \psi(s_i) = t'_i = s'_i.$$

If there is an arrow  $i \rightarrow k$  in  $\Gamma$ , then there is an arrow  $k \rightarrow i$  in  $\Gamma'$  and therefore

$$\begin{aligned} \psi(\varphi(s'_i)) &= \psi(t_i) \\ &= \psi(s_k s_i s_k) \\ &= \psi(s_k) \psi(s_i) \psi(s_k) \\ &= t'_k t'_i t'_k \\ &= s'_k s'_i s'_k s'_k = s'_i. \end{aligned}$$

So  $\psi \circ \varphi = \text{id}_{W_{\Gamma'}}$ , and, similarly,  $\varphi \circ \psi = \text{id}_{W_{\Gamma}}$ , showing that  $\varphi$  and  $\psi$  are isomorphisms and (a) is proved. Part (b) follows from part (a) and Remark 3.1.  $\square$

## 6. COMPANION BASES

We first recall some results from [1]. Recall that a matrix  $A$  is said to be *symmetrisable* if there is diagonal matrix  $D$ , with positive entries on the diagonal, such that  $DA$  is symmetric. A symmetrisable matrix  $A$  is said to be *quasi-Cartan* if its diagonal entries are all equal to 2, and it is said to be *positive* if  $DA$  is positive definite for some positive diagonal matrix  $D$  (this does not depend on the choice of  $D$ ). If  $B$  is a skew-symmetrisable matrix, a *quasi-Cartan companion* of  $B$  is a quasi-Cartan matrix  $A$  such that  $|A_{ij}| = |B_{ij}|$  for all  $i, j$ . Such matrices are used in [1] to give a characterisation of cluster algebras of finite type:

**Theorem 6.1.** [1] *Let  $B$  be a skew-symmetrisable matrix. Then the cluster algebra  $\mathcal{A}(B)$  is of finite type if and only if every chordless cycle in  $\Gamma(B)$  is cyclically oriented, and  $B$  has a positive quasi-Cartan companion.*

We also mention the following result, which we shall need later:

**Proposition 6.2.** [1] *Let  $A$  be a positive quasi-Cartan companion of a skew-symmetrisable matrix  $B$ . Then, for every chordless cycle*

$$i_0 \rightarrow i_1 \rightarrow \cdots \rightarrow i_{d-1} \rightarrow i_0$$

in  $\Gamma(B)$ , the following holds:

$$\prod_{a=0}^{d-1} (-1)^{(-A_{i_a, i_{a+1}})} < 0.$$

Let  $\Phi$  be a root system of rank  $n$  associated to a Dynkin diagram  $\Delta$  and let  $B$  be the exchange matrix in a seed  $(\mathbf{x}, B)$  of the cluster algebra of type  $\Delta$ . Motivated by [1], Parsons [10, Defn. 4.1] defines a *companion basis* of  $B$  to be a subset  $\beta_1, \beta_2, \dots, \beta_n$  of  $\Phi$  which is a  $\mathbb{Z}$ -basis of  $\mathbb{Z}\Phi$  such that the matrix  $A$  with  $i, j$  entry  $A_{ij} = (\beta_i, \beta_j)$  for all  $i, j$  is a quasi-Cartan companion of  $B$ . In the general case (not necessarily simply-laced), we use  $A_{ij} = (\beta_i, \beta_j^\vee)$ . Parsons uses companion bases to obtain the dimension vectors of the indecomposable modules over the cluster-tilted algebra associated to  $(\mathbf{x}, B)$  by [3] and [4], in the type  $A$  case (they are obtained by expanding arbitrary roots in terms of the basis; see [10, Theorem 5.3]). We remark that independent later work of Ringel [11] also obtains such dimension vectors (in a more general setting which includes all the finite type cases).

Suppose that  $\mathcal{B} = \{\beta_1, \beta_2, \dots, \beta_n\}$  is a companion basis for an exchange matrix  $B$  in a cluster algebra of finite type as above, and fix  $1 \leq k \leq n$ . Parsons [10, Thm. 6.1] defines the (inward) *mutation*  $\mathcal{B}' = \mu_k(\mathcal{B})$  of  $\mathcal{B}$  at  $k$  to be the subset  $\{\beta'_1, \beta'_2, \dots, \beta'_n\}$  of  $\Phi$  defined by

$$\beta'_i = \begin{cases} s_{\beta_k}(\beta_i) & \text{if there is an arrow from } i \text{ to } k \text{ in } \Gamma(B); \\ \beta_i & \text{otherwise.} \end{cases}$$

Outward mutation of  $\mathcal{B}$  is defined similarly.

**Proposition 6.3.** [10, Thm. 6.1] *Let  $\mathcal{B}$  be a companion basis for an exchange matrix  $B$  in a cluster algebra of finite simply-laced type and fix  $1 \leq k \leq n$ . Then  $\mu_k(\mathcal{B})$  is a companion basis for  $\mu_k(B)$ .*

We remark that mutations of quasi-Cartan companion matrices were considered in [1].

Proposition 6.3 can be extended to all finite type cases using a similar argument:

**Proposition 6.4.** *Let  $\mathcal{B}$  be a companion basis for an exchange matrix  $B$  in a cluster algebra of finite type and fix  $1 \leq k \leq n$ . Then  $\mu_k(\mathcal{B})$  is a companion basis for  $\mu_k(B)$ .*

*Proof.* We must check that if  $\mathcal{B} = \{\beta_1, \beta_2, \dots, \beta_n\}$  is a companion basis for an exchange matrix  $B$  underlying a diagram  $\Gamma$  then  $\mathcal{B}' = \{\gamma_1, \gamma_2, \dots, \gamma_n\} = \mu_k(\mathcal{B})$  is a

companion basis for the mutation  $B' = \mu_k(B)$ , which underlies  $\mu_k(\Gamma)$ . Note that [10, Thm. 6.1] covers the cases  $(\gamma_p, \gamma_q)$  where  $p = k$  or  $q = k$  or where at most one of  $p$  and  $q$  is connected to  $k$  by an edge (in this case the assumption of simply-laced type is not used).

Hence, since [10, Thm. 6.1] covers the simply-laced case, we are reduced to cases (d),(e) and (f) in Corollary 2.3. In each case, we are only left with checking that  $|(\gamma_i, \gamma_j)| = |B'_{ij}|$ .

We denote by  $A$  the  $3 \times 3$  matrix

$$\begin{pmatrix} (\beta_i, \beta_i^\vee) & (\beta_i, \beta_k^\vee) & (\beta_i, \beta_j^\vee) \\ (\beta_k, \beta_i^\vee) & (\beta_k, \beta_k^\vee) & (\beta_k, \beta_j^\vee) \\ (\beta_j, \beta_i^\vee) & (\beta_j, \beta_k^\vee) & (\beta_j, \beta_j^\vee) \end{pmatrix}.$$

We consider first case (d), mutating from left to right. There are two possibilities for  $A$ :

$$\begin{pmatrix} 2 & \pm 1 & 0 \\ \pm 2 & 2 & \pm 1 \\ 0 & \pm 1 & 2 \end{pmatrix}, \begin{pmatrix} 2 & \pm 2 & 0 \\ \pm 1 & 2 & \pm 1 \\ 0 & \pm 1 & 2 \end{pmatrix}.$$

In the first case, the mutation of  $B$  is:

$$\begin{pmatrix} 0 & 1 & 0 \\ -2 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix} \rightarrow \begin{pmatrix} 0 & -1 & 1 \\ 2 & 0 & -1 \\ -2 & 1 & 0 \end{pmatrix}.$$

We have

$$\begin{aligned} (\gamma_i, \gamma_j^\vee) &= (s_{\beta_k}(\beta_i), \beta_j^\vee) \\ &= (\beta_i - (\beta_i, \beta_k^\vee)\beta_k, \beta_j^\vee) \\ &= (\beta_i, \beta_j^\vee) - (\beta_i, \beta_k^\vee)(\beta_k, \beta_j^\vee) = \pm(\beta_i, \beta_k^\vee) = \pm 1, \end{aligned}$$

and

$$\begin{aligned} (\gamma_j, \gamma_i^\vee) &= (\beta_j, (s_{\beta_k}(\beta_i))^\vee) \\ &= (s_{\beta_k}(\beta_j), \beta_i^\vee) \\ &= (\beta_j - (\beta_j, \beta_k^\vee)\beta_k, \beta_i^\vee) \\ &= (\beta_j, \beta_i^\vee) - (\beta_j, \beta_k^\vee)(\beta_k, \beta_i^\vee) = \pm(\beta_k, \beta_i^\vee) = \pm 2, \end{aligned}$$

and this case is done.

In the second case, the mutation is:

$$\begin{pmatrix} 0 & 2 & 0 \\ -1 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix} \rightarrow \begin{pmatrix} 0 & -2 & 2 \\ 1 & 0 & -1 \\ -1 & 1 & 0 \end{pmatrix}$$

Arguing as above, we have  $(\gamma_i, \gamma_j^\vee) = \pm 2$  and  $(\gamma_j, \gamma_i^\vee) = \pm 1$  as required.

Next, we consider case (d), mutating from right to left. Note that the skew-symmetrisability of  $B$  imposes restrictions; see also [7, Lemma 7.6]. The only possibilities for  $B$  are thus

$$\begin{pmatrix} 0 & -1 & 1 \\ 2 & 0 & -1 \\ -2 & 1 & 0 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 0 & -2 & 2 \\ 1 & 0 & -1 \\ -1 & 1 & 0 \end{pmatrix}.$$

In the first case, the mutation is

$$\begin{pmatrix} 0 & -1 & 1 \\ 2 & 0 & -1 \\ -2 & 1 & 0 \end{pmatrix} \rightarrow \begin{pmatrix} 0 & 1 & 0 \\ -2 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}.$$

We have  $(\gamma_i, \gamma_j^\vee) = (\beta_i, \beta_j^\vee) - (\beta_i, \beta_k^\vee)(\beta_k, \beta_j^\vee)$ . Moreover,  $|(\beta_i, \beta_j^\vee)| = |(\beta_i, \beta_k^\vee)| = |(\beta_k, \beta_j^\vee)| = 1$ . Note that the matrix  $A$  must be positive (as the  $\beta_i$  form a basis of  $\mathbb{Z}\Phi$ ). Hence, by Proposition 6.2, an odd number of the signs of  $(\beta_i, \beta_j^\vee)$ ,  $(\beta_i, \beta_k^\vee)$  and  $(\beta_k, \beta_j^\vee)$  must be positive. We obtain the following table of possible values:

$(\beta_i, \beta_j^\vee)$	$(\beta_i, \beta_k^\vee)$	$(\beta_k, \beta_j^\vee)$	$(\gamma_i, \gamma_j^\vee)$
1	1	1	0
1	-1	-1	0
-1	1	-1	0
-1	-1	1	0

In the second case, the mutation is

$$\begin{pmatrix} 0 & -2 & 2 \\ 1 & 0 & -1 \\ -1 & 1 & 0 \end{pmatrix} \rightarrow \begin{pmatrix} 0 & 2 & 0 \\ -1 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}.$$

As before,  $(\gamma_i, \gamma_j^\vee) = (\beta_i, \beta_j^\vee) - (\beta_i, \beta_k^\vee)(\beta_k, \beta_j^\vee)$ . We have  $|(\beta_i, \beta_j^\vee)| = |(\beta_i, \beta_k^\vee)| = 2$  and  $|(\beta_k, \beta_j^\vee)| = 1$ . As in the previous case, we obtain a table of possible values:

$(\beta_i, \beta_j^\vee)$	$(\beta_i, \beta_k^\vee)$	$(\beta_k, \beta_j^\vee)$	$(\gamma_i, \gamma_j^\vee)$
2	2	1	0
2	-2	-1	0
-2	2	-1	0
-2	-2	1	0

and case (d) is complete.

Case (e) is very similar to case (d), so we omit it. We are left with case (f) to consider. By symmetry, we only need to consider the mutation from the left hand side of the figure to the right hand side. Using the skew-symmetrisability of  $B$  (see [7, Lemma 7.6]) we conclude that there are only two possible cases for  $B$ , namely

$$\begin{pmatrix} 0 & 1 & -1 \\ -2 & 0 & 2 \\ 1 & -1 & 0 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 0 & 2 & -1 \\ -1 & 0 & 1 \\ 1 & -2 & 0 \end{pmatrix}.$$

In the first case, the mutation is

$$\begin{pmatrix} 0 & 1 & -1 \\ -2 & 0 & 2 \\ 1 & -1 & 0 \end{pmatrix} \rightarrow \begin{pmatrix} 0 & -1 & 1 \\ 2 & 0 & -2 \\ -1 & 1 & 0 \end{pmatrix}.$$

We have  $(\gamma_i, \gamma_j^\vee) = (s_{\beta_k}(\beta_i), \beta_j^\vee) = (\beta_i, \beta_j^\vee) - (\beta_i, \beta_k^\vee)(\beta_k, \beta_j^\vee)$ . Again, using Proposition 6.2, we obtain the following table of possible values:

$(\beta_i, \beta_j^\vee)$	$(\beta_i, \beta_k^\vee)$	$(\beta_k, \beta_j^\vee)$	$(\gamma_i, \gamma_j^\vee)$
1	1	2	-1
1	-1	-2	-1
-1	1	-2	1
-1	-1	2	1

A similar argument shows that  $(\gamma_j, \gamma_i^\vee) = \pm 1$ . In the second case, the mutation is:

$$\begin{pmatrix} 0 & 2 & -1 \\ -1 & 0 & 1 \\ 1 & -2 & 0 \end{pmatrix} \rightarrow \begin{pmatrix} 0 & -2 & 1 \\ 1 & 0 & -1 \\ -1 & 2 & 0 \end{pmatrix}.$$

As before,  $(\gamma_i, \gamma_j^\vee) = (\beta_i, \beta_j^\vee) - (\beta_i, \beta_k^\vee)(\beta_k, \beta_j^\vee)$ . and we obtain the following table of possibilities:

$(\beta_i, \beta_j^\vee)$	$(\beta_i, \beta_k^\vee)$	$(\beta_k, \beta_j^\vee)$	$(\gamma_i, \gamma_j^\vee)$
1	2	1	-1
1	-2	-1	-1
-1	2	-1	1
-1	-2	1	1

A similar argument shows that  $(\gamma_j, \gamma_i^\vee) = \pm 1$ . The proof is complete.  $\square$

**Remark 6.5.** Every companion basis for  $\mu_k(B)$  is of the form  $\mu_k(\mathcal{B})$  for some companion basis  $\mathcal{B}$  for  $B$ , since it can be mutated back to a companion basis for  $B$  using outward mutation, and outward and inward mutation are easily seen to be inverses of each other.

**Corollary 6.6.** [1] *Let  $B$  be an exchange matrix in a cluster algebra of finite type. Then  $B$  has a companion basis.*

*Proof.* This follows from results in [1] (see also [10, Cor. 3.10]). Alternatively, it can be proved using Proposition 6.4.  $\square$

We can now prove:

**Theorem 6.7.** *Let  $\mathcal{B} = \{\beta_1, \beta_2, \dots, \beta_n\} \subseteq \Phi$  be a companion basis for an exchange matrix  $B$  in a cluster algebra of finite type  $\Delta$  (with associated root system  $\Phi$ ). Then there is an isomorphism between  $W$  and  $W_{\Gamma(B)}$  as defined in Section 3 taking  $s_{\beta_i}$  to  $s_i$ . In particular, the defining relations for  $W_{\Gamma(B)}$  can be regarded as giving a presentation of  $W$  in terms of the generators  $s_{\beta_1}, s_{\beta_2}, \dots, s_{\beta_n}$ .*

*Proof.* Note that any simple system in  $\Phi$  is a companion basis for an exchange matrix  $B$  whose Cartan counterpart is a Cartan matrix of type  $\Delta$ . By Remark 3.1 it follows that the result holds for this case.

Suppose that the result holds for every companion basis associated to an exchange matrix  $B$ . Fix  $1 \leq k \leq n$  and let  $\mathcal{B}' = \{\beta'_1, \beta'_2, \dots, \beta'_n\}$  be a companion basis for  $\mu_k(B)$ . By Remark 6.5,  $\mathcal{B}'$  is of the form  $\mu_k(\mathcal{B})$  for some companion basis  $\mathcal{B} = \{\beta_1, \beta_2, \dots, \beta_n\}$  for  $B$ .

By assumption, the result holds for  $\mathcal{B}$ , i.e. there is a group isomorphism  $\tau_{\mathcal{B}} : W \rightarrow W_{\Gamma(B)}$  taking  $s_{\beta_i}$  to  $s_i$  for  $1 \leq i \leq n$ . If there is no arrow from  $i$  to  $k$  in  $\Gamma(B)$  then  $\tau_{\mathcal{B}}(s_{\beta'_i}) = \tau_{\mathcal{B}}(s_{\beta_i}) = s_i = t_i$ . If there is an arrow from  $i$  to  $k$  in  $\Gamma(B)$ , we have:

$$s_{\beta'_i} = s_{s_{\beta_k}(\beta_i)} = s_{\beta_k} s_{\beta_i} s_{\beta_k}^{-1} = s_{\beta_k} s_{\beta_i} s_{\beta_k},$$

so  $\tau_{\mathcal{B}}(s_{\beta'_i}) = s_k s_i s_k = t_i$ .

From the proof of Theorem 5.4, there is an isomorphism  $\psi : W_{\Gamma(B)} \rightarrow W_{\Gamma(B')}$ . If the defining generators of  $W_{\Gamma'}$  are denoted  $s'_i$ , then we have  $\psi(t_i) = s'_i$ . Hence  $\tau_{\mathcal{B}'} := \psi \circ \tau_{\mathcal{B}} : W \rightarrow W_{\Gamma'}$  is a group isomorphism taking  $s_{\beta'_i}$  to  $s'_i$ . Therefore, the result holds for  $\mathcal{B}'$ .

The result follows in general by using induction on the number of mutations needed to obtain  $B$  from a fixed exchange matrix whose Cartan counterpart is a Cartan matrix of type  $\Delta$ . □

Finally, we explain how the perspective of companion bases allows an alternative proof of Theorem 5.4 in the simply-laced case using results from [5]. Let  $B$  be an exchange matrix of a cluster algebra of finite simply-laced type with corresponding diagram  $\Gamma$ . Let  $\mathcal{B} = \{\beta_1, \beta_2, \dots, \beta_n\}$  be a companion basis for  $B$  and let  $A$  be the corresponding matrix with  $A_{ij} = (\beta_i, \beta_j)$ .

Let  $G$  be the unoriented graph on vertices  $1, 2, \dots, n$  with an edge between  $i$  and  $j$  whenever  $A_{ij} \neq 0$ . We define a map  $f$  from the set of edges of  $G$  to  $\{1, -1\}$  by setting  $f(\{i, j\})$  to be the sign of  $A_{ij}$  whenever  $A_{ij} \neq 0$ . In this way we obtain a signed graph  $\mathcal{G}(\mathcal{B}) = (G, f)$  corresponding to  $\mathcal{B}$ , whose underlying unsigned graph coincides with the underlying unoriented graph of  $\Gamma$ .

Note that  $A$  is positive definite, since  $\mathcal{B}$  is a companion basis. By [1, Prop. 1.4] or a remark in the proof of [5, Thm. 6.10], this implies that every induced cycle in  $(G, f)$  has an odd number of positive signs.

We recall the definition of local switching from [5]. Since we are in the positive definite case, this takes a simpler form; see the comment after Remark 4.6 in [5].

**Definition 6.8.** [5, 4.3] *Let  $(G, f)$  be a signed graph and suppose that the symmetric matrix  $A = A_{ij}$  with  $A_{ij} = f(i, j)$  and  $A_{ii} = 2$  for all  $i, j$  is positive definite. Fix a vertex  $k$  of  $G$  and let  $I$  be a subset of the neighbours of  $k$  in  $G$ . Let  $J$  be the set of neighbours of  $k$  in  $G$  that do not lie in  $I$ . Then the local switching of  $(G, f)$  at  $k$  is given by the following operations:*

- (i) *Delete all edges of  $G$  between  $I$  and  $J$ .*

- (ii) For any  $i \in I$  and  $j \in J$  not originally joined in  $G$ , introduce an edge  $\{i, j\}$  with sign chosen so that the 3-cycle between  $i, j, k$  has an even number (i.e. zero or two) of positive signs.
- (iii) Change the signs of all edges between  $k$  and elements of  $I$ .
- (iv) Leave all other edges and signs unchanged.

**Lemma 6.9.** Fix a vertex  $k$  of  $\Gamma$ . Let  $I$  be the set of vertices  $i$  of  $\Gamma$  for which there is an arrow  $i \rightarrow k$  in  $\Gamma$ , so that  $J$  is the set of vertices  $j$  of  $\Gamma$  for which there is an arrow  $k \rightarrow j$  in  $\Gamma$ . Then  $(G', f') = \mathcal{G}(\mu_k(\mathcal{B}))$  coincides with the result of locally switching  $(G, f)$  at  $k$  with respect to  $I$ .

*Proof.* By Proposition 6.3,  $G'$  is the underlying unoriented graph of  $\mu_k(\Gamma)$ . Let  $(G'', f')$  be the result of locally switching  $(G, f)$  at  $k$  with respect to  $I$ . Then it is easy to see from Definition 6.8 that  $G''$  and  $G'$  coincide as unoriented graphs. Let  $\mu_k(\mathcal{B}) = (\beta'_1, \dots, \beta'_n)$ . If  $i \in I$  then

$$(\beta'_i, \beta'_k) = (s_{\beta_k}(\beta_i), \beta_k) = -(\beta_i, \beta_k)$$

and we see that the signs on the edges of  $G'$  incident with  $k$  and a vertex in  $I$  coincide with the signs on the corresponding edges of  $G''$ . If  $\{i, j\}$  is an edge in  $G''$  arising from step (ii) in Definition 6.8 its sign is chosen to ensure that the 3-cycle on  $i, j, k$  has an odd number of positive signs after step (iii) has been applied. Since  $(G', f')$  must also have this property (using Proposition 6.2 and the fact that  $\mathcal{B}'$  is a companion basis). It is easy to see that the other signs on the edges of  $G'$  and  $G''$  coincide, and the result follows.  $\square$

Since the signed graph  $\mathcal{G}(\mathcal{B})$  has an odd number of positive signs in every cycle we have that the group  $\text{Cox}(G, f)$  defined in [5, Defn. 6.2] coincides with the group  $W_\Gamma$  defined in Section 3, in the simply-laced case. Then Theorem 5.4 (in the simply-laced case) follows by combining [5, Thm. 6.10] and [10, Thm. 6.1] (see Proposition 6.3) with Lemma 6.9.

**Acknowledgements:** This work was partly carried out while both authors were visiting the Institute for Mathematical Research (FIM) at the ETH in Zurich. Both authors would like to thank the FIM for its support and, in particular, Karin Baur for her kind hospitality during the visit. The second named author would like to thank Konstanze Rietsch for some helpful conversations.

#### REFERENCES

- [1] M. Barot, C. Geiss and A. Zelevinsky, *Cluster algebras of finite type and positive symmetrizable matrices*, J. London Math. Soc. (2) **73** (2006), no. 3, 545–564.
- [2] M. Barot and D. Rivera, *Generalized Serre relations for Lie algebras associated with positive unit forms*, J. Pure Appl. Algebra **211** (2007), no. 2, 360–373.
- [3] A. B. Buan, R. J. Marsh, M. Reineke, I. Reiten and G. Todorov, *Tilting theory and cluster combinatorics*, Adv. Math. **204** (2006), no. 2, 572–618.
- [4] P. Caldero, F. Chapoton and R. Schiffler, *Quivers with relations arising from clusters ( $A_n$  case)*, Trans. Amer. Math. Soc. **358** (2006), no. 3, 1347–1364.
- [5] P. J. Cameron, J. J. Seidel and S. V. Tsaranov, *Signed graphs, root lattices, and Coxeter groups*, J. Algebra **164** (1994), no. 1, 173–209.

- [6] S. Fomin and A. Zelevinsky, *Cluster algebras. I. Foundations*, J. Amer. Math. Soc. **15** (2002), no. 2, 497–529.
- [7] S. Fomin and A. Zelevinsky, *Cluster algebras. II. Finite type classification*, Invent. Math. **154** (2003), no. 1, 63–121.
- [8] J. E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge Studies in Advanced Mathematics, **29**. Cambridge University Press, Cambridge, 1990.
- [9] M. J. Parsons, *On indecomposable modules over cluster-tilted algebras of type A*, PhD Thesis, University of Leicester, 2007.
- [10] M. J. Parsons, *Companion bases for cluster-tilted algebras*, Preprint arXiv:1111.0450v1 [math.RT] (2011).
- [11] C. M. Ringel, *Cluster-concealed algebras*, Adv. Math. **226** (2011), no. 2, 1513–1537.

INSTITUTO DE MATEMÁTICAS UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO CIUDAD UNIVERSITARIA, MÉXICO, DISTRITO FEDERAL, C.P. 04510 MÉXICO

*E-mail address:* `barot@matem.unam.mx`

SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS LS2 9JT, ENGLAND

*E-mail address:* `marsh@maths.leeds.ac.uk`