

A CATEGORY OF WIDE SUBCATEGORIES

ASLAK BAKKE BUAN AND ROBERT J. MARSH

ABSTRACT. An algebra is said to be τ -tilting finite provided it has only a finite number of τ -rigid objects up to isomorphism. We associate a category to each such algebra. The objects are the wide subcategories of its category of finite dimensional modules, and the morphisms are indexed by support τ -tilting pairs.

INTRODUCTION AND MAIN RESULT

A full subcategory \mathbf{B} of an abelian category \mathbf{A} is called *wide* if it is an exact abelian subcategory, or equivalently it is closed under kernels, cokernels and extensions.

Let Λ be a finite dimensional algebra over a field k , and $\text{mod } \Lambda$ the category of finitely generated left Λ -modules. Let τ denote the Auslander-Reiten translate in $\text{mod } \Lambda$. Following [1], we call a Λ -module M with $\text{Hom}(M, \tau M) = 0$ a τ -rigid module. The algebra Λ is called τ -tilting finite [5] if there are only a finite number of isomorphism classes of indecomposable τ -rigid Λ -modules. By [1] this is equivalent to Λ having finitely many isomorphism classes of basic τ -tilting modules, as defined in [1]. In particular, all algebras of finite representation type, as well as all preprojective algebras of Dynkin type are τ -tilting finite [10]; see [5] for further examples.

For a module U , let $U^\perp = \{X \in \text{mod } \Lambda \mid \text{Hom}(U, X) = 0\}$, and define ${}^\perp U$ similarly. Jasso [9] proved that, if U is τ -rigid, then the subcategory $J(U) = U^\perp \cap {}^\perp(\tau U)$ is equivalent to a module category, and by [6] we have that $J(U)$ is a wide subcategory of $\text{mod } \Lambda$. For a wide subcategory \mathbf{W} which is equivalent to a module category, and a module V which is τ -rigid in \mathbf{W} , we let $J_{\mathbf{W}}(V) = V^\perp \cap {}^\perp(\tau_{\mathbf{W}} V) \cap \mathbf{W}$. Note that the AR-translations τ in $\text{mod } \Lambda$ and $\tau_{\mathbf{W}}$ in \mathbf{W} will usually be different.

Let $C(\Lambda) = C(\text{mod } \Lambda)$ be the full subcategory of the bounded derived category $D^b(\text{mod } \Lambda)$ with objects corresponding to $\text{mod } \Lambda \amalg (\text{mod } \Lambda)[1]$. For a full subcategory \mathcal{Y} of $\text{mod } \Lambda$, we shall denote by $C(\mathcal{Y})$ the full subcategory $\mathcal{Y} \amalg \mathcal{Y}[1]$ of $C(\Lambda)$. As in [4], we say $\mathcal{U} = U \amalg P[1]$ is *support τ -rigid* in $C(\text{mod } \Lambda)$ if U, P are modules, P is projective, U is τ -rigid and $\text{Hom}(P, U) = 0$. Analogously, if \mathbf{W} is a wide subcategory of $\text{mod } \Lambda$ equivalent to a module category, we will say that an object $\mathcal{U} = U \amalg P[1]$ in $C(\mathbf{W})$, where $U, P \in \mathbf{W}$, the object P is projective in \mathbf{W} , the object U is τ -rigid in \mathbf{W} and $\text{Hom}(P, U) = 0$, is support τ -rigid in $C(\mathbf{W})$. We let $J(\mathcal{U}) = J(U) \cap P^\perp$. We then have the following.

Theorem 0.1. *Let Λ be a finite dimensional algebra, then the following hold.*

This work was supported by FRINAT grant number 231000, from the Norwegian Research Council. The work for this paper was done during several visits of A. B. Buan to Leeds in 2017-2018 and he would like to thank R. J. Marsh and the School of Mathematics at the University of Leeds for their warm hospitality.

- (a) [6, Thm. 3.28], [9, Thm. 3.8] *If \mathcal{U} is support τ -rigid in $C(\text{mod } \Lambda)$, then the subcategory $J(\mathcal{U})$ is wide, and it is equivalent to a module category of a finite dimensional algebra.*
- (b) [6, Thm. 3.34] *If Λ is τ -tilting finite, then any wide subcategory of $\text{mod } \Lambda$ is of the form $J(\mathcal{U})$ for some support τ -rigid object \mathcal{U} in $C(\Lambda)$.*

The aim of the paper is to prove the following result.

Theorem 0.2. *Assume Λ is τ -tilting finite. Then there is a category \mathfrak{B}_Λ whose objects are all wide subcategories of $\text{mod } \Lambda$ and such that the maps from \mathfrak{W}_1 to \mathfrak{W}_2 are indexed by all basic τ -rigid objects T in $C(\mathfrak{W}_1)$ such that $\mathfrak{W}_2 = J_{\mathfrak{W}_1}(T)$.*

Our results are inspired by a recent paper of Igusa and Todorov [7], where they defined a similar category in the setting of hereditary finite dimensional algebras.

In Section 1 we state the main results of the paper and explain how they are used to prove Theorem 0.2.

1. KEY STEPS FOR THE PROOF OF THE MAIN RESULT

For a (skeletally small) Krull-Schmidt category \mathcal{X} , let $\text{ind } \mathcal{X}$ denote the set of isomorphism classes of indecomposable objects in \mathcal{X} and for any basic object X in \mathcal{X} let $\delta(X)$ denote the number of indecomposable direct summands of X . We generally assume all objects are basic and we always assume subcategories are full and closed under isomorphism.

Firstly, we need the following, which is a generalization of [4, Propositions 5.6 and 5.10], and can be seen as a refinement of [9, Theorem 3.15]. This is crucial.

Theorem 1.1 (Theorem 3.6). *Let \mathcal{U} be a support τ -rigid object in $C(\Lambda)$. Then there are bijections*

$$\{\mathcal{X} \in \text{ind}(C(\Lambda)) \mid \mathcal{X} \amalg \mathcal{U} \text{ } \tau\text{-rigid}\} \setminus \text{ind } \mathcal{U}$$

$$\mathcal{E}_{\mathcal{U}} \downarrow \uparrow \mathcal{E}_{\mathcal{U}}$$

$$\{\mathcal{X} \in \text{ind}(C(J(\mathcal{U}))) \mid \mathcal{X} \text{ is support } \tau\text{-rigid in } C(J(\mathcal{U}))\}.$$

The map $\mathcal{E}_{\mathcal{U}}$ can be extended additively, giving the following:

Theorem 1.2 (Theorem 3.7). *Let \mathcal{U} be a support τ -rigid object in $C(\Lambda)$ with $\delta(\mathcal{U}) = t'$. For any positive integer $t \leq n - t'$, the map $\mathcal{E}_{\mathcal{U}}$ induces a bijection between:*

- (a) *The set of support τ -rigid objects \mathcal{X} in $C(\Lambda)$ such that $\delta(\mathcal{X}) = t$, the object $\mathcal{X} \amalg \mathcal{U}$ is support τ -rigid and $\text{add } \mathcal{X} \cap \text{add } \mathcal{U} = 0$, and*
- (b) *The set of support τ -rigid objects \mathcal{X} in $C(J(\mathcal{U}))$ such that $\delta(\mathcal{X}) = t$.*

From now on we assume Λ is τ -tilting finite. Then, using Theorem 0.1, we obtain the following as a direct consequence of Theorems 1.1 and 1.2.

Corollary 1.3. *Assume Λ is τ -tilting finite, and let \mathfrak{W} be a wide subcategory of $\text{mod } \Lambda$. Let \mathcal{U} be a support τ -rigid object in $C(\mathfrak{W})$. Then there is a bijection $\mathcal{E}_{\mathcal{U}}^{\mathfrak{W}}$ from*

$$\{\mathcal{X} \in \text{ind}(C(\mathfrak{W})) \mid \mathcal{X} \amalg \mathcal{U} \text{ } \tau\text{-rigid}\} \setminus \text{ind } \mathcal{U}$$

to

$$\{\mathcal{X} \in \text{ind}(C(J_W(\mathcal{U})) \mid \mathcal{X} \text{ is support } \tau\text{-rigid in } C(J_W(\mathcal{U}))\}.$$

Furthermore, the map $\mathcal{E}_{\mathcal{U}}^W$ induces a bijection between:

- (a) The set of support τ -rigid objects \mathcal{X} in $C(W)$ such that $\mathcal{X} \amalg \mathcal{U}$ is support τ -rigid, with $\delta(\mathcal{X}) = t$ and $\text{add } \mathcal{X} \cap \text{add } \mathcal{U} = 0$, and
- (b) The set of support τ -rigid objects \mathcal{X} in $C(J_W(\mathcal{U}))$ with $\delta(\mathcal{X}) = t$.

Note that $[1]$ always denotes the shift in $D^b(\text{mod } \Lambda)$ rather than the shift in $D^b(W)$ for some wide subcategory W .

The next main ingredient is the following:

Theorem 1.4 (Theorem 4.3). *Assume Λ is τ -tilting finite. Let \mathcal{U} and \mathcal{V} be support τ -rigid objects in $C(\Lambda)$ with no common direct summands, and suppose that $\mathcal{U} \amalg \mathcal{V}$ is support τ -rigid. Then $\mathcal{E}_{\mathcal{U}}(\mathcal{V})$ is support τ -rigid in $C(J(\mathcal{U}))$ and the following equation holds:*

$$J_{J(\mathcal{U})}(\mathcal{E}_{\mathcal{U}}(\mathcal{V})) = J(\mathcal{U} \amalg \mathcal{V}).$$

This has the following direct consequence, using Theorem 0.1.

Corollary 1.5. *Assume Λ is τ -tilting finite and let W be a wide subcategory of $\text{mod } \Lambda$. Let \mathcal{U} and \mathcal{V} be support τ -rigid objects in $C(W)$ with no common direct summands. Then $\mathcal{E}_{\mathcal{U}}^W(\mathcal{V})$ is support τ -rigid in $C(J_W(\mathcal{U}))$ and the following equation holds:*

$$J_{J_W(\mathcal{U})}(\mathcal{E}_{\mathcal{U}}^W(\mathcal{V})) = J_W(\mathcal{U} \amalg \mathcal{V}).$$

For a τ -tilting finite algebra Λ , we can now define \mathfrak{B}_{Λ} as follows. The objects of \mathfrak{B}_{Λ} are the wide subcategories of $\text{mod } \Lambda$. Suppose W_1 and W_2 are two such wide subcategories. If $W_2 \not\subseteq W_1$, then we set $\text{Hom}(W_1, W_2) = \emptyset$. Suppose that $W_2 \subseteq W_1$. Then we set

$$\text{Hom}(W_1, W_2) = \left\{ g_T^{W_1} \mid \begin{array}{l} T \text{ is a basic support } \tau\text{-rigid object in } C(W_1) \\ \text{and } W_2 = J_{W_1}(T) \end{array} \right\},$$

where $g_T^{W_1}$ is a formal symbol associated to W_1 and T . Thus, in general g_T^W is a morphism in \mathfrak{B}_{Λ} from W to $J_W(T)$.

Suppose that W_1 , W_2 and W_3 are wide subcategories of Λ and $W_3 \subseteq W_2 \subseteq W_1$. Let $a \in \text{Hom}(W_1, W_2)$ and $b \in \text{Hom}(W_2, W_3)$. Then there are support τ -rigid objects \mathcal{U} in W_1 and $\overline{\mathcal{V}}$ in W_2 such that $a = g_{\mathcal{U}}^{W_1}$ and $b = g_{\overline{\mathcal{V}}}^{W_2}$, so that $W_2 = J_{W_1}(\mathcal{U})$ and $W_3 = J_{W_2}(\overline{\mathcal{V}})$. By Theorem 1.2, we can write $\overline{\mathcal{V}} = \mathcal{E}_{\mathcal{U}}^{W_1}(\mathcal{V})$ for some support τ -rigid object \mathcal{V} in $C(W_1)$ such that $\mathcal{U} \amalg \mathcal{V}$ is support τ -rigid and $\text{add } \mathcal{V} \cap \text{add } \mathcal{U} = 0$. Thus, we have $b = g_{\mathcal{E}_{\mathcal{U}}^{W_1}(\mathcal{V})}^{W_2}$.

By Theorem 1.5,

$$J_{W_1}(\mathcal{U} \amalg \mathcal{V}) = J_{J_{W_1}(\mathcal{U})}(\mathcal{E}_{\mathcal{U}}^{W_1}(\mathcal{V})) = J_{W_2}(\mathcal{E}_{\mathcal{U}}^{W_1}(\mathcal{V})) = J_{W_2}(\overline{\mathcal{V}}) = W_3,$$

so we may define:

$$b \circ a = g_{\mathcal{E}_{\mathcal{U}}^{W_1}(\mathcal{V})}^{W_2} \circ g_{\mathcal{U}}^{W_1} = g_{\mathcal{U} \amalg \mathcal{V}}^{W_1},$$

since this is a morphism from W_1 to W_3 .

For associativity of composition in \mathfrak{B}_{Λ} we need the following theorem.

Theorem 1.6 (Theorem 5.9). *Assume Λ is τ -tilting finite, and let \mathcal{U} and \mathcal{V} be support τ -rigid objects in $C(\Lambda)$ with no common direct summands. Then*

$$\mathcal{E}_{\mathcal{E}_u(\mathcal{V})}^{J(\mathcal{U})} \mathcal{E}_u = \mathcal{E}_{u \amalg \mathcal{V}}$$

The following is then a direct consequence, using Theorem 0.1.

Corollary 1.7. *Assume Λ is τ -tilting finite, and let \mathcal{W} be a wide subcategory of $\text{mod } \Lambda$. Let \mathcal{U} and \mathcal{V} be support τ -rigid objects in $C(\mathcal{W})$ with no common direct summands, and suppose that $\mathcal{U} \amalg \mathcal{V}$ is support τ -rigid in $C(\mathcal{W})$. Then*

$$\mathcal{E}_{\mathcal{E}_u^{\mathcal{W}}(\mathcal{V})}^{J_{\mathcal{W}}(\mathcal{U})} \mathcal{E}_u^{\mathcal{W}} = \mathcal{E}_{u \amalg \mathcal{V}}^{\mathcal{W}}$$

We are then in position to prove the following.

Corollary 1.8. *The composition operation defined above is associative.*

Proof. For a wide subcategory \mathcal{W} of $\text{mod } \Lambda$ and support τ -rigid object \mathcal{U} in $C(\mathcal{W})$, let $\mathcal{F}_u^{\mathcal{W}}$ denote the inverse of the bijection $\mathcal{E}_u^{\mathcal{W}}$.

Consider now maps

$$\mathcal{W}_1 \xrightarrow{g_u^{W_1}} \mathcal{W}_2 \xrightarrow{g_{\mathcal{V}}^{W_2}} \mathcal{W}_3 \xrightarrow{g_{\mathcal{W}}^{W_3}} \mathcal{W}_4$$

where $\mathcal{W}_2 = J_{\mathcal{W}_1}(\mathcal{U})$, $\mathcal{W}_3 = J_{\mathcal{W}_2}(\mathcal{V})$ and $\mathcal{W}_4 = J_{\mathcal{W}_3}(\mathcal{W})$. Thus \mathcal{U} is a support τ -rigid object in $C(\mathcal{W}_1)$, the object \mathcal{V} is support τ -rigid in $C(\mathcal{W}_2)$ and \mathcal{W} is a support τ -rigid object in $C(\mathcal{W}_3)$, and $\mathcal{W}_4 \subseteq \mathcal{W}_3 \subseteq \mathcal{W}_2 \subseteq \mathcal{W}_1$.

We then have that $g_{\mathcal{V}}^{W_2} \circ g_u^{W_1} = g_{u \amalg \mathcal{F}_u^{W_1}(\mathcal{V})}^{W_1}$ and $g_{\mathcal{W}}^{W_3} \circ g_{\mathcal{V}}^{W_2} = g_{\mathcal{V} \amalg \mathcal{F}_{\mathcal{V}}^{W_2}(\mathcal{W})}^{W_2}$. Hence it follows that

$$g_{\mathcal{W}}^{W_3} \circ (g_{\mathcal{V}}^{W_2} \circ g_u^{W_1}) = g_{u \amalg \mathcal{F}_u^{W_1}(\mathcal{V}) \amalg \mathcal{F}_{u \amalg \mathcal{F}_u^{W_1}(\mathcal{V})}^{W_1}(\mathcal{W})}^{W_1}$$

and that

$$(g_{\mathcal{W}}^{W_3} \circ g_{\mathcal{V}}^{W_2}) \circ g_u^{W_1} = g_{u \amalg \mathcal{F}_u^{W_1}(\mathcal{V} \amalg \mathcal{F}_{\mathcal{V}}^{W_2}(\mathcal{W}))}^{W_1} = g_{u \amalg \mathcal{F}_u^{W_1}(\mathcal{V}) \amalg \mathcal{F}_u^{W_1} \mathcal{F}_{\mathcal{V}}^{W_2}(\mathcal{W})}^{W_1}.$$

It follows from Theorem 1.6 that

$$\mathcal{F}_{u \amalg \mathcal{F}_u^{W_1}(\mathcal{V})}^{W_1} = \mathcal{F}_u^{W_1} \mathcal{F}_{\mathcal{V}}^{W_2}$$

and the claim follows. \square

Finally, we note that for each wide subcategory \mathcal{W} , we can consider the trivial support τ -rigid object 0 in $C(\mathcal{W})$ which gives rise to a map $g_0^{\mathcal{W}} : \mathcal{W} \rightarrow \mathcal{W}$. It is easy to check that this satisfies the axioms required for an identity map. This completes the proof of the main result, Theorem 0.2.

The paper is organized as follows. First, in Section 2, we give some background and notation. In Section 3 we prove Theorem 1.1 and Theorem 1.2, while in Section 4 we deal with Theorem 1.4. Sections 5 - 9 are devoted to the proof of Theorem 1.6. In Section 10, we consider the morphisms in \mathfrak{B}_{Λ} from a wide subcategory to a subcategory of corank 1, and in Section 11 we show how to interpret signed τ -exceptional sequences in terms of factorizations of morphisms in \mathfrak{B}_{Λ} . We conclude with an example in Section 12.

2. BACKGROUND AND NOTATION

Let $\mathcal{P}(\Lambda)$ denote the full subcategory of projective objects in $\text{mod } \Lambda$ and if \mathcal{X} is a subcategory of $\text{mod } \Lambda$, let $\mathcal{P}(\mathcal{X})$ denote the full subcategory of \mathcal{X} consisting of the Ext-projective objects in \mathcal{X} , i.e. the objects P in \mathcal{X} such that $\text{Ext}^1(P, X) = 0$ for all $X \in \mathcal{X}$.

For an object U in an additive category \mathcal{C} , let $\text{add } U$ denote the additive subcategory of \mathcal{C} generated by U , i.e. the full subcategory of all direct summands in direct sums of copies of U . If \mathbf{A} is abelian, we denote by $\text{Gen } U$ the full subcategory of \mathbf{A} consisting of all objects which are factor objects of objects in $\text{add } U$. We assume throughout that Λ is basic and denote $\delta(\Lambda)$ by n . We now recall notation and definitions of from [1].

We consider $\text{mod } \Lambda$ as a full subcategory of $D^b(\text{mod } \Lambda)$ by regarding a module as a stalk complex concentrated in degree 0, and we consider the full subcategory $C(\Lambda) = \text{mod } \Lambda \amalg \text{mod } \Lambda[1]$ of $D^b(\text{mod } \Lambda)$. For a module M , we denote by \mathbb{P}_M its minimal projective presentation, considered as a two-term object in $K^b(\mathcal{P}(\Lambda)) \subseteq D^b(\text{mod } \Lambda)$. Here, a two-term object in \mathbf{K} is a complex of the form

$$\cdots 0 \rightarrow 0 \rightarrow P^{-1} \rightarrow P^0 \rightarrow 0 \rightarrow 0 \rightarrow \cdots$$

The following summarizes some facts which we will use throughout the paper.

Lemma 2.1. *Let U, X be in $\text{mod } \Lambda$.*

- (a) [1, Lemma 3.4] $\text{Hom}(U, \tau X) = 0$ if and only if $\text{Hom}_{\mathcal{D}}(\mathbb{P}_X, \mathbb{P}_U[1]) = 0$. In particular, the module U is τ -rigid if and only if $\text{Hom}_{\mathcal{D}}(\mathbb{P}_U, \mathbb{P}_U[1]) = 0$.
- (b) [3, Theorem 5.10] $\text{Hom}(U, \tau X) = 0$ if and only if $\text{Ext}^1(X, \text{Gen } U) = 0$
- (c) Let \mathbb{X} and \mathbb{Y} be two-term objects in \mathbf{K} . Then H^0 induces an epimorphism $\text{Hom}_{\mathbf{K}}(\mathbb{X}, \mathbb{Y}) \rightarrow \text{Hom}(H^0(\mathbb{X}), H^0(\mathbb{Y}))$ with kernel consisting of the maps factoring through $\text{add } \Lambda[1]$.

We recall that if U is a τ -rigid module in $\text{mod } \Lambda$ then, by [3, Theorem 5.8] there is a torsion pair $(\text{Gen } U, U^\perp)$ in $\text{mod } \Lambda$. We denote the corresponding torsion functors by $t_U : \text{mod } \Lambda \rightarrow \text{Gen } U$ and $f_U : \text{mod } \Lambda \rightarrow U^\perp$. If U is τ -rigid in \mathbf{W} , where \mathbf{W} is a wide subcategory equivalent to a module category, we denote the corresponding torsion functors by $t_U^{\mathbf{W}}$ and $f_U^{\mathbf{W}}$ respectively.

3. BIJECTION

Let \mathcal{U} be an arbitrary (not necessarily indecomposable) support τ -rigid object in $C(\Lambda)$. Then $\mathcal{U} = U \amalg P[1]$, where U is a τ -rigid module, P is in $\mathcal{P}(\Lambda)$ and $\text{Hom}(P, U) = 0$. In this section, we will show that there is a bijection $\mathcal{E}_{\mathcal{U}}$ from the set

$$\{\mathcal{X} \in \text{ind}(C(\Lambda)) \mid \mathcal{X} \amalg \mathcal{U} \text{ } \tau\text{-rigid}\} \setminus \text{ind } \mathcal{U}$$

to

$$\{\mathcal{X} \in \text{ind}(C(J(\mathcal{U}))) \mid \mathcal{X} \text{ } \tau\text{-rigid}\}$$

Such a map has already been defined in [4, Section 4-6] for the case \mathcal{U} is either a τ -rigid module or a shift of a projective module, so we first summarize the construction given there.

Definition 3.1. Let \mathcal{U} be a support τ -rigid object in $C(\Lambda)$ which is either a module or a shift of a projective module. Suppose X lies in the set

$$\{X \in \text{ind}(C(\Lambda)) \mid X \amalg \mathcal{U} \text{ } \tau\text{-rigid}\} \setminus \text{ind } \mathcal{U}.$$

Define $\mathcal{E}_{\mathcal{U}}(X)$ in the following way.

Case I: $\mathcal{U} = U$ is a module.

Case I(a): If X is in $\text{ind}(\text{mod } \Lambda)$, $X \amalg U$ is τ -rigid and $X \notin \text{Gen } U$, then

$$\mathcal{E}_U(X) = f_U(X).$$

Case I(b): If X is in $\text{ind}(\text{mod } \Lambda)$ with $X \amalg U$ τ -rigid and X is in $\text{Gen } U$, then

$$\mathcal{E}_U(X) = f_U(H^0(\mathbb{R}_X)[1])$$

where the triangle

$$\mathbb{R}_X \rightarrow \mathbb{P}_{U_X} \rightarrow \mathbb{P}_X \rightarrow$$

arises from the completion of the minimal right $\text{add } \mathbb{P}_U$ -approximation $\mathbb{P}_{U_X} \rightarrow \mathbb{P}_X$ to a triangle. We have that $\mathbb{R}_X = \mathbb{P}_{B_X}$, for an indecomposable direct summand B_X of the Bongartz complement B of U . Then we have

(i) The triangle

$$(1) \quad \mathbb{P}_{B_X} \rightarrow \mathbb{P}_{U_X} \rightarrow \mathbb{P}_X \rightarrow$$

where the first map is a minimal left $\text{add } \mathbb{P}_U$ -approximation and the second map is a minimal right $\text{add } \mathbb{P}_U$ -approximation;

(ii) The exact sequence obtained from taking the homology of (1):

$$B_X \rightarrow U_X \rightarrow X \rightarrow 0,$$

where the first map is a minimal left $\text{add } U$ -approximation and the second map is a minimal right $\text{add } U$ -approximation,

and we have

$$\mathcal{E}_U(X) = f_U(B_X)[1].$$

The object B_X is shown to be in $\mathcal{P}({}^\perp \tau U)$ and $f_U(B_X)$ is in $\mathcal{P}(J(U))$, so $\mathcal{E}_U(X)$ is in $\text{ind } \mathcal{P}(J(U))[1]$.

Case I(c): If X is in $\text{ind}(\mathcal{P}(\Lambda) \cap {}^\perp U)[1]$, write $X = Q[1]$, with Q in $\text{ind } \mathcal{P}(\Lambda) \cap {}^\perp U$. We have the triangle

$$\mathbb{P}_{B_X} \rightarrow \mathbb{P}_{U_X} \rightarrow Q[1] \rightarrow$$

as in case (b), where the first map is a minimal left $\text{add } \mathbb{P}_U$ -approximation and the second map is a minimal right $\text{add } \mathbb{P}_U$ -approximation. Taking homology gives the exact sequence

$$Q \rightarrow B_X \rightarrow U_X \rightarrow 0$$

where the first map is a minimal left $\mathcal{P}({}^\perp(\tau U))$ -approximation and the second map is a minimal left $\text{add } U$ -approximation.

We set

$$\mathcal{E}_U(X) = f_U(B_X)[1].$$

Case II: $\mathcal{U} = P[1]$, where P is a projective module.

Case II(a): If X is τ -rigid in $\text{ind}(\text{mod } \Lambda)$ and $\text{Hom}(P, X) = 0$, then set $\mathcal{E}_{P[1]}(X) = X$.

Case II(b): If $X = Q[1]$ with Q in $(\text{ind}(\mathcal{P}(\Lambda) \setminus \text{ind } P)[1])$, then set $\mathcal{E}_{P[1]}(X) = \mathcal{E}_{P[1]}(Q[1]) = f_P(Q)[1]$.

Theorem 3.2. [4, Proposition 5.6 and 5.10] *Let \mathcal{U} be a support τ -rigid object in $\mathcal{C}(\Lambda)$. Then we have the following.*

(a) *If $\mathcal{U} = U$ is a module, then \mathcal{E}_U gives a bijection between*

(i)

$$\{X \in \text{ind}(\text{mod } \Lambda) \mid X \amalg U \text{ } \tau\text{-rigid}, X \notin \text{Gen } U\}$$

and

$$\{X \in \text{ind}(J(U)) \mid X \text{ } \tau\text{-rigid in } J(U)\}.$$

(ii)

$$\{X \in \text{ind}(\text{mod } \Lambda) \mid X \amalg U \text{ } \tau\text{-rigid}, X \in \text{Gen } U\} \cup \{(\text{ind } \mathcal{P}(\Lambda) \cap {}^\perp U)[1]\}$$

and

$$\{\text{ind } \mathcal{P}(J(U))[1]\}.$$

(b) *If $\mathcal{U} = P[1]$ is the shift of a projective module, then \mathcal{E}_U gives a bijection between*

$$(\{X \in \text{ind}(\text{mod } \Lambda) \mid X \text{ } \tau\text{-rigid}\} \cap P^\perp) \cup (\text{ind } \mathcal{P}(\Lambda) \setminus \text{ind } P)[1]$$

and

$$\{X \in \text{ind}(J(\mathcal{U})) \mid X \text{ } \tau\text{-rigid}\} \cup \text{ind}(\mathcal{P}(J(\mathcal{U}))[1])$$

(noting that $J(\mathcal{U}) = P^\perp$ in this case).

We now consider the general case, where $\mathcal{U} = U \amalg P[1]$, for modules P, U with P projective, is an arbitrary support τ -rigid object in $\mathcal{C}(\Lambda)$. Note first that

$$\{X \in \text{ind}(\mathcal{C}(\Lambda)) \mid X \amalg \mathcal{U} \text{ } \tau\text{-rigid}\} \setminus \text{ind } \mathcal{U}$$

is the union of the sets

$$(\{X \in \text{ind}(\text{mod } \Lambda) \mid X \amalg U \text{ } \tau\text{-rigid}\} \cap P^\perp) \setminus \text{ind } U$$

and

$$((\text{ind } \mathcal{P}(\Lambda) \cap {}^\perp U) \setminus \text{ind } P)[1],$$

and that

$$\begin{aligned} & \{X \in \text{ind}(\mathcal{C}(J(\mathcal{U})) \mid X \text{ support } \tau\text{-rigid in } \mathcal{C}(J(U))\} \\ & = \{X \in \text{ind}(J(\mathcal{U})) \mid X \text{ } \tau\text{-rigid in } J(U)\} \cup \text{ind}(\mathcal{P}(J(\mathcal{U}))[1]), \end{aligned}$$

so we next analyse the behaviour of \mathcal{E}_U when applied to a module $X \in P^\perp$.

Lemma 3.3. *Let U be a τ -rigid module. Then:*

(a) *The map \mathcal{E}_U restricts to a bijection between*

$$\{X \in \text{ind}(\text{mod } \Lambda) \mid X \amalg U \text{ } \tau\text{-rigid}, X \notin \text{Gen } U\} \cap P^\perp$$

and

$$\{X \in \text{ind}(J(U)) \mid X \text{ } \tau\text{-rigid in } J(U)\} \cap P^\perp$$

(b) *The map \mathcal{E}_U restricts to a bijection between*

$$\{X \in \text{ind}(\text{mod } \Lambda) \mid X \amalg U \text{ } \tau\text{-rigid}, X \in \text{Gen } U\} \cap P^\perp \cup ((\text{ind } \mathcal{P}(\Lambda) \setminus \text{ind } P) \cap {}^\perp U)[1]$$

and

$$\text{ind}(\mathcal{P}(J(U)))[1] \setminus \text{ind } \mathcal{E}_U(\text{add } P[1]).$$

(c) *The map \mathcal{E}_U restricts to a bijection between*

$$(\{X \in \text{ind}(\text{mod } \Lambda) \mid X \amalg U \text{ } \tau\text{-rigid}\} \cap P^\perp) \cup ((\text{ind } \mathcal{P}(\Lambda) \setminus \text{ind } P) \cap {}^\perp U)[1]$$

and

$$(\{X \in \text{ind}(J(U)) \mid X \text{ } \tau\text{-rigid in } J(U)\} \cap P^\perp) \cup (\text{ind } \mathcal{P}(J(U)))[1] \setminus \text{ind } \mathcal{E}_U(P[1])$$

Proof. (a) Let X be in $\text{ind}(\text{mod } \Lambda)$ with $X \amalg U$ a τ -rigid module, and such that X is not in $\text{Gen } U$. Then $\mathcal{E}_U(X) = f_U(X)$. Since $\text{Hom}(P, U) = 0$, we also have $\text{Hom}(P, \text{Gen } U) = 0$ and in particular $\text{Hom}(P, t_U(X)) = 0$. Since P is projective, it then follows that $\text{Hom}(P, X) \simeq \text{Hom}(P, f_U(X)) = \text{Hom}(P, \mathcal{E}_U(X))$. Hence the claim follows, using Theorem 3.2(a).

(b) Since $\text{Hom}(P, U) = 0$, we have $\text{Hom}(P, \text{Gen } U) = 0$, and the claim follows Theorem 3.2(b).

(c) The claim follows directly from combining (a) and (b). \square

If \mathcal{W} is a wide subcategory of $\text{mod } \Lambda$ which is equivalent to a module category, and \mathcal{U} is a support τ -rigid object in $\mathcal{C}(\mathcal{W})$ which is either a module or the shift of a projective object in \mathcal{W} , then we denote by $\mathcal{E}_{\mathcal{U}}^{\mathcal{W}}$ the map corresponding to that defined in Definition 3.1.

Note that $\mathcal{E}_U(P[1]) = P[1]$, so we have the map

$$\mathcal{E}_{P[1]}^{J(U)} = \mathcal{E}_{\mathcal{E}_U(P[1])}^{J(U)}.$$

Lemma 3.4. *Let U be a τ -rigid module. Then the set*

$$(\{X \in \text{ind}(J(U)) \mid X \text{ } \tau\text{-rigid in } J(U)\} \cap P^\perp) \cup (\text{ind } \mathcal{P}(J(U)))[1] \setminus \text{ind } \mathcal{E}_U(P[1])$$

is the domain of $\mathcal{E}_{\mathcal{E}_U(P[1])}^{J(U)}$.

Proof. Let Q in $\mathcal{P}(J(U))$ be such that $\mathcal{E}_U(P[1]) = Q[1]$. Recall (see Definition 3.1, Case I(c)) that $Q = f_U(Y_P)$, where $P \rightarrow Y_P$ is a minimal left ${}^\perp(\tau U)$ -approximation, and there is an exact sequence

$$(2) \quad P \rightarrow Y_P \rightarrow U_P \rightarrow 0$$

with U_P in $\text{add } U$.

We claim that

$$(3) \quad J(U) \cap P^\perp = J(U) \cap Q^\perp.$$

It is clear by the definition of $\mathcal{E}_{\mathcal{E}_U(P[1])}^{J(U)}$ that the assertion of the lemma follows from this claim.

In order to prove the claim, let M be in $J(U) \cap P^\perp$ and apply the right exact functor $\text{Hom}(_, M)$ to the sequence (2), to obtain the exact sequence

$$0 \rightarrow \text{Hom}(U_P, M) \rightarrow \text{Hom}(Y_P, M) \rightarrow \text{Hom}(P, M).$$

We have by assumption that $\text{Hom}(U_P, M) = 0 = \text{Hom}(P, M)$, and hence also $\text{Hom}(Y_P, M) = 0$. It then follows that $\text{Hom}(Q, M) = 0$, since there is an epimorphism $Y_P \rightarrow f_U(Y_P) = Q$. So we have $J(U) \cap P^\perp \subseteq J(U) \cap Q^\perp$.

Conversely, suppose M is in $J(U) \cap Q^\perp$. Consider the canonical sequence

$$0 \rightarrow t_U(Y_P) \rightarrow Y_P \rightarrow f_U(Y_P)(= Q) \rightarrow 0$$

for Y_P , and apply $\text{Hom}(_, M)$ to obtain the exact sequence

$$0 \rightarrow \text{Hom}(Q, M) \rightarrow \text{Hom}(Y_P, M) \rightarrow \text{Hom}(t_U(Y_P), M)$$

We have by assumption $\text{Hom}(Q, M) = 0$, and $\text{Hom}(t_U(Y_P), M) = 0$, since $t_U(Y_P)$ is in $\text{Gen } U$ and M is in U^\perp . Hence, also $\text{Hom}(Y_P, M) = 0$. By Lemma 2.1, we then have

$$(4) \quad \text{Hom}(\mathbb{P}_{Y_P}, \mathbb{P}_M) / (\text{add } \Lambda[1]) = 0.$$

We have the following triangle (from the computation of $\mathcal{E}_U(P[1]) = Q[1]$; see Definition 3.1, Case I(c)).

$$(5) \quad \mathbb{P}_{U_P}[-1] \rightarrow P \rightarrow \mathbb{P}_{Y_P} \rightarrow \mathbb{P}_{U_P}$$

Now let $\alpha: P \rightarrow \mathbb{P}_M$ be arbitrary. Since M is in $J(U)$, we have $\text{Hom}(M, \tau U) = 0$ and so by Lemma 2.1, we have $\text{Hom}(\mathbb{P}_{U_P}, \mathbb{P}_M[1]) = 0$. Applying $\text{Hom}(_, \mathbb{P}_M)$ to the triangle (5), we obtain that $\text{Hom}(\mathbb{P}_{Y_P}, \mathbb{P}_M) \rightarrow \text{Hom}(P, \mathbb{P}_M)$ is surjective, and hence that α factors through a map $\mathbb{P}_{Y_P} \rightarrow \mathbb{P}_M$ and hence through $\Lambda[1]$ by (4). We have $\text{Hom}(P, \Lambda[1]) = 0$ and hence we obtain $\text{Hom}(P, \mathbb{P}_M) = 0$. So we have $J(U) \cap Q^\perp \subseteq J(U) \cap P^\perp$, and this finishes the proof of the claim that $J(U) \cap Q^\perp = J(U) \cap P^\perp$, and hence the proof of the lemma. \square

By Lemmas 3.3(c) and 3.4, the composition $\mathcal{E}_{\mathcal{E}_U(P[1])}^{J(U)} \mathcal{E}_U$ is a well-defined map with domain

$$\{X \in \text{ind}(C(\Lambda)) \mid X \amalg \mathcal{U} \text{ } \tau\text{-rigid}\} \setminus \text{ind } \mathcal{U}.$$

We make the following definition:

Definition 3.5. Let U and P be modules such that $\mathcal{U} = U \amalg P[1]$ is a support τ -rigid object in $C(\Lambda)$. We set $\mathcal{E}_{\mathcal{U}} := \mathcal{E}_{\mathcal{E}_U(P[1])}^{J(U)} \mathcal{E}_U$.

We can now prove the main result of this section.

Theorem 3.6. Let $\mathcal{U} = U \amalg P[1]$ be a support τ -rigid object in $C(\Lambda)$. Then the map $\mathcal{E}_{\mathcal{U}}$ is a bijection between the sets

$$\{X \in \text{ind}(C(\Lambda)) \mid X \amalg \mathcal{U} \text{ } \tau\text{-rigid}\} \setminus \text{ind } \mathcal{U}$$

and

$$\{X \in \text{ind}(C(J(\mathcal{U}))) \mid X \text{ support } \tau\text{-rigid in } C(J(\mathcal{U}))\}.$$

Proof. First note that if $P = 0$ or $U = 0$, this is proved in [4, Proposition 5.6 and 5.10].

Using Lemma 3.3(c) and (3) and the fact that $\mathcal{E}_U(P[1]) = Q[1]$, we have that \mathcal{E}_U restricts to a bijection between

$$(\{X \in \text{ind}(\text{mod } \Lambda) \mid X \amalg U \text{ } \tau\text{-rigid}\} \cap P^\perp) \cup ((\text{ind } \mathcal{P}(\Lambda) \setminus \text{ind } P) \cap {}^\perp U)[1]$$

and

$$(\{X \in \text{ind}(J(U)) \mid X \text{ } \tau\text{-rigid}\} \cap Q^\perp) \cup (\text{ind } \mathcal{P}(J(U)) \setminus \text{ind } Q)[1].$$

The target of this map is the domain of $\mathcal{E}_{Q[1]}^{J(U)} = \mathcal{E}_{\mathcal{E}_U(P[1])}^{J(U)}$. Moreover (see Case II in Definition 3.1), the map $\mathcal{E}_{Q[1]}^{J(U)}$ gives a bijection between

$$\{\{X \in \text{ind}(J(U)) \mid X \text{ } \tau\text{-rigid}\} \cap Q^\perp\} \cup (\text{ind } \mathcal{P}(J(U))) \setminus \text{ind } Q[1]$$

and

$$\{X \in \text{ind}(J(U)) \mid X \text{ } \tau\text{-rigid}\} \cup \text{ind}(\mathcal{P}(J(\mathcal{U}))[1]).$$

This finishes the proof of the claim. \square

Note that we have so far only defined $\mathcal{E}_{\mathcal{U}}(X)$ for an object X in the set

$$\{X \in \text{ind}(C(\Lambda)) \mid X \amalg \mathcal{U} \text{ } \tau\text{-rigid}\} \setminus \text{ind } \mathcal{U}.$$

However, we will also need to consider $\mathcal{E}_{\mathcal{U}}$ as a map from the set of all basic objects X (not necessarily indecomposable) in $C(\Lambda)$ such that $X \amalg \mathcal{U}$ is support τ -rigid and $\text{add } X \cap \text{add } \mathcal{U} = 0$, to the set of all support τ -rigid objects in $C(J(\mathcal{U}))$. So for such $X = X_1 \amalg \cdots \amalg X_t$ where the X_i are indecomposable, we define $\mathcal{E}_{\mathcal{U}}(X) = \mathcal{E}_{\mathcal{U}}(X_1) \amalg \cdots \amalg \mathcal{E}_{\mathcal{U}}(X_t)$.

Theorem 3.7. *Let \mathcal{U} be a support τ -rigid object in $C(\Lambda)$ with $\delta(\mathcal{U}) = t'$. For any positive integer $t \leq n - t'$, the map $\mathcal{E}_{\mathcal{U}}$ induces a bijection between the set of basic support τ -rigid objects X in $C(\Lambda)$ such that $\delta(X) = t$, with $X \amalg \mathcal{U}$ support τ -rigid and $\text{add } X \cap \text{add } \mathcal{U} = 0$, and the set of basic support τ -rigid objects \mathcal{Y} in $C(J(\mathcal{U}))$ with $\delta(\mathcal{Y}) = t$.*

Proof. Recall that by definition $\mathcal{E}_{\mathcal{U}} = \mathcal{E}_{\mathcal{E}_U(P[1])}^{J(U)} \mathcal{E}_U$, so the result follows from [4, Prop. 6.7, Prop. 6.10]. \square

Lemma 3.8. *Let U be a τ -rigid module, and P a projective module with $\text{Hom}(P, U) = 0$. Then*

$${}^\perp(\tau_{P^\perp} U) \cap P^\perp = \cap^\perp(\tau U) \cap P^\perp.$$

Proof. We have

$$(6) \quad {}^\perp(\tau_{P^\perp} U) \cap P^\perp = \{Y \in \text{mod } \Lambda \mid \text{Ext}^1(U, \text{Gen}_{P^\perp} Y) = 0\} \cap P^\perp$$

$$(7) \quad \begin{aligned} &= \{Y \in \text{mod } \Lambda \mid \text{Ext}^1(U, \text{Gen } Y) = 0\} \cap P^\perp \\ &= {}^\perp(\tau U) \cap P^\perp, \end{aligned}$$

where (6) holds by Lemma 2.1, and (7) holds since $\text{Gen}_{P^\perp} Y = \text{Gen } Y$ for Y in P^\perp . \square

4. COMPOSITION

The aim of this section is to prove Theorem 1.4.

If \mathbf{A} is (a category equivalent to) a module category, we let $r(\mathbf{A})$ denote the rank of the Grothendieck group of \mathbf{A} , that is: the number of simple objects in \mathbf{A} up to isomorphism. Recall that $\delta(X)$ denotes the number of indecomposable summands in a basic object X . We always write $r(\text{mod } \Lambda) = n$. Recall the following important facts.

Proposition 4.1. *Let \mathcal{U} be a τ -rigid object in $\text{mod } \Lambda$. Then the following hold.*

- (a) [6, Theorem 3.28] $J(\mathcal{U})$ is a wide subcategory of $\text{mod } \Lambda$.
- (b) [9, Theorem 3.8] $J(\mathcal{U})$ is equivalent to a module category with rank $r(J(\mathcal{U})) = n - \delta(\mathcal{U})$.

The following results are crucial.

Proposition 4.2. *Assume that Λ is τ -tilting finite.*

- (a) *For each wide subcategory \mathcal{W} of $\text{mod } \Lambda$, there is a support τ -rigid object \mathcal{U} in $C(\Lambda)$ such that $\mathcal{W} = J(\mathcal{U})$.*
- (b) *If Λ is τ -tilting finite, then each wide subcategory \mathcal{W} of $\text{mod } \Lambda$ is τ -tilting finite.*

Proof. (a) This is contained in Theorem 3.34 in [6].

(b) This is a direct consequence of Theorem 3.6, using (a). \square

We will from now on assume Λ is a τ -tilting finite algebra.

In this section we prove the following (Theorem 1.4).

Theorem 4.3. *Let \mathcal{U} and \mathcal{V} be objects in $C(\Lambda)$ with no common direct summands, and suppose that $\mathcal{U} \amalg \mathcal{V}$ is support τ -rigid. Then $\mathcal{E}_{\mathcal{U}}(\mathcal{V})$ is support τ -rigid in $C(J(\mathcal{U}))$ and the following equation holds*

$$J_{J(\mathcal{U})}(\mathcal{E}_{\mathcal{U}}(\mathcal{V})) = J(\mathcal{U} \amalg \mathcal{V}).$$

This theorem is the key for proving that composition is well-defined in the category \mathfrak{B}_{Λ} . Note first that by Theorem 3.7, we have that $\mathcal{E}_{\mathcal{U}}(\mathcal{V})$ is support τ -rigid in $C(J(\mathcal{U}))$. The remainder of this section is devoted to proving the second assertion of the Theorem.

We first make the following observation.

Lemma 4.4. *In the setting of Theorem 4.3 we have*

$$r(J_{J(\mathcal{U})}(\mathcal{E}_{\mathcal{U}}(\mathcal{V}))) = r(J(\mathcal{U} \amalg \mathcal{V})).$$

Proof. Let $r(\text{mod } \Lambda) = n$. By [9] we have $r(J(\mathcal{T})) = n - \delta(\mathcal{T})$ for any support τ -rigid object \mathcal{T} in $C(\Lambda)$. So $r(J(\mathcal{U})) = n - \delta(\mathcal{U})$ and $r(J(\mathcal{U} \amalg \mathcal{V})) = n - \delta(\mathcal{U}) - \delta(\mathcal{V})$. Furthermore $r(J_{J(\mathcal{U})}(\mathcal{E}_{\mathcal{U}}(\mathcal{V}))) = (n - \delta(\mathcal{U})) - (\delta(\mathcal{E}_{\mathcal{U}}(\mathcal{V}))) = n - \delta(\mathcal{U}) - \delta(\mathcal{V})$, and the claim follows. \square

Lemma 4.5. *Let \mathcal{A} be an abelian category and $\mathcal{A}'' \subseteq \mathcal{A}'$ wide subcategories of \mathcal{A} . Then \mathcal{A}'' is a wide subcategory of \mathcal{A}' .*

Proof. This follows directly from the fact that a subcategory is wide if and only if it is closed under kernels, cokernels and extensions. \square

Proof of Theorem 4.3. We first claim it is sufficient to prove

$$J(\mathcal{U} \amalg \mathcal{V}) \subseteq J_{J(\mathcal{U})}(\mathcal{E}_{\mathcal{U}}(\mathcal{V})).$$

If this holds then, by Lemma 4.5, we have that $J(\mathcal{U} \amalg \mathcal{V})$ is a wide subcategory of $J_{J(\mathcal{U})}(\mathcal{E}_{\mathcal{U}}(\mathcal{V}))$. Then, by Proposition 4.2, there is a support τ -rigid object \mathcal{V}' in $C(J_{J(\mathcal{U})}(\mathcal{E}_{\mathcal{U}}(\mathcal{V})))$ such that

$$J(\mathcal{U} \amalg \mathcal{V}) = J_{J(\mathcal{U})}(\mathcal{E}_{\mathcal{U}}(\mathcal{V}))(\mathcal{V}')$$

We have $r(J_{J(\mathcal{U})}(\mathcal{E}_{\mathcal{U}}(\mathcal{V}))(\mathcal{V}')) = n - \delta(\mathcal{U}) - \delta(\mathcal{V}) - \delta(\mathcal{V}')$ by Proposition 4.1(b) and Theorem 3.7. Hence $r(\mathcal{V}') = 0$, so $\mathcal{V}' = 0$, and we have

$$J(\mathcal{U} \amalg \mathcal{V}) = J_{J(\mathcal{U})}(\mathcal{E}_{\mathcal{U}}(\mathcal{V})).$$

In order to prove

$$(8) \quad J(\mathcal{U} \amalg \mathcal{V}) \subseteq J_{J(\mathcal{U})}(\mathcal{E}_{\mathcal{U}}(\mathcal{V}))$$

we first discuss various special cases.

Case I: Let U be τ -rigid in $\text{mod } \Lambda$, and $V \notin \text{Gen } U$, such that $\bar{V} = \mathcal{E}_U(V)$ is τ -rigid in $J(U)$. Then $\bar{V} = f_U(V)$, and there is an epimorphism $V \rightarrow \bar{V}$.

Let M be in $J(U \amalg V)$. Then we have $M \in J(V) \subseteq V^\perp$, and since $0 \rightarrow \text{Hom}(\bar{V}, M) \rightarrow \text{Hom}(V, M)$ is exact, we also have $\text{Hom}(\bar{V}, M) = 0$.

We next need to show $\text{Hom}(M, \tau_{J(U)}\bar{V}) = 0$. By Lemma 2.1, this is equivalent to showing $\text{Ext}^1(\bar{V}, \text{Gen}_{J(U)} M) = 0$. We have $\text{Gen}_{J(U)} M = \text{Gen } M \cap J(U)$, and hence it is sufficient to prove $\text{Ext}^1(\bar{V}, \text{Gen } M \cap J(U)) = 0$. Let M' be in $\text{Gen } M \cap J(U)$. Apply $\text{Hom}(_, M')$ to the canonical sequence $0 \rightarrow t_U(V) \rightarrow V \rightarrow f_U(V) = \bar{V} \rightarrow 0$ for V , to obtain the exact sequence

$$(9) \quad \text{Hom}(t_U(V), M') \rightarrow \text{Ext}^1(\bar{V}, M') \rightarrow \text{Ext}^1(V, M').$$

The first term in (9) vanishes, since $t_U(V)$ is in $\text{Gen } U$ and M' is in U^\perp . We have $\text{Hom}(M, \tau V) = 0$, since M is in $J(V)$, so $\text{Hom}(M', \tau V) = 0$, since M' is in $\text{Gen } M$. Using the AR-formula, we obtain that the third term in (9) also vanishes, and hence also the second term vanishes. Hence we have that $\text{Ext}^1(\bar{V}, \text{Gen } M \cap J(U)) = 0$ and so $\text{Hom}(M, \tau_{J(U)}\bar{V}) = 0$. So M is in $J_{J(U)}(\bar{V})$, and we have shown inclusion (8) in this case.

Case II (a): Let U be τ -rigid in $\text{mod } \Lambda$, and V in $\text{Gen } U$ such that $\mathcal{E}_U(V) = \bar{V}$ is in $\mathcal{P}(J(U))[1]$. Recall that \bar{V} is computed as follows. We have a triangle

$$\mathbb{P}_{B_V} \xrightarrow{a} \mathbb{P}_{U_V} \xrightarrow{b} \mathbb{P}_V \xrightarrow{c} \mathbb{P}_{B_V}[1]$$

where a is a minimal left add \mathbb{P}_U -approximation and b is a minimal right add \mathbb{P}_U -approximation, and taking homology gives the exact sequence

$$B_V \xrightarrow{a'} U_V \xrightarrow{b'} V \longrightarrow 0$$

where a' is a minimal left add U -approximation and b' is a minimal right add U -approximation. Let $\bar{Q} = f_U(B_V)$. Then $\bar{V} = \bar{Q}[1]$.

Now suppose that M lies in $J(U \amalg V)$. Note that $J_{J(U)}(\mathcal{E}_U(V)) = J(U) \cap \bar{Q}^\perp$. Since M is in $J(U)$, it is sufficient to show that $\text{Hom}(\bar{Q}, M) = 0$. Since \bar{Q} is a quotient of B_V , it is sufficient to show that $\text{Hom}(B_V, M) = 0$. For this let $g : \mathbb{P}_{B_V} \rightarrow \mathbb{P}_M$ be an arbitrary map. By Lemma 2.1, we have that $\text{Hom}(\mathbb{P}_V, \mathbb{P}_M[1]) = 0$, since $\text{Hom}(M, \tau V) = 0$. Hence, the composition $g \circ c[-1] : \mathbb{P}_V[-1] \rightarrow \mathbb{P}_M$ vanishes, and there is a factorization $g = ha$ for some $h : \mathbb{P}_{U_V} \rightarrow \mathbb{P}_M$:

$$\begin{array}{ccccccc} \mathbb{P}_V[-1] & \xrightarrow{c[-1]} & \mathbb{P}_{B_V} & \xrightarrow{a} & \mathbb{P}_{U_V} & \xrightarrow{b} & \mathbb{P}_V \xrightarrow{c} \mathbb{P}_{B_V}[1] \\ & & \downarrow g & \swarrow h & & & \\ & & \mathbb{P}_M & & & & \end{array}$$

Since $\text{Hom}(U, M) = 0$, we have $\text{Hom}(\mathbb{P}_{U_V}, \mathbb{P}_M)/\Lambda[1] = 0$, and it follows that $\text{Hom}(\mathbb{P}_{B_V}, \mathbb{P}_M)/\Lambda[1] = 0$, and hence by Lemma 2.1, we have $\text{Hom}(B_V, M) = 0$. Hence we have shown inclusion (8) in this case.

Case II (b): Let U be τ -rigid in $\text{mod } \Lambda$, and $V \in (\mathcal{P}(\Lambda) \cap {}^\perp U)[1]$. Assume $V = Q[1]$ for Q in $\mathcal{P}(\Lambda) \cap {}^\perp U$.

Recall that $\mathcal{E}_U(V) = \bar{V}$ is computed as follows. There is an exact sequence

$$Q \rightarrow B_V \rightarrow U_V \rightarrow 0$$

where the first map is a minimal left ${}^\perp(\tau U)$ -approximation (or, equivalently, a minimal left $\mathcal{P}({}^\perp(\tau U))$ -approximation), and $\bar{V} = f_U(B_V)[1]$; we set $\bar{Q} = f_U(B_V)$.

Now let M be in $J(U \amalg V)$, that is M is in $J(U)$ and $\text{Hom}(Q, M) = 0$. We need to prove that $\text{Hom}(\bar{Q}, M) = 0$. Since $\bar{Q} = f_U(B_V)$ is a quotient of B_V , it is sufficient to show that $\text{Hom}(B_V, M) = 0$. Recall that there is a triangle

$$Q \rightarrow \mathbb{P}_{B_V} \rightarrow \mathbb{P}_{U_V} \rightarrow$$

and consider an arbitrary map $\mathbb{P}_{B_V} \rightarrow \mathbb{P}_M$. The composition $Q \rightarrow \mathbb{P}_{B_V} \rightarrow \mathbb{P}_M$ vanishes, since $\text{Hom}(Q, M) = 0$ and hence $\text{Hom}(Q, \mathbb{P}_M) = 0$, by Lemma 2.1. Therefore, the map $\mathbb{P}_{B_V} \rightarrow \mathbb{P}_M$ factors $\mathbb{P}_{B_V} \rightarrow \mathbb{P}_{U_V} \rightarrow \mathbb{P}_M$. Since M is in $J(U) \subseteq U^\perp$, we have $\text{Hom}(U_V, M) = 0$, so $\text{Hom}(\mathbb{P}_{U_V}, \mathbb{P}_M) / \text{add } \Lambda[1] = 0$. Hence also $\text{Hom}(\mathbb{P}_{B_V}, \mathbb{P}_M) / \text{add } \Lambda[1] = 0$ and $\text{Hom}(B_V, M) = 0$ as required. Hence we have shown that the inclusion (8) holds also in this case.

Case III: Let $U = P[1]$ with P in $\mathcal{P}(\Lambda)$, and let V be τ -rigid. Then $J(U) = P^\perp$ and $\mathcal{E}_U(V) = \bar{V} = V$ is also $\tau_{J(U)}$ -rigid, by [1, Lemma 2.1]. Furthermore, by Lemma 3.8 we have

$$J_{P^\perp}(V) = P^\perp \cap V^\perp \cap {}^\perp(\tau_{P^\perp} V) = P^\perp \cap V^\perp \cap {}^\perp(\tau V) = J(U \amalg V),$$

which finishes the proof of case III.

Case IV: Now let $U = P[1]$ and $V = Q[1]$, for $P, Q \in \mathcal{P}(\Lambda)$. Then $\mathcal{E}_U(V) = \bar{V} = (f_P Q)[1]$. For an object M in P^\perp , apply $\text{Hom}(\cdot, M)$ to the exact sequence

$$0 \rightarrow t_P(Q) \rightarrow Q \rightarrow f_P(Q) \rightarrow 0$$

to obtain the exact sequence

$$0 \rightarrow \text{Hom}(f_P(Q), M) \rightarrow \text{Hom}(Q, M) \rightarrow \text{Hom}(t_P(Q), M)$$

The last term vanishes, since $t_P(Q)$ is in $\text{Gen } P$, so $\text{Hom}(f_P(Q), M) \simeq \text{Hom}(Q, M)$. Hence, we have $J_{J(U)}(\mathcal{E}_U(V)) = J_{P^\perp}(\bar{V}) = P^\perp \cap (f_P Q)^\perp = P^\perp \cap Q^\perp = J(U \amalg V)$, which finishes the proof of case IV.

General case. Let $\mathcal{U} = U \amalg P[1]$ and $\mathcal{V} = V \amalg Q[1]$, for U, V τ -rigid modules and P, Q in $\mathcal{P}(\Lambda)$. We assume that $\mathcal{U} \amalg \mathcal{V}$ is support τ -rigid in $\mathcal{C}(\Lambda)$. We proceed by induction on the rank $n = r(\text{mod } \Lambda)$. We therefore first assume $U \neq 0$, so $r(J(U)) < n$.

Then

$$(10) \quad J(\mathcal{U} \amalg \mathcal{V}) = J(U \amalg V) \cap P^\perp \cap Q^\perp$$

and

$$(11) \quad \begin{aligned} J_{J(\mathcal{U})}(\mathcal{E}_{\mathcal{U}}(\mathcal{V})) &= J_{J(U) \cap P^\perp}(\mathcal{E}_{U \amalg P[1]}(V \amalg Q[1])) \\ &= J_{J(U) \cap P^\perp}(\mathcal{E}_{\mathcal{E}_U(P[1])}^{J(U)} \mathcal{E}_U(V \amalg Q[1])) \end{aligned}$$

$$(12) \quad = J_{J(U) \cap P^\perp}(\mathcal{E}_{\mathcal{E}_U(P[1])}^{J(U)} \mathcal{E}_U(V)) \cap J_{J(U) \cap P^\perp}(\mathcal{E}_{\mathcal{E}_U(P[1])}^{J(U)} \mathcal{E}_U(Q[1]))$$

Where (11) is by definition by of $\mathcal{E}_{\mathcal{U}} = \mathcal{E}_{U \amalg P[1]}$.

Note that we have

$$(13) \quad J(U) \cap P^\perp = J(U \amalg P[1]) = J_{J(U)}(\mathcal{E}_U(P[1]))$$

by case II(b).

We next compute the terms of (12) separately. For the first term, we obtain

$$(14) \quad J_{J(U) \cap P^\perp}(\mathcal{E}_{\mathcal{E}_U(P[1])}^{J(U)} \mathcal{E}_U(V)) = J_{J_{J(U)}(\mathcal{E}_U(P[1]))}(\mathcal{E}_{\mathcal{E}_U(P[1])}^{J(U)} \mathcal{E}_U(V))$$

$$(15) \quad \begin{aligned} &= J_{J(U)}(\mathcal{E}_U(P[1]) \amalg \mathcal{E}_U(V)) \\ &= J_{J(U)}(\mathcal{E}_U(P[1])) \cap J_{J(U)}(\mathcal{E}_U(V)) \end{aligned}$$

$$(16) \quad \begin{aligned} &= J(U \amalg P[1]) \cap J_{J(U)}(\mathcal{E}_U(V)) \\ &= J(U) \cap P^\perp \cap J_{J(U)}(\mathcal{E}_U(V)) \end{aligned}$$

$$(17) \quad = P^\perp \cap J_{J(U)}(\mathcal{E}_U(V))$$

where (14) follows from (13), and (15) is obtained by using the induction assumption for the proper subcategory $J(U)$, while (16) holds by case II(b) and (17) holds by $J_{J(U)}(\mathcal{E}_U(V)) \subseteq J(U)$.

Similarly, for the second term in (12), we obtain

$$(18) \quad \begin{aligned} J_{J(U) \cap P^\perp}(\mathcal{E}_{\mathcal{E}_U(P[1])}^{J(U)} \mathcal{E}_U(Q[1])) &= J_{J_{J(U)}(\mathcal{E}_U(P[1]))}(\mathcal{E}_{\mathcal{E}_U(P[1])}^{J(U)} \mathcal{E}_U(Q[1])) \\ &= J_{J(U)}(\mathcal{E}_U(P[1]) \amalg \mathcal{E}_U(Q[1])) \\ &= J_{J(U)}(\mathcal{E}_U(P[1] \amalg Q[1])) \\ &= J(U \amalg P[1] \amalg Q[1]) \\ &= J(U) \cap P^\perp \cap Q^\perp \end{aligned}$$

We then obtain

$$(19) \quad J_{J(\mathcal{U})}(\mathcal{E}_{\mathcal{U}}(\mathcal{V})) = J_{J(U) \cap P^\perp}(\mathcal{E}_{\mathcal{E}_U(P[1])}^{J(U)} \mathcal{E}_U(V)) \cap J_{J(U) \cap P^\perp}(\mathcal{E}_{\mathcal{E}_U(P[1])}^{J(U)} \mathcal{E}_U(Q[1]))$$

$$(20) \quad = J_{J(U)}(\mathcal{E}_U(V)) \cap P^\perp \cap J(U) \cap P^\perp \cap Q^\perp$$

$$(21) \quad = J(U \amalg V) \cap P^\perp \cap Q^\perp$$

$$(22) \quad = J(\mathcal{U} \amalg \mathcal{V})$$

where (19) is (12) and where (20) follows from combining (17) and (18). Furthermore (21) follows from Cases I and II(a) and (22) follows from (10) respectively.

So we have that the claim of the theorem holds in the general case, with the assumption that $U \neq 0$.

Now, consider the case where $U = 0$.

We then have

$$(23) \quad \begin{aligned} J(\mathcal{U} \amalg \mathcal{V}) &= J(P[1] \amalg V \amalg Q[1]) \\ &= J(V) \cap P^\perp \cap Q^\perp \end{aligned}$$

and

$$J_{J(\mathcal{U})}(\mathcal{E}_{\mathcal{U}}(\mathcal{V})) = J_{J(P[1])}(\mathcal{E}_{P[1]}(V \amalg Q[1]))$$

$$\begin{aligned}
(24) \quad &= J_{J(P[1])}(\mathcal{E}_{P[1]}(V)) \cap J_{J(P[1])}(\mathcal{E}_{P[1]}(Q[1])) \\
&= J(V \amalg P[1]) \cap J(P[1] \amalg Q[1]) \\
&= J(V) \cap P^\perp \cap Q^\perp \\
&= J(\mathcal{U} \amalg \mathcal{V})
\end{aligned}$$

where for (24), we use cases III and IV. This finishes the proof for the case $U = 0$, and hence the proof of the theorem. \square

5. ASSOCIATIVITY

The aim of this section is to prove that the composition operation defined in Section 4 is associative. The main step is to prove Theorem 1.6. We prepare for this, by giving several useful lemmas.

Lemma 5.1. *Let U, X, Y be in $\text{mod } \Lambda$ where U is τ -rigid and $\text{Hom}(U, \tau X) = 0$. Then the induced map $\alpha: \text{Hom}(X, Y) \rightarrow \text{Hom}(f_U(X), f_U(Y))$ is an epimorphism.*

Proof. Consider the canonical sequences for X and Y ,

$$0 \rightarrow t_U(X) \rightarrow X \rightarrow f_U(X) \rightarrow 0$$

and

$$0 \rightarrow t_U(Y) \rightarrow Y \rightarrow f_U(Y) \rightarrow 0$$

Applying $\text{Hom}(_, f_U(Y))$ to the canonical sequence for X gives the exact sequence

$$0 \rightarrow \text{Hom}(f_U(X), f_U(Y)) \xrightarrow{a} \text{Hom}(X, f_U(Y)) \rightarrow \text{Hom}(t_U(X), f_U(Y))$$

Noting that the last term vanishes, this gives that a is an isomorphism.

Applying $\text{Hom}(X, _)$ to the canonical sequence for Y gives the exact sequence

$$\text{Hom}(X, Y) \xrightarrow{b} \text{Hom}(X, f_U(Y)) \rightarrow \text{Ext}^1(X, t_U(Y)).$$

Since $\text{Hom}(U, \tau X) = 0$ we have by Lemma 2.1 that $\text{Ext}^1(X, \text{Gen } U) = 0$, so in particular $\text{Ext}^1(X, t_U(Y)) = 0$. Hence the map b is an epimorphism. The induced map $\alpha = a^{-1} \circ b$ is then also an epimorphism. \square

We have the following similar lemma:

Lemma 5.2. *Let U, X, Y be in $\text{mod } \Lambda$ where U is τ -rigid and $\text{Hom}(U, Y) = 0$. Then the induced map $\text{Hom}(X, Y) \rightarrow \text{Hom}(f_U(X), f_U(Y))$ is an isomorphism.*

Proof. Since $\text{Hom}(U, Y) = 0$, we have $t_U(Y) = 0$, so $f_U(Y) \simeq Y$. We have the canonical sequence for X :

$$0 \longrightarrow t_U(X) \longrightarrow X \longrightarrow f_U(X) \longrightarrow 0$$

Applying $\text{Hom}(_, Y)$ to this we obtain the exact sequence

$$0 \longrightarrow \text{Hom}(f_U(X), Y) \longrightarrow \text{Hom}(X, Y) \longrightarrow \text{Hom}(t_U(X), Y).$$

The last term vanishes since $\text{Hom}(U, Y) = 0$ implies that $\text{Hom}(\text{Gen } U, Y) = 0$. So we have $\text{Hom}(f_U(X), f_U(Y)) \simeq \text{Hom}(f_U(X), Y) \simeq \text{Hom}(X, Y)$. \square

Lemma 5.1 has the following consequence in terms of approximations:

Lemma 5.3. *Let \mathcal{T} be a subcategory of $\text{mod } \Lambda$. Let U be τ -rigid and assume $\text{Hom}(U, \tau B) = 0$. If $a: B \rightarrow A$ is a left \mathcal{T} -approximation, then $f_U(a): f_U(B) \rightarrow f_U(A)$ is a left $f_U(\mathcal{T})$ -approximation.*

Proof. Let $f_U(T)$ be in $f_U(\mathcal{T})$, and consider a map $b': f_U(B) \rightarrow f_U(T)$. By Lemma 5.1, there is $b: B \rightarrow T$ such that $f_U(b) = b'$. Since $a: B \rightarrow A$ is a left \mathcal{T} -approximation, there is $c: A \rightarrow T$ such that $b = ca$. It follows that $f_U(b) = f_U(c)f_U(a)$, which proves the claim. \square

Lemma 5.3 is used in the proof of part (b) of the following lemma.

Lemma 5.4. *Let U, V be in $\text{mod } \Lambda$, where $U \amalg V$ is τ -rigid. Assume no indecomposable summand in V lies in $\text{Gen } U$ and let $\bar{V} = f_U(V)$. Let $\mathcal{T} = {}^\perp(\tau U \amalg \tau V)$ and let $\mathcal{T}' = {}^\perp(\tau_{J(U)}\bar{V}) \cap J(U)$. Then the following hold.*

- (a) *We have $f_U(\mathcal{T}) = \mathcal{T}'$.*
- (b) *If $B \rightarrow A$ is a left \mathcal{T} -approximation in $\text{mod } \Lambda$ and $\text{Hom}(U, \tau B) = 0$, then $f_U(B) \rightarrow f_U(A)$ is a left \mathcal{T}' -approximation (in $\text{mod } \Lambda$).*

Proof. (a) We first show $f_U(\mathcal{T}) \subseteq \mathcal{T}'$. Since U is in \mathcal{T} , we have $\text{Gen } U \subseteq \mathcal{T}$, and clearly $\mathcal{T} \subseteq {}^\perp(\tau U)$. By [9, Theorem 3.14], we have that $f_U(\mathcal{T}) = \mathcal{T} \cap U^\perp$ is a torsion class in $J(U)$. So

$$f_U(\mathcal{T}) = \mathcal{T} \cap U^\perp = {}^\perp(\tau U \amalg \tau V) \cap U^\perp = {}^\perp(\tau V) \cap J(U),$$

and we want to show that $f_U(\mathcal{T}) = {}^\perp(\tau V) \cap J(U) \subseteq {}^\perp(\tau_{J(U)}V) \cap J(U) = \mathcal{T}'$.

Now let Y be in $f_U(\mathcal{T})$, and consider the canonical sequence

$$0 \rightarrow t_U(V) \rightarrow V \rightarrow f_U(V) \rightarrow 0$$

which, after applying $\text{Hom}(\cdot, \text{Gen } Y \cap J(U))$ gives rise to an exact sequence

$$\text{Hom}(t_U(V), \text{Gen } Y \cap J(U)) \rightarrow \text{Ext}^1(\bar{V}, \text{Gen } Y \cap J(U)) \rightarrow \text{Ext}^1(V, \text{Gen } Y \cap J(U)).$$

Since Y is in ${}^\perp(\tau V)$, we have $\text{Ext}^1(V, \text{Gen } Y) = 0$ by Lemma 2.1, so in particular $\text{Ext}^1(V, \text{Gen } Y \cap J(U)) = 0$. Since $t_U(V)$ is in $\text{Gen } U$ and $J(U) \subseteq U^\perp$, we have that $\text{Hom}(t_U(V), \text{Gen } Y \cap J(U)) = 0$. Hence, we also have $\text{Ext}^1(\bar{V}, \text{Gen } Y \cap J(U)) = 0$, so $\text{Ext}^1(\bar{V}, \text{Gen}_{J(U)} Y) = 0$ which implies $\text{Hom}(Y, \tau_{J(U)}\bar{V}) = 0$, by Lemma 2.1. Hence we have that Y is in $\mathcal{T}' = {}^\perp(\tau_{J(U)}V) \cap J(U)$, which gives $f_U(\mathcal{T}) \subseteq \mathcal{T}'$.

For full subcategories \mathcal{X} and \mathcal{Y} of $\text{mod } \Lambda$, we let $\mathcal{X} * \mathcal{Y}$ denote the full subcategory

$$\{M \in \text{mod } \Lambda \mid \text{There is an exact sequence } 0 \rightarrow X \rightarrow M \rightarrow Y \rightarrow 0 \text{ with } X \in \mathcal{X}, Y \in \mathcal{Y}\}.$$

Since $f_U(\mathcal{T}) \subseteq \mathcal{T}'$, we have $\text{Gen } U * f_U(\mathcal{T}) \subseteq \text{Gen } U * \mathcal{T}'$. Since $f_U(\mathcal{T}) = \mathcal{T} \cap U^\perp$, it follows from [9, Theorem 3.12] that $\text{Gen } U * f_U(\mathcal{T}) = \mathcal{T}$, so we have $\mathcal{T} \subseteq \text{Gen } U * \mathcal{T}'$, and we aim to prove equality.

We first claim that U is Ext-projective in $\text{Gen } U * \mathcal{T}'$. Since $\text{Hom}(U, \tau U) = 0$, we have $\text{Ext}^1(U, \text{Gen } U) = 0$. We have $\mathcal{T}' \subseteq J(U)$, so $\text{Hom}(\mathcal{T}', \tau U) = 0$ and hence $\text{Ext}^1(U, \mathcal{T}') = 0$. From this we obtain that also $\text{Ext}^1(U, \text{Gen } U * \mathcal{T}') = 0$, as required.

We next claim that V is Ext-projective in $\text{Gen } U * \mathcal{T}'$. Note first that by [9, Theorem 3.12] we have $(\text{Gen } U * \mathcal{T}') \cap U^\perp = \mathcal{T}'$. Since \bar{V} is $\tau_{J(U)}$ -rigid in $J(U)$ and $\mathcal{T}' = {}^\perp(\tau_{J(U)}\bar{V}) \cap J(U)$, we have that \bar{V} is in $\mathcal{P}(\mathcal{T}')$ by [1, Theorem 2.10]. By [9, Theorem 3.15] we have $\mathcal{P}(\mathcal{T}') = f_U\mathcal{P}(\text{Gen } U * \mathcal{T}')$, and hence there is V' in $\mathcal{P}(\text{Gen } U * \mathcal{T}')$ such

that $\bar{V} = f_U(V')$. We claim that $V' \amalg U$ is τ -rigid. Since V' is in $\mathcal{P}(\text{Gen } U * \mathcal{T}')$, we have $\text{Hom}(V', \tau V') = 0$ by [1, Proposition 1.2(c)] (noting that \mathcal{T}' is functorially finite in $J(U)$ by [1, Theorem 2.10] and therefore $\text{Gen } U * \mathcal{T}'$ is functorially finite in $\text{mod } \Lambda$ by [9, Theorem 3.14]).

Since $\text{Ext}^1(V', \text{Gen } U) = 0$, we have that $\text{Hom}(U, \tau V') = 0$. We also have that $\text{Hom}(\text{Gen } U, \tau U) = 0$, since U is τ -rigid and since $\mathcal{T}' \subseteq J(U)$ we have $\text{Hom}(\mathcal{T}', \tau U) = 0$. Since V' is in $\text{Gen } U * \mathcal{T}'$ we hence have $\text{Hom}(V', \tau U) = 0$, so we have proved the claim that $V' \amalg U$ is τ -rigid. Since $f_U(\text{Gen } U) = 0$, we may assume that V' has no direct summands in $\text{Gen } U$. We have $\bar{V} = f_U(V) = f_U(V')$. It follows from [4, Lemmas 5.6, 5.7] that \bar{V} is basic, since V is basic by assumption. Similarly, also V' is basic and $V \simeq V'$. So we have proved the claim that V is in $\mathcal{P}(\text{Gen } U * \mathcal{T}')$.

Now, using that $U \amalg V$ is in $\mathcal{P}(\text{Gen } U * \mathcal{T}')$ in combination with [1, Proposition 2.9], gives that $\text{Gen } U * \mathcal{T}' \subseteq \mathcal{T} = {}^\perp(\tau U \amalg \tau V)$, and hence we obtain $\mathcal{T} = \text{Gen } U * \mathcal{T}'$, which implies $f_U(\mathcal{T}) = f_U(\text{Gen } U * \mathcal{T}') = \mathcal{T}'$, and this finishes the proof of (a).

Part (b) follows from part (a) and Lemma 5.3. \square

Lemma 5.5. *Let $U \amalg V$ be τ -rigid in $\text{mod } \Lambda$, let $\mathcal{T} = \text{Gen}(U \amalg V)$ and let $\mathcal{T}' = \text{Gen } f_U(V) \cap J(U) = \text{Gen}_{J(U)} f_U(V)$. Then $f_U(\mathcal{T}) = \mathcal{T}'$.*

Proof. Since $\text{Hom}(U \amalg V, \tau U) = 0$, we have $\mathcal{T} \subseteq {}^\perp(\tau U)$, so we have

$$\text{Gen } U \subseteq \mathcal{T} \subseteq {}^\perp(\tau U).$$

By [9, Theorem 3.15], we have that $f_U(\mathcal{T}) = \mathcal{T} \cap U^\perp$ is a torsion class in $J(U)$.

Let Y be in $f_U(\mathcal{T})$ and let $T \in \mathcal{T}$ be such that $Y = f_U(T)$. There is an epimorphism $U' \amalg V' \xrightarrow{a} T$ with $U' \in \text{add } U$ and $V' \in \text{add } V$. The canonical maps $U' \amalg V' \xrightarrow{c} f_U(V')$ and $T \xrightarrow{d} f_U(T)$ are also epimorphisms, and there is a commutative diagram

$$\begin{array}{ccc} U' \amalg V' & \xrightarrow{a} & T \\ \downarrow c & & \downarrow d \\ f_U(V') & \xrightarrow{b} & f_U(T) \end{array}$$

where $b = f_U(a)$.

Since $bc = da$ is an epimorphism, also b must be an epimorphism, and hence $Y = f_U(T)$ is in $\text{Gen } f_U(V)$. Since $f_U(\mathcal{T}) \subseteq J(U)$, we have that Y is in $J(U)$ and hence in $\text{Gen } f_U(V) \cap J(U)$.

Conversely suppose Y is in $\text{Gen } f_U(V) \cap J(U)$. Since $f_U(V)$ is a factor module of V , we have that Y is in $\text{Gen } V$, so Y is in \mathcal{T} . Since Y is in $J(U) \subseteq U^\perp$, we hence have that Y is in $\mathcal{T} \cap U^\perp = f_U(\mathcal{T})$. This finishes the proof of the lemma. \square

We also need the following reformulation of Lemma 3.8.

Lemma 5.6. *Let P, V be in $\text{mod } \Lambda$, with P projective and $\text{Hom}(P, V) = 0$, and let $\bar{V} = f_P V = V$. Let $\mathcal{T} = {}^\perp(\tau V)$ and let $\mathcal{T}' = {}^\perp(\tau_{P^\perp} \bar{V}) \cap P^\perp$. Then we have $f_P(\mathcal{T}) = \mathcal{T}'$.*

Proof. This follows directly from Lemma 3.8, using that $f_P(\mathcal{T}) = P^\perp \cap {}^\perp(\tau V)$. \square

Finally, we need the following. Suppose that \mathcal{U} and \mathcal{V} are objects in $C(\Lambda)$, with $\text{add } \mathcal{U} \cap \text{add } \mathcal{V} = 0$ and such that $\mathcal{U} \amalg \mathcal{V}$ is support τ -rigid. Note that the domain of $\mathcal{E}_{\mathcal{U} \amalg \mathcal{V}}$ is:

$$\{X \in \text{ind } C(\Lambda) : X \amalg \mathcal{U} \amalg \mathcal{V} \text{ support } \tau\text{-rigid and } \text{add } X \cap \text{add}(\mathcal{U} \amalg \mathcal{V}) = 0\}.$$

Then:

Lemma 5.7. *Let \mathcal{U} and \mathcal{V} be objects in $C(\Lambda)$ such that $\mathcal{U} \amalg \mathcal{V}$ is support τ -rigid and $\text{add } \mathcal{U} \cap \text{add } \mathcal{V} = 0$. Then $\mathcal{E}_{\mathcal{U}}$ induces a bijection between the sets:*

$$\{X \in \text{ind } C(\Lambda) : X \amalg \mathcal{U} \amalg \mathcal{V} \text{ support } \tau\text{-rigid and } \text{add } X \cap \text{add}(\mathcal{U} \amalg \mathcal{V}) = 0\},$$

and

$$\{X \in \text{ind } C(J(\mathcal{U})) : X \amalg \mathcal{E}_{\mathcal{U}}(\mathcal{V}) \text{ support } \tau\text{-rigid and } \text{add } X \cap \text{add}(\mathcal{E}_{\mathcal{U}}(\mathcal{V})) = 0\}.$$

Proof. This follows from Theorem 3.7. \square

Corollary 5.8. *The composition $\mathcal{E}_{\mathcal{E}_{\mathcal{U}}(\mathcal{V})}^{J(\mathcal{U})} \mathcal{E}_{\mathcal{U}}$ is a well-defined map with domain coinciding with the domain of $\mathcal{E}_{\mathcal{U} \amalg \mathcal{V}}$.*

Proof. This follows from Lemma 5.7 and the fact that target set in Lemma 5.7 is exactly the domain of $\mathcal{E}_{\mathcal{E}_{\mathcal{U}}(\mathcal{V})}^{J(\mathcal{U})}$. \square

The following sections will be devoted to proving the following theorem (Theorem 1.6 from Section 0.2).

Theorem 5.9. *Let \mathcal{U} and \mathcal{V} be support τ -rigid objects in $C(\Lambda)$ with no common direct summands, and suppose that $\mathcal{U} \amalg \mathcal{V}$ is support τ -rigid in $C(\Lambda)$. Then*

$$(25) \quad \mathcal{E}_{\mathcal{E}_{\mathcal{U}}(\mathcal{V})}^{J(\mathcal{U})} \mathcal{E}_{\mathcal{U}} = \mathcal{E}_{\mathcal{U} \amalg \mathcal{V}}$$

Proof We assume that $\mathcal{U} = U \amalg P[1]$ and $\mathcal{V} = V \amalg Q[1]$, with U, V, P, Q modules and P, Q projective, $\text{add}(\mathcal{U}) \cap \text{add}(\mathcal{V}) = 0$ and $\mathcal{U} \amalg \mathcal{V}$ support τ -rigid. In view of Corollary 5.8, we need to show that $\mathcal{E}_{\mathcal{E}_{\mathcal{U}}(\mathcal{V})}^{J(\mathcal{U})} \mathcal{E}_{\mathcal{U}}(X) = \mathcal{E}_{\mathcal{U} \amalg \mathcal{V}}(X)$ for each indecomposable object X in the domain

$$\{X \in \text{ind } C(\Lambda) : X \amalg \mathcal{U} \amalg \mathcal{V} \text{ support } \tau\text{-rigid and } \text{add } X \cap \text{add}(\mathcal{U} \amalg \mathcal{V}) = 0\}$$

of each of the maps $\mathcal{E}_{\mathcal{E}_{\mathcal{U}}(\mathcal{V})}^{J(\mathcal{U})} \mathcal{E}_{\mathcal{U}}$ and $\mathcal{E}_{\mathcal{U} \amalg \mathcal{V}}$.

Our strategy is to employ a case analysis, based on the properties of \mathcal{U} , \mathcal{V} and X , since the maps $\mathcal{E}_{\mathcal{U}}$, $\mathcal{E}_{\mathcal{E}_{\mathcal{U}}(\mathcal{V})}^{J(\mathcal{U})}$ and $\mathcal{E}_{\mathcal{U} \amalg \mathcal{V}}$ are defined via cases. We will consider the following cases for \mathcal{U} and \mathcal{V} .

Case I	$\mathcal{U} = U$ and $\mathcal{V} = V$
Case II	$\mathcal{U} = U$ and $\mathcal{V} = Q[1]$
Case III	$\mathcal{U} = P[1]$ and $\mathcal{V} = V$
Case IV	$\mathcal{U} = P[1]$ and $\mathcal{V} = Q[1]$

In case II, we assume that $\mathcal{U} = U$ lies in $\text{mod } \Lambda$ and that $\mathcal{V} = Q[1]$, where Q lies in $\mathcal{P}(\Lambda) \cap {}^\perp U$. In this case the claim that equation (25) holds follows directly, since we have

$$\mathcal{E}_{\mathcal{U} \amalg \mathcal{V}} = \mathcal{E}_{U \amalg P[1]} = \mathcal{E}_{\mathcal{E}_U(P[1])}^{J(U)} \mathcal{E}_U = \mathcal{E}_{\mathcal{E}_U(\mathcal{V})}^{J(U)} \mathcal{E}_U = \mathcal{E}_{\mathcal{E}_U(\mathcal{V})}^{J(\mathcal{U})} \mathcal{E}_U$$

where the second equality holds by definition of $\mathcal{E}_{U \amalg [1]}$. So it remains to consider cases I, III and IV.

For each of the cases I, III and IV we will also need to further subdivide according to the properties of X in $\text{ind } \mathcal{C}(\Lambda)$. We consider Case I in Section 6, Case III in Section 7 and Case IV in Section 8. Finally, we must consider the ‘mixed case’, where \mathcal{U} and \mathcal{V} have both module and shifted projective direct summands; this is considered in Section 9.

6. PROOF OF THEOREM 5.9, CASE I

We assume that $\mathcal{U} = U$ and $\mathcal{V} = V$ for U, V in $\text{mod } \Lambda$ where $U \amalg V$ is τ -rigid. We divide Case I into the following subcases.

- Case I*: $\mathcal{U} = U$ and $\mathcal{V} = V$ where U and V lie in $\text{mod } \Lambda$ and $\text{add } V \cap \text{Gen } U = 0$.
- Case I**: $\mathcal{U} = U$ and $\mathcal{V} = V$ where U and V lie in $\text{mod } \Lambda$ and $\mathcal{V} \in \text{Gen } U$.

We firstly note that Case I will follow from these two cases:

Proposition 6.1. *Assume that (25) holds in both cases I* and I**. Then (25) holds in Case I.*

Proof. Write $V = V_1 \amalg V_2$, where V_1 is in $\text{Gen } U$, and $\text{add } V_2 \cap \text{Gen } U = 0$. Then $\mathcal{E}_{U \amalg V} = \mathcal{E}_{(U \amalg V_1) \amalg V_2}$. If $U = 0$, then also $V_1 = 0$, and the result is trivial. We therefore assume $U \neq 0$. We proceed by induction on $n = r(\text{mod } \Lambda)$. Hence we can assume that it holds for $J(U)$, since $r(J(U)) < n$.

Note that we have $\text{Gen}(U \amalg V_1) = \text{Gen } U$, so $\text{add } V_2 \cap \text{Gen}(U \amalg V_1) = 0$. Hence we have:

$$\begin{aligned}
 \mathcal{E}_{U \amalg V} &= \mathcal{E}_{(U \amalg V_1) \amalg V_2} \\
 (26) \quad &= \mathcal{E}_{\mathcal{E}_{U \amalg V_1}(V_2)}^{J(U \amalg V_1)} \mathcal{E}_{U \amalg V_1} \\
 (27) \quad &= \mathcal{E}_{\mathcal{E}_{U \amalg V_1}(V_2)}^{J(U \amalg V_1)} \mathcal{E}_{\mathcal{E}_U(V_1)}^{J(U)} \mathcal{E}_U \\
 (28) \quad &= \mathcal{E}_{\mathcal{E}_{U \amalg V_1}(V_2)}^{J_{J(U)}(\mathcal{E}_U(V_1))} \mathcal{E}_{\mathcal{E}_U(V_1)}^{J(U)} \mathcal{E}_U \\
 (29) \quad &= \mathcal{E}_{\mathcal{E}_{\mathcal{E}_U(V_1)}(V_2)}^{J_{J(U)}(\mathcal{E}_U(V_1))} \mathcal{E}_{\mathcal{E}_U(V_1)}^{J(U)} \mathcal{E}_U \\
 (30) \quad &= \mathcal{E}_{\mathcal{E}_U(V_1) \amalg \mathcal{E}_U(V_2)}^{J(U)} \mathcal{E}_U \\
 (31) \quad &= \mathcal{E}_{\mathcal{E}_U(V_1 \amalg V_2)}^{J(U)} \mathcal{E}_U \\
 &= \mathcal{E}_{\mathcal{E}_U(V)}^{J(U)} \mathcal{E}_U
 \end{aligned}$$

where (26) holds by Case I*, the equations (27) and (29) hold by Case I**, and (28) holds by Theorem 4.3. Furthermore (30) holds (in $J(U)$) by the induction assumption, while (31) holds by definition. \square

For each of the subcases I* and I**, we will need to consider the following cases for X .

- (a) $X \in \text{ind } \Lambda$ and $X \notin \text{Gen}(U \amalg V)$
- (b) $X \in \text{ind } \Lambda$ and $X \in \text{Gen}(U \amalg V) \setminus \text{Gen } U$

- (c) $X \in \text{ind } \Lambda$ and $X \in \text{Gen } U$
- (d) $X \in \text{ind } \mathcal{P}(\Lambda)[1]$

Case I*: We assume that $\mathcal{U} = U$ and $\mathcal{V} = V$ where U and V lie in $\text{mod } \Lambda$ and add $V \cap \text{Gen } U = 0$, i.e. V has no direct summands in $\text{Gen } U$. We set $\bar{V} = f_U(V)$.

Lemma 6.2. *With the above assumptions on U and V , we have that $f_U(X)$ is not in $\text{Gen } \bar{V}$ and that*

$$f_{\bar{V}}^{J(U)}(f_U(X)) \simeq f_{U \amalg V}(X)$$

for any X not in $\text{Gen}(U \amalg V)$.

Proof. Consider the composition

$$X \xrightarrow{a} f_U(X) \xrightarrow{b} f_{\bar{V}}^{J(U)} f_U(X)$$

We first claim that ba is a left $(U \amalg V)^\perp$ -approximation. Let $c: X \rightarrow Y$ be a map, with Y in $(U \amalg V)^\perp$. Since Y is in U^\perp , and a is left U^\perp -approximation, there is a map $d: f_U(X) \rightarrow Y$ such that $da = c$. Applying $\text{Hom}(_, Y)$ to the canonical sequence

$$0 \rightarrow f_{\bar{V}}^{J(U)} f_U(X) \xrightarrow{e} f_U(X) \xrightarrow{b} f_{\bar{V}}^{J(U)} f_U(X) \rightarrow 0$$

of $f_U(X)$ gives the exact sequence

$$(32) \quad \text{Hom}(f_{\bar{V}}^{J(U)} f_U(X), Y) \rightarrow \text{Hom}(f_U(X), Y) \rightarrow \text{Hom}(f_{\bar{V}}^{J(U)} f_U(X), Y).$$

We have that $f_{\bar{V}}^{J(U)} f_U(X)$ is in $\text{Gen}_{J(U)} \bar{V} = \text{Gen } \bar{V} \cap J(U) \subseteq \text{Gen } V$, since $\bar{V} = f_U(V)$ is a factor module of V . Since Y is in V^\perp , we then have that $de = 0$. Hence, by the sequence (32), there is a map $g: f_{\bar{V}}^{J(U)} f_U(X) \rightarrow Y$, such that $gb = d$. Hence $c = gba$, which proves that ba is a left $(U \amalg V)^\perp$ -approximation as claimed. We have that ba is minimal, since both a and b are epimorphisms.

The canonical map $X \rightarrow f_{U \amalg V}(X)$ is also a minimal left $(U \amalg V)^\perp$ -approximation. So we have

$$(33) \quad f_{\bar{V}}^{J(U)} f_U(X) \simeq f_{U \amalg V}(X)$$

In particular, since by assumption X is not in $\text{Gen}(U \amalg V)$, we have that $f_{\bar{V}}^{J(U)} f_U(X) \simeq f_{U \amalg V}(X)$ is non-zero, so $f_U(X)$ is not in $\text{Gen } \bar{V}$. \square

Case I* (a): We assume that X is indecomposable in $\text{mod } \Lambda$, that $X \amalg U \amalg V$ is a τ -rigid module, and that X does not lie in $\text{Gen}(U \amalg V)$.

We then have that $\mathcal{E}_U(X) = f_U(X)$, since X does not lie in $\text{Gen } U$. Using the first claim of Lemma 6.2 we have that $f_U(X)$ is not in $\text{Gen } \bar{V}$. Hence,

$$\mathcal{E}_{\bar{V}}^{J(U)}(\mathcal{E}_U X) = f_{\bar{V}}^{J(U)}(f_U(X)).$$

Since X is not in $\text{Gen}(U \amalg V)$, we have $\mathcal{E}_{U \amalg V}(X) = f_{U \amalg V}(X)$, and equation (25) now follows from the second claim of Lemma 6.2.

Case I* (b): We assume that X is indecomposable in $\text{mod } \Lambda$, that $X \amalg U \amalg V$ is a τ -rigid module, that X is in $\text{Gen}(U \amalg V)$, and that X does not lie in $\text{Gen}(U)$.

We have $\mathcal{E}_U(X) = f_U(X)$, since by assumption, X is not in $\text{Gen } U$. By Lemma 5.5, we have that $f_U \text{Gen}(U \amalg V) = \text{Gen } f_U(V) \cap J(U) = \text{Gen}_{J(U)}(f_U(V))$. Hence we have that $f_U(X)$ is in $\text{Gen}_{J(U)}(f_U(V))$.

There is a right exact sequence

$$(34) \quad Y_{f_U(X)} \rightarrow \overline{V}'_{f_U(X)} \rightarrow f_U(X) \rightarrow 0,$$

where the first map is a minimal left add U -approximation, the second map is a minimal right add U -approximation, and $Y_{f_U(X)}$ lies in $\mathcal{P}(\perp(\tau_{J(U)}\overline{V}) \cap J(U))$. We then have $\mathcal{E}_{\overline{V}}^{J(U)}\mathcal{E}_U(X) = f_{\overline{V}}^{J(U)}(Y_{f_U(X)})$.

To compute $\mathcal{E}_{U \amalg V}(X)$, consider the right exact sequence

$$(35) \quad Y'_X \rightarrow U' \amalg V' \xrightarrow{\alpha} X \rightarrow 0$$

where the first map is a minimal left add $(U \amalg V)$ -approximation, the second map is a minimal right add $(U \amalg V)$ -approximation and Y'_X lies in $\mathcal{P}(\perp(\tau U \amalg \tau V))$. We then have $\mathcal{E}_{U \amalg V}(X) = f_{U \amalg V}(Y'_X)$.

We now aim to prove the following.

Claim 6.3. *Applying f_U to the right exact sequence (35) gives the right exact sequence (34).*

To prepare for the proof of Claim 6.3, we consider first a more general set-up.

Lemma 6.4. *Let $(\mathcal{T}, \mathcal{F})$ be an arbitrary torsion pair in $\text{mod } \Lambda$. Assume that there is commutative diagram*

$$\begin{array}{ccccccc} A & \xrightarrow{a} & B & \xrightarrow{b} & C & \longrightarrow & 0 \\ \downarrow x & & \downarrow y & & \downarrow z & & \\ A' & \xrightarrow{a'} & B' & \xrightarrow{b'} & C' & \longrightarrow & 0 \end{array}$$

where the vertical maps are minimal left \mathcal{F} -approximations (and hence epimorphisms), and the upper row is a right exact sequence. Then the map b' is an epimorphism and for any Z in \mathcal{F} , and any map $t: B' \rightarrow Z$ with $ta' = 0$, there is a map $u: C' \rightarrow Z$, such that $ub' = t$.

Proof. For the first claim, note that $zb = b'y$ is an epimorphism, hence also b' is an epimorphism. We have that $zba = 0$, and this implies $b'a'x = 0$ and hence $b'a' = 0$, since x is an epimorphism. Now $ta' = 0$ implies $ta'x = 0$, and hence $tya = 0$. Since b is the cokernel of a , there is a map $u': C \rightarrow Z$ such that $ty = u'b$. Since z is an \mathcal{F} -approximation, and by assumption Z is in \mathcal{F} , there is $u: C' \rightarrow Z$ such that $u' = uz$. It then follows that $ty = u'b = uz b = ub'y$, and since y is an epimorphism, we have $t = ub'$, as claimed. \square

Proof of Claim 6.3. Apply f_U to the sequence (35), and consider the commutative diagram

$$\begin{array}{ccccccc}
& & & 0 & & 0 & \\
& & & \downarrow & & \downarrow & \\
& & & t_U(U' \amalg V') & \xrightarrow{\gamma} & t_U(X) & \\
& & & \downarrow c & & \downarrow d & \\
Y'_X & \xrightarrow{s'} & U' \amalg V' & \xrightarrow{\alpha} & X & \longrightarrow & 0 \\
\downarrow r & & \downarrow p & & \downarrow q & & \\
f_U(Y'_X) & \xrightarrow{s} & f_U(V') & \xrightarrow{\beta} & f_U(X) & \longrightarrow & 0 \\
\downarrow & & \downarrow & & \downarrow & & \\
0 & & 0 & & 0 & &
\end{array}$$

where the second row is sequence (35), so is exact, and the second and third columns are the canonical sequences for $U' \amalg V'$ and X , respectively. Note that the map γ exists since $q\alpha c = \beta p c = 0$, so αc factors through d .

We first claim that all objects in the third row are in $J(U)$. This follows from the fact that all objects in sequence (35) are in ${}^\perp(\tau U)$, and hence the same hold for all objects in the third row, since ${}^\perp(\tau U)$ is closed under factor objects. All objects in the third row are by definition in U^\perp , and hence also in $J(U) = U^\perp \cap {}^\perp(\tau U)$.

We next claim that β is the cokernel of s . We have $\beta s = f_U(\alpha s') = f_U(0) = 0$. It now follows from applying Lemma 6.4, with the torsion pair $(\text{Gen } U, U^\perp)$, and using that $J(U) \subseteq U^\perp$ that β is the cokernel of s in $J(U)$, that is the sequence

$$(36) \quad f_U(Y'_X) \xrightarrow{s} f_U(V') \xrightarrow{\beta} f_U(X) \rightarrow 0$$

is exact in $J(U)$ (and hence also an exact sequence in $\text{mod } \Lambda$).

We now claim that the map s is a minimal left $\text{add } \bar{V}$ -approximation. For this let $b: f_U(Y'_X) \rightarrow \bar{V}''$ be a map with \bar{V}'' in $\text{add } \bar{V}$. Let V'' in $\text{add } V$ be such that $f_U(V'') = \bar{V}''$, and let $g: V'' \rightarrow \bar{V}''$ be the canonical epimorphism. Consider the canonical exact sequence for V'' ,

$$0 \rightarrow t_U(V'') \rightarrow V'' \xrightarrow{g} \bar{V}'' \rightarrow 0$$

and note that since $\text{Gen } U \subseteq {}^\perp(\tau U \amalg \tau V)$, all terms in the sequence are in ${}^\perp(\tau U \amalg \tau V)$. Applying $\text{Hom}(Y'_X, _)$ gives the exact sequence

$$\text{Hom}(Y'_X, V'') \rightarrow \text{Hom}(Y'_X, \bar{V}'') \rightarrow \text{Ext}^1(Y'_X, t_U(V'')).$$

Using that Y'_X is in $\mathcal{P}({}^\perp(\tau U \amalg \tau V))$ and that $t_U(V'') \in \text{Gen } U \subseteq {}^\perp(\tau U \amalg \tau V)$, we have that the last term vanishes, and hence the first map is surjective, and therefore there is a map $f': Y'_X \rightarrow V''$ such that $gf' = br$.

The map s' is an $\text{add}(U \amalg V)$ -approximation, therefore there is a map $f: U' \amalg V' \rightarrow V''$, such that $f' = fs'$. Then we have $br = gf' = gfs'$. Now consider the canonical sequence

$$(37) \quad 0 \rightarrow t_U(U' \amalg V') \rightarrow U' \amalg V' \xrightarrow{p} f_U(U' \amalg V') = f_U(V') \rightarrow 0$$

Note that $\text{Hom}(t_U(U' \amalg V'), \bar{V}'') = 0$ since \bar{V} is in ${}^\perp(\tau U)$. Applying $\text{Hom}(\cdot, \bar{V}'')$ to (37), we obtain

$$\text{Hom}(f_U(V'), \bar{V}'') \simeq \text{Hom}(U' \amalg V', \bar{V}'').$$

Hence there is a map $e: f_U(V') \rightarrow \bar{V}''$, such that $ep = gf$. We then have $br = gfs' = eps' = esr$, that is $(b - es)r = 0$. Since r is an epimorphism, this implies $b = es$, and we have proved the claim that s is a left add \bar{V} -approximation.

We next claim that s is a left minimal map. Let

$$s = \begin{pmatrix} s_1 \\ 0 \end{pmatrix}: f_U(Y'_X) \rightarrow f_U(V')$$

where s_1 is left minimal. Then $\text{coker}(s) \simeq \text{coker}(s_1) \amalg M$ for some module M . Note that X indecomposable implies that $\text{coker}(s) \simeq f_U(X)$ is indecomposable by [4, Lemma 4.6], hence we have $\text{coker}(s_1) = 0$ or $M = 0$. If $\text{coker}(s_1) = 0$, then $f_U(X)$ is in $\text{add } f_U(V')$. Then there is an indecomposable direct summand V_0 of V , such that $f_U(X) \simeq f_U(V_0)$. But then, by [4, Lemma 5.7] we have $X \simeq V_0$, but this is a contradiction, since X is by assumption not in $\text{add } V$. Hence $M = 0$, and therefore $s = s_1$ is left minimal.

We claim that $f_U(Y'_X)$ is in $\mathcal{P}(\mathcal{T}')$, where $\mathcal{T}' = {}^\perp(\tau_{J(U)}\bar{V}) \cap J(U)$ is a torsion class in $J(U)$. For this consider the torsion class $\mathcal{T} = {}^\perp(\tau U \amalg \tau V)$ in $\text{mod } \Lambda$. By Lemma 5.4 we have that $f_U(\mathcal{T}) = \mathcal{T}'$. It then follows from [9, Theorem 3.15], that $f_U(\mathcal{P}(\mathcal{T})) = \mathcal{P}(\mathcal{T}')$, and hence $f_U(Y'_X)$ is in $\mathcal{P}(\mathcal{T}')$, since $Y'_X \in \mathcal{P}(\mathcal{T})$.

We can now apply [4, Proposition 4.7] to obtain that the sequences (34) and (36) are isomorphic, and this concludes the proof of the claim. \square

Using Claim 6.3 we obtain

$$\mathcal{E}_{\bar{V}}^{J(U)} \mathcal{E}_U(X) = \mathcal{E}_{\bar{V}}^{J(U)} f_U(X) = f_{\bar{V}}^{J(U)}(Y_{f_U(X)}) = f_{\bar{V}}^{J(U)} f_U(Y'_X).$$

Moreover, we have that Y'_X is not in $\text{Gen}(U \amalg V)$, since $\mathcal{E}_{U \amalg V}(X) = f_{U \amalg V}(Y'_X) \neq 0$. Therefore, using Lemma 6.2, we obtain

$$\mathcal{E}_{U \amalg V}(X) = f_{U \amalg V}(Y'_X) \simeq f_{\bar{V}}^{J(U)} f_U(Y'_X).$$

This finishes the proof that equation (25) holds in this case.

Case I* (c): We assume that X is an indecomposable module in $\text{mod } \Lambda$, that $X \amalg U \amalg V$ is a τ -rigid module and that X is in $\text{Gen}(U)$.

Let $\mathcal{T} = {}^\perp(\tau U \amalg \tau V)$ and $\mathcal{T}' = {}^\perp(\tau_{J(U)}\bar{V}) \cap J(U)$, and consider the exact sequence

$$(38) \quad Y'_X \rightarrow U_X \rightarrow X \rightarrow 0,$$

where the first map is a minimal left add U -approximation, and the second map is a minimal right add U -approximation. We then have that $\mathcal{E}_U(X) = \mathcal{E}_U(X) = f_U(Y'_X)[1]$. Note that by Theorem 3.7 the object $\mathcal{E}_U(X \amalg V) = f_U(Y'_X)[1] \amalg \bar{V}$ is support τ -rigid in $\mathcal{C}(J(\mathcal{U}))$, and hence we have that $f_U(Y'_X)$ is in ${}^\perp \bar{V}$.

We have $\mathcal{E}_{\mathcal{E}_U(V)}^{J(\mathcal{U})} \mathcal{E}_U(X) = \mathcal{E}_{\bar{V}}^{J(\mathcal{U})} \mathcal{E}_U(X) = f_{\bar{V}}^{J(U)} Y''_X$, where $f_U(Y'_X) \rightarrow Y''_X$ is a minimal left \mathcal{T}' -approximation.

To compute $\mathcal{E}_{U \amalg V}(X)$ we consider the right exact sequence

$$(39) \quad Y_X''' \rightarrow U_X' \amalg V_X' \rightarrow X \rightarrow 0$$

where the first map is a minimal left $\text{add}(U \amalg V)$ -approximation, and the second map is a minimal right $\text{add}(U \amalg V)$ -approximation. Then $\mathcal{E}_{U \amalg V}(X) = \mathcal{E}_{U \amalg V}(X) = f_{U \amalg V}(Y_X''')$.

Since $\mathcal{E}_{U \amalg V}(X) \neq 0$, we have that Y_X''' is not in $\text{Gen}(U \amalg V)$. By Lemma 6.2, we hence have that $\mathcal{E}_{U \amalg V}(X) = f_{U \amalg V}(Y_X''')[1] = f_{\bar{V}}^{J(U)} f_U(Y_X''')[1]$.

It is therefore sufficient to prove that

$$f_{\bar{V}}^{J(U)}(Y_X''') \simeq f_{\bar{V}}^{J(U)} f_U(Y_X''')$$

The main steps are as follows:

- Claim 6.5.** (a) *There is a map $Y_X' \rightarrow U_X \amalg Y_X'''$, which is a left \mathcal{T} -approximation.*
 (b) *The induced map $f_U(Y_X') \rightarrow f_U(Y_X''')$ is a left \mathcal{T}' -approximation.*
 (c) *We have that Y_X'' is a direct summand in $f_U(Y_X''')$*
 (d) *We have $f_{\bar{V}}^{J(U)}(Y_X''') \simeq f_{\bar{V}}^{J(U)} f_U(Y_X''')$.*

Proof. (a): Consider the diagram

$$\begin{array}{ccccccc} \mathbb{P}_{Y_X'} & \longrightarrow & \mathbb{P}_{U_X} & \longrightarrow & \mathbb{P}_X & \longrightarrow & \mathbb{P}_{Y_X'}[1] \\ & & & & \downarrow & = & \\ \mathbb{P}_{Y_X'''} & \longrightarrow & \mathbb{P}_{U_X' \amalg V_X'} & \longrightarrow & \mathbb{P}_X & \longrightarrow & \mathbb{P}_{Y_X'''}[1] \end{array}$$

where the rows are triangles giving rise (by taking homology) to the exact sequences (38) and (39), respectively (see Section 3). We have that U, Y_X''' are in $\mathcal{P}(\mathcal{T})$, so in particular $\text{Hom}(Y_X''', \tau U) = 0$. Hence by Lemma 2.1 we have that $\text{Hom}(\mathbb{P}_{U_X}, \mathbb{P}_{Y_X'''}[1]) = 0$. Therefore (see [11, Section 1.4]) the above diagram can be completed to a commutative diagram in such a way that there is an induced triangle

$$\mathbb{P}_{U_X' \amalg V_X'}[-1] \xrightarrow{g} \mathbb{P}_{Y_X'} \xrightarrow{h} \mathbb{P}_{U_X' \amalg Y_X'''} \rightarrow \mathbb{P}_{U_X' \amalg V_X'}$$

Now, let $k: \mathbb{P}_{Y_X'} \rightarrow \mathbb{P}_T$ be a map with $T \in \mathcal{T} = {}^\perp(\tau U \amalg \tau V)$. Then $\text{Hom}(T, \tau U \amalg \tau V) = 0$, and hence by Lemma 2.1, we have $\text{Hom}(\mathbb{P}_{U_X' \amalg V_X'}, \mathbb{P}_T[1]) = 0$. Hence we have $kg = 0$, so by exactness of $\text{Hom}(\cdot, \mathbb{P}_T)$ we have that there is map $l: \mathbb{P}_{U_X' \amalg Y_X'''} \rightarrow \mathbb{P}_T$, such that $lh = k$. It then follows that the map $H^0(k): Y_X' \rightarrow T$ factors through $H^0(l)$, and by Lemma 2.1, we have that any map $Y_X' \rightarrow T$ factors through $Y_X' \rightarrow U_X' \amalg Y_X'''$. This finishes the proof of the claim.

(b): This follows directly from Lemma 5.4, using that $U, Y_X''' \in \mathcal{P}(\mathcal{T})$ and therefore $\text{Hom}(U, \tau Y_X''') = 0$.

(c): This follows directly from (b), using that $f_U(Y_X') \rightarrow Y_X''$ is a minimal left \mathcal{T}' -approximation.

(d): This follows directly from (c), using that both $f_{\bar{V}}^{J(U)} Y_X''$ and $f_{\bar{V}}^{J(U)} f_U(Y_X''')$ are indecomposable. \square

So equation (25) is proved for this case.

Case I* (d): We assume that X is of the form $R[1]$, where R is an indecomposable module in $(\mathcal{P}(\Lambda) \cap {}^\perp(U \amalg V))$.

Let $\mathcal{T} = {}^\perp(\tau U \amalg \tau V)$ and let $\mathcal{T}' = {}^\perp(\tau_{J(U)}\bar{V}) \cap J(U)$.

We first compute $\mathcal{E}_{U \amalg V}(X)$. For this, let $R \rightarrow Y_R$ be a minimal left \mathcal{T} -approximation. Then $\mathcal{E}_{U \amalg V}(X) = f_{U \amalg V}(Y_R)[1] = f_{\bar{V}}^{J(U)} f_U(Y_R)[1]$, where the last equation follows from Lemma 6.2.

Similarly, we compute $\mathcal{E}_U(X)$ by letting $R \rightarrow Y'_R$ be a minimal left \mathcal{T} -approximation and then $\mathcal{E}_U(X) = f_U(Y'_R)[1]$.

To compute $\mathcal{E}_{\bar{V}}^{J(U)} \mathcal{E}_U(X)$, let $f_U(Y'_R) \rightarrow Y''_R$ be a minimal \mathcal{T}' -approximation. Then we have $\mathcal{E}_{\bar{V}}^{J(U)} \mathcal{E}_U(X) = f_{\bar{V}}^{J(U)}(Y''_R)[1]$.

Claim 6.6. *We have that $f_U(Y_R) \simeq Y''_R$.*

Proof. We first claim that the composition $R \xrightarrow{h} Y_R \xrightarrow{k} f_U(Y_R)$ is a left $\mathcal{P}(\mathcal{T}')$ -approximation. Consider a map $f: R \rightarrow N'$ with N' in $\mathcal{P}(\mathcal{T}')$. By Lemma 5.4(b), we have that $f_U \mathcal{P}(\mathcal{T}) = \mathcal{P}(\mathcal{T}')$, so there is a module N in $\mathcal{P}(\mathcal{T})$ satisfying $f_U(N) = N'$. Since R is projective, there is a map $u: R \rightarrow N$, such that $gu = f$. Since $h: R \rightarrow Y_R$ is a $\mathcal{P}(\mathcal{T})$ -approximation and N is in $\mathcal{P}(\mathcal{T})$, there is $v: Y_R \rightarrow N$ such that $u = vh$. Note that N is in $\mathcal{P}(\mathcal{T}) \subseteq J(U) \subseteq U^\perp$. Since k is a left U^\perp -approximation, there is $w: f_U(Y_R) \rightarrow N$ such that $v = wk$. So, we have $f = gu = gvh = gwhk$. Note that N' is in $\mathcal{P}(\mathcal{T}') \subseteq J(U) \subseteq U^\perp$. Since k is a left U^\perp -approximation, there is $w': f_U(Y_R) \rightarrow N$ such that $gv = wk$. So, we have $f = gu = gvh = wkh$, and hence kh is a left $\mathcal{P}(\mathcal{T}')$ -approximation.

Next, we claim that the composition

$$R \xrightarrow{b} Y'_R \xrightarrow{c} f_U(Y'_R) \xrightarrow{d} Y''_R$$

is a left $\mathcal{P}(\mathcal{T}')$ -approximation. Let N be in $\mathcal{P}(\mathcal{T}')$, and let $a: R \rightarrow N$ be a map. Since N is in ${}^\perp(\tau U)$ and b is a left ${}^\perp(\tau U)$ -approximation, there is a map $a': Y'_R \rightarrow N$ such that $a = a'b$. Since N is in U^\perp and c is a left U^\perp -approximation, there is a map $a'': f_U(Y'_R) \rightarrow N$ such that $a' = a''c$. Since N is in \mathcal{T}' and d is a left \mathcal{T}' -approximation, there is a map $a''': Y''_R \rightarrow N$ such that $a'' = a'''d$. So we have $a'''dcb = a''cb = a'b = a$, so dcb is a left $\mathcal{P}(\mathcal{T}')$ -approximation as claimed.

Note that both Y_R and Y''_R are indecomposable by [4, Prop. 3.7], hence also $f_U(Y_R)$ is indecomposable, by [4, Lemma 4.6]. It then follows that both

$$X \xrightarrow{h} Y_R \xrightarrow{k} f_U(Y_R)$$

and

$$X \xrightarrow{b} Y'_R \xrightarrow{c} f_U(Y'_R) \xrightarrow{d} Y''_R$$

are minimal left \mathcal{T}' -approximations. So we obtain $f_U(Y_R) \simeq Y''_R$. \square

By Claim 6.6 we now have that

$$\mathcal{E}_{U \amalg V}(X) = f_{\bar{V}}^{J(U)} f_U(Y_R)[1] = f_{\bar{V}}^{J(U)}(Y''_R)[1] = \mathcal{E}_{\bar{V}}^{J(U)} \mathcal{E}_U(X)$$

and hence equation (25) holds also in this case.

Case I:** We assume that $\mathcal{U} = U$ and $\mathcal{V} = V$ for U, V in $\text{mod } \Lambda$, that $U \amalg V$ is a τ -rigid module, and that V lies in $\text{Gen } U$.

Let $\bar{V} = \bar{Q}[1]$, where \bar{Q} is in $\mathcal{P}(J(U))$. We first make the following observation.

Lemma 6.7. *With the assumptions of Case I**, we have that $f_{U \amalg V} = f_U$.*

Proof. Since $V \in \text{Gen } U$, we have that $\text{Gen}(U \amalg V) = \text{Gen } U$. It follows that $t_{U \amalg V} = t_U$, and then by uniqueness of canonical sequences, that also $f_{U \amalg V} = f_U$. \square

Case I (a):** We assume that X is an indecomposable module in $\text{mod } \Lambda$, that $X \amalg U \amalg V$ is a τ -rigid module, and that X does not lie in $\text{Gen}(U \amalg V) = \text{Gen } U$.

We have that $\mathcal{E}_U(X) = f_U(X)$, and

$$\mathcal{E}_{\bar{V}}^{J(U)} \mathcal{E}_U(X) = \mathcal{E}_{\bar{V}}^{J(U)}(f_U(X)) = \mathcal{E}_{\bar{Q}[1]}^{J(U)}(f_U(X)) = f_U(X).$$

Note that the last equation holds since \bar{Q} is in $\mathcal{P}(J(U))$ and we have $\text{Hom}(\bar{Q}, f_U(X)) = 0$ since $\mathcal{E}_U(V \amalg X)$ is support τ -rigid in $\mathcal{C}(J(U))$ by Theorem 3.7.

By Lemma 6.7 we have

$$\mathcal{E}_{U \amalg V}(X) = f_{U \amalg V}(X) = f_U(X) = \mathcal{E}_{\bar{V}}^{J(U)} \mathcal{E}_U(X)$$

and the claim that equation (25) holds, is proved in this case.

Case I (b):** Since $\text{Gen}(U \amalg V) = \text{Gen } U$, this case (where X lies in $\text{ind } \Lambda$ and $X \in \text{Gen}(U \amalg V) \setminus \text{Gen } U$) cannot occur.

Case I (c):** We assume that X is an indecomposable τ -rigid module, that $X \amalg U \amalg V$ is a τ -rigid module, and that X lies in $\text{Gen } U = \text{Gen}(U \amalg V)$.

In order to compute $\mathcal{E}_U(X)$, we consider the exact sequence

$$(40) \quad Y_X \rightarrow U_X \rightarrow X \rightarrow 0$$

where the first map is a minimal left add U -approximation, and the second map is a minimal right add U -approximation. Then $\mathcal{E}_U(X) = f_U(Y_X)[1]$.

We have

$$\mathcal{E}_{\mathcal{E}_U(\mathcal{V})}^{J(U)} \mathcal{E}_U(X) = \mathcal{E}_{\mathcal{E}_{\bar{V}}}^{J(U)} \mathcal{E}_U(X) = f_{\bar{Q}}^{J(U)} f_U(Y_X)[1].$$

Next, to compute $\mathcal{E}_{U \amalg V}(X)$, we consider the exact sequence

$$(41) \quad Y'_X \rightarrow U'_X \amalg V'_X \rightarrow X \rightarrow 0,$$

where the first map is a minimal left add $(U \amalg V)$ -approximation, and the second map is a minimal right add $(U \amalg V)$ -approximation. Then $\mathcal{E}_{U \amalg V}(X) = f_{U \amalg V}(Y'_X)[1]$.

By Lemma 6.7, it now follows that

$$\mathcal{E}_{U \amalg V}(X) = f_{U \amalg V}(Y'_X)[1] = f_U(Y'_X)[1].$$

By the above, it will be sufficient to prove that

$$(42) \quad f_{\bar{Q}}^{J(U)} f_U(Y_X) \simeq f_U(Y'_X).$$

The main steps in the proof are as follows:

Claim 6.8. Let $\mathcal{T} = {}^\perp(\tau U \amalg \tau V)$ and let $\mathcal{T}' = \overline{Q}^\perp \cap J(U)$. Then the following hold.

- (a) We have $f_{\overline{Q}}^{J(U)} f_U(\mathcal{T}) = f_U(\mathcal{T}) = \mathcal{T}'$.
- (b) There is a map $Y_X \xrightarrow{\gamma} Y'_X \amalg U_X$ which is a left \mathcal{T} -approximation.
- (c) The map $f_U(Y_X) \xrightarrow{f_U(\gamma)} f_U(Y'_X)$ is a left \mathcal{T}' -approximation.
- (d) The map $f_{\overline{Q}}^{J(U)} f_U(Y_X) \xrightarrow{f_{\overline{Q}}^{J(U)} f_U(\gamma)} f_{\overline{Q}}^{J(U)} f_U(Y'_X)$ is a left $f_{\overline{Q}}^{J(U)} \mathcal{T}' = \mathcal{T}'$ -approximation.
- (e) The map $f_{\overline{Q}}^{J(U)} f_U(\gamma)$ is an isomorphism.
- (f) We have $f_{\overline{Q}}^{J(U)} f_U(Y'_X) \simeq f_U(Y'_X)$.
- (g) We have $f_{\overline{Q}}^{J(U)} f_U(Y_X) \simeq f_U(Y'_X)$.

Proof. (a): Since V is in $\text{Gen } U$, we have by Lemma 6.7 that $f_U = f_{U \amalg V}$, and we have

$$\begin{aligned} f_U(\mathcal{T}) &= f_{U \amalg V}(\mathcal{T}) = f_{U \amalg V}({}^\perp(\tau U \amalg \tau V)) = (U \amalg V)^\perp \cap {}^\perp(\tau U \amalg \tau V) \\ &= J(U \amalg V) \stackrel{(*)}{=} J_{J(U)}(\mathcal{E}_U(V)) = J_{J(U)}(\overline{Q}[1]) = \overline{Q}^\perp \cap J(U) = \mathcal{T}' \end{aligned}$$

where $(*)$ holds by Theorem 4.3. This proves the second equality. But $f_{\overline{Q}}^{J(U)}$ clearly acts as the identity on objects in $\overline{Q}^\perp \cap J(U)$, and this proves the first equality.

(b): Consider the diagram

$$\begin{array}{ccccccc} \mathbb{P}_{Y_X} & \longrightarrow & \mathbb{P}_{U_X} & \longrightarrow & \mathbb{P}_X & \longrightarrow & \mathbb{P}_{Y_X}[1] \\ & & & & \downarrow = & & \\ \mathbb{P}_{Y'_X} & \longrightarrow & \mathbb{P}_{U'_X \amalg V'_X} & \longrightarrow & \mathbb{P}_X & \longrightarrow & \mathbb{P}_{Y'_X}[1] \end{array}$$

where the rows are triangles giving rise (by taking homology) to the exact sequences (40) and (41), respectively (see Section 2 for details). Since Y'_X is in $\mathcal{T} \subseteq {}^\perp(\tau U)$, we have $\text{Hom}(Y'_X, \tau U) = 0$, and hence $\text{Hom}(\mathbb{P}_{U_X}, \mathbb{P}_{Y'_X}[1]) = 0$, by Lemma 2.1. Hence there are maps $\mathbb{P}_{U_X} \rightarrow \mathbb{P}_{U'_X \amalg V'_X}$ and $\mathbb{P}_{Y_X} \rightarrow \mathbb{P}_{Y'_X}$ completing the above diagram in such a way that there is a triangle (see [11, Section 1.4])

$$\mathbb{P}_{U'_X \amalg V'_X}[-1] \xrightarrow{g} \mathbb{P}_{Y_X} \xrightarrow{h} \mathbb{P}_{U_X \amalg Y'_X} \rightarrow \mathbb{P}_{U'_X \amalg V'_X}.$$

Now, let $k: \mathbb{P}_{Y_X} \rightarrow \mathbb{P}_T$ be a map with T in $\mathcal{T} = {}^\perp(\tau U \amalg \tau V)$. Then $\text{Hom}(T, \tau U \amalg \tau V) = 0$, and hence by Lemma 2.1, we have $\text{Hom}(\mathbb{P}_{U'_X \amalg V'_X}, \mathbb{P}_T[1]) = 0$. Hence we have $kg = 0$, so by exactness of $\text{Hom}(\cdot, \mathbb{P}_T)$ we have that there is map $l: \mathbb{P}_{U_X \amalg Y'_X} \rightarrow \mathbb{P}_T$ such that $lh = k$. It then follows that the map $H^0(k): Y_X \rightarrow T$ factors through $H^0(l)$, and by Lemma 2.1, it then follows that any map $Y_X \rightarrow T$ factors through $Y_X \rightarrow U_X \amalg Y'_X$. This finishes the proof of the claim.

(c): We have $U \in \mathcal{P}({}^\perp(\tau U))$ by [1, Proposition 2.9], and $Y_X \in \mathcal{P}({}^\perp(\tau U))$ by construction (see Definition 3.1, Case I(b)). Hence, in particular, $\text{Hom}(U, \tau Y_X) = 0$, since $\mathcal{P}({}^\perp(\tau U))$ is τ -rigid by [1, Theorem 2.10]. Then the assertion follows from Lemma 5.3.

(d): We have that $f_U(Y_X)$ is in $\mathcal{P}(J(U))$, and hence $\tau_{J(U)}f_U(Y_X) = 0$. Hence, the assertion follows from Lemma 5.3.

(e): We have that the image of the map $f_{\overline{Q}}^{J(U)}f_U$ is in $\overline{Q}^\perp \cap J(U) = \mathcal{T}'$, so in particular $f_{\overline{Q}}^{J(U)}f_U(Y_X)$ is in \mathcal{T}' . By [4, Lemma 5.6]), $f_{\overline{Q}}^{J(U)}f_U(Y'_X)$ is indecomposable, so the map $f_{\overline{Q}}^{J(U)}f_U(\gamma)$ in (d) is left minimal. The assertion follows.

(f): This follows from $f_U(Y'_X) \in f_U(\mathcal{T}) = \mathcal{T}' = \overline{Q}^\perp \cap J(U)$.

(g): This now follows from (e) and (f). \square

We have now proved that (42) holds, and hence (25) holds in this case.

Case I (d):** We assume that X is of the form $R[1]$, where R is an indecomposable module in $(\mathcal{P}(\Lambda) \cap {}^\perp(U \amalg V))$.

Note first that we have $\overline{\mathcal{V}} = \overline{Q}[1]$, where $\overline{Q} = f_U(Y_V)$ and where there is an exact sequence

$$Y_V \rightarrow U_V \rightarrow V \rightarrow 0,$$

where the first map is a minimal left add U -approximation, and the second map is a minimal right add U -approximation.

We have that $\mathcal{E}_U(X) = f_U(Y_R)[1]$, where $R \xrightarrow{\beta} Y_R$ is a minimal left ${}^\perp(\tau U)$ -approximation. Furthermore, we have

$$\mathcal{E}_{\overline{\mathcal{V}}}^{J(U)}\mathcal{E}_U(X) = \mathcal{E}_{\overline{Q}[1]}^{J(U)}\mathcal{E}_U(X) = f_{\overline{Q}}^{J(U)}f_U(Y_R)[1].$$

We have $f_U = f_{U \amalg V}$, by Lemma 6.7. We hence have that $\mathcal{E}_{U \amalg V}(X) = f_{U \amalg V}(Y'_R)[1] = f_U(Y'_R)[1]$, where $R \xrightarrow{\alpha} Y'_R$ is a left minimal ${}^\perp(\tau U \amalg \tau V)$ -approximation.

Hence, it will be sufficient to prove that

$$(43) \quad f_{\overline{Q}}^{J(U)}f_U(Y_R) \simeq f_U(Y'_R).$$

The main steps in the proof are as follows.

Claim 6.9. *Let $\mathcal{T} = {}^\perp(\tau U \amalg \tau V)$ and let $\mathcal{T}' = \overline{Q}^\perp \cap J(U)$. Then the following hold.*

- (a) *We have $f_{\overline{Q}}^{J(U)}f_U(\mathcal{T}) = f_U(\mathcal{T}) = \mathcal{T}'$.*
- (b) *There is a map $Y_R \xrightarrow{a} Y'_R$ such that $f_U(a)$ is a left $f_U(\mathcal{T})$ -approximation.*
- (c) *We have $f_{\overline{Q}}^{J(U)}f_U(Y'_R) \simeq f_U(Y'_R)$.*
- (d) *The map $f_{\overline{Q}}^{J(U)}f_U(Y_R) \xrightarrow{f_{\overline{Q}}^{J(U)}f_U(a)} f_{\overline{Q}}^{J(U)}f_U(Y'_R)$ is a left $f_{\overline{Q}}^{J(U)}\mathcal{T}' = \mathcal{T}'$ -approximation.*
- (e) *The map $f_{\overline{Q}}^{J(U)}f_U(a)$ is an isomorphism.*
- (f) *We have $f_{\overline{Q}}^{J(U)}f_U(Y'_R) \simeq f_U(Y'_R)$.*
- (g) *We have $f_{\overline{Q}}^{J(U)}f_U(Y_R) \simeq f_U(Y'_R)$.*

Proof. (a): See Claim 6.8(a).

(b): Since $\beta: R \rightarrow Y_R$ is a left ${}^\perp(\tau U)$ -approximation, and $Y'_R \in \mathcal{T} \subseteq {}^\perp(\tau U)$, there is a map $a: Y_R \rightarrow Y'_R$, such that $a\beta = \alpha$.

We claim that $f_U(a)$ is a left $f_U(\mathcal{T})$ -approximation. Consider a map $y: f_U(Y_R) \rightarrow f_U(T)$, where $f_U(T)$ is in $f_U(\mathcal{T})$. Since $U, Y_R \in \mathcal{P}({}^\perp(\tau U))$, we have in particular that $\text{Hom}(U, \tau Y_R) = 0$. It then follows from Lemma 5.1 that $y = f_U(x)$ for some $x: Y_R \rightarrow T$. Consider the diagram

$$\begin{array}{ccccc}
 & & \alpha & & \\
 & & \curvearrowright & & \\
 R & \xrightarrow{\beta} & Y_R & \xrightarrow{a} & Y'_R \\
 & \searrow^{x\beta} & \downarrow x & \swarrow c & \\
 & & T & &
 \end{array}$$

where $c: Y'_R \rightarrow T$ such that $x\beta = ca\beta$ exists since $\alpha = a\beta$ is a left \mathcal{T} -approximation. It follows that $(ca - x)\beta = 0$. Applying $\text{Hom}(_, T)$ to the right exact sequence $R \xrightarrow{\beta} Y_R \xrightarrow{a} Y'_R \rightarrow 0$ gives the left exact sequence

$$0 \rightarrow \text{Hom}(U_R, T) \rightarrow \text{Hom}(Y_R, T) \rightarrow \text{Hom}(R, T).$$

Now $(ca - x)\beta = 0$ implies that there is a map $n: U_R \rightarrow T$, such that $ca - x = nl$, and so $x = ca + nl$. Since $f_U(nl) = 0$, this gives $y = f_U(x) = f_U(c)f_U(a)$. Hence we have that $f_U(a)$ is a left $f_U(\mathcal{T})$ -approximation as claimed.

(c): This follows directly from (a), since $f_{\overline{Q}}^{J(U)}$ acts as the identity on $\overline{Q}^\perp \cap J(U)$.

(d,e,f,g): See Claim 6.8(d,e,f,g). □

We have now proved that (43) holds, and hence that equation (25) holds in this case.

7. PROOF OF THEOREM 5.9, CASE III

We have already dealt with Case II (see the end of Section 5), so we must next deal with Case III. We assume that $\mathcal{U} = P[1]$, where P lies in $\mathcal{P}(\Lambda)$, that $\mathcal{V} = V$ is a τ -rigid module satisfying $\text{Hom}(P, V) = 0$.

Then $\mathcal{E}_{\mathcal{U}}(\mathcal{V}) = \overline{\mathcal{V}} = \overline{V} = V$ is τ -rigid in $J(U) = P^\perp$.

We need in this case to consider three possible cases for X :

- (a) X lies in $\text{ind}(\Lambda)$, X does not lie in $\text{Gen } V$, $X \amalg V$ is τ -rigid and $\text{Hom}(P, X) = 0$.
- (b) X lies in $\text{ind}(\Lambda)$, X lies in $\text{Gen } V$, $X \amalg V$ is τ -rigid and $\text{Hom}(P, X) = 0$.
- (c) X is of the form $R[1]$ where R lies in $\text{ind } \mathcal{P}(\Lambda)$ and $\text{Hom}(R, V) = 0$.

Case III (a): We assume that X is an indecomposable τ -rigid module in $\text{mod } \Lambda$, that $X \amalg V$ is τ -rigid, that $\text{Hom}(P, X) = 0$ and that X does not lie in $\text{Gen } V$.

Then $\mathcal{E}_{\mathcal{U}}(X) = \mathcal{E}_{P[1]}(X) = X$, and X is τ -rigid in $J(\mathcal{U}) = P^\perp$. We have

$$\mathcal{E}_{\mathcal{E}_{\mathcal{U}}(\mathcal{V})}^{J(\mathcal{U})} \mathcal{E}_{\mathcal{U}}(X) = \mathcal{E}_{\overline{\mathcal{V}}}^{J(\mathcal{U})} \mathcal{E}_{\mathcal{U}}(X) = \mathcal{E}_V^{J(U)}(X) = f_V^{J(P[1])}(X) = f_V^{P^\perp}(X).$$

We next compute $\mathcal{E}_{\mathcal{U}\amalg\mathcal{V}}(X)$. We have $\mathcal{E}_{\mathcal{V}}(\mathcal{U}) = \mathcal{E}_{\mathcal{V}}(P[1]) = f_V(Y_P)[1]$, where $P \rightarrow Y_P$ is a left ${}^\perp(\tau V)$ -approximation. We then obtain

$$\mathcal{E}_{\mathcal{U}\amalg\mathcal{V}}(X) = \mathcal{E}_{V\amalg P[1]}(X) = \mathcal{E}_{\mathcal{E}_{\mathcal{V}}(P[1])}^{J(V)} \mathcal{E}_V(X) = \mathcal{E}_{f_V(Y_P)[1]}^{J(V)} \mathcal{E}_V(X) = \mathcal{E}_{f_V(Y_P)[1]}^{J(V)} f_V(X) = f_V(X),$$

where the second equality holds by definition. The last equation holds since $\mathcal{E}_V(X \amalg P[1]) = f_V(X) \amalg f_V(Y_P)[1]$ is support τ -rigid in $\mathcal{C}(J(U))$ by Theorem 3.7, so $f_V(Y_P)$ is in $\mathcal{P}(J(U))$ with $\text{Hom}(f_V(Y_P), f_V(X)) = 0$.

We next claim that $f_V^{P^\perp}(X) = f_V(X)$. For this, consider the canonical sequence of X in $\text{mod } \Lambda$ with respect to the torsion pair $(\text{Gen } V, V^\perp)$:

$$0 \rightarrow t_V(X) \rightarrow X \rightarrow f_V(X) \rightarrow 0.$$

Since $\text{Hom}(P, X) = 0$ by assumption, and P is projective, we also have $\text{Hom}(P, f_V(X)) = 0$, and clearly also $\text{Hom}(P, t_V(X)) = 0$. We have $t_V(X) \in \text{Gen } V \cap P^\perp = \text{Gen}_{P^\perp} V$ and $f_V(X) \in P^\perp \cap V^\perp$, so this sequence is also the canonical sequence of X in P^\perp with respect to the torsion pair $(\text{Gen}_{P^\perp} V, P^\perp \cap V^\perp)$. Hence $f_V^{P^\perp}(X) = f_V(X)$ and it follows that

$$\mathcal{E}_{\mathcal{U}\amalg\mathcal{V}}(X) = f_V(X) = f_V^{P^\perp}(X) = \mathcal{E}_{\overline{V}}^{J(U)} \mathcal{E}_{\mathcal{U}}(X),$$

and equation (25) holds also in this case.

Case III (b): We assume that $X \amalg V$ is a τ -rigid module in $\text{mod } \Lambda$ such that X lies in $\text{Gen } V$ and $\text{Hom}(P, X) = 0$.

First note that $\mathcal{E}_{\mathcal{U}}(X) = \mathcal{E}_{P[1]}(X) = X$. Consider the right exact sequence in $J(\mathcal{U}) = P^\perp$,

$$(44) \quad Y_X^{P^\perp} \rightarrow V_X^{P^\perp} \rightarrow X \rightarrow 0,$$

where the first map is a minimal left add V -approximation in P^\perp , $Y_X \in \mathcal{P}({}^\perp(\tau V))$ and the second map is a minimal right add V -approximation in P^\perp .

We then have that

$$\mathcal{E}_{\mathcal{E}_{\mathcal{U}}(\mathcal{V})}^{J(\mathcal{U})} \mathcal{E}_{\mathcal{U}}(X) = \mathcal{E}_{\overline{V}}^{P^\perp} \mathcal{E}_{\mathcal{U}}(X) = \mathcal{E}_{\overline{V}}^{P^\perp}(X) = f_V^{P^\perp}(Y_X^{P^\perp})[1].$$

We next compute $\mathcal{E}_{\mathcal{U}\amalg\mathcal{V}}(X)$. First note that $\mathcal{E}_{\mathcal{V}}(P[1]) = f_V(Y_P)[1]$, where $P \rightarrow Y_P$ is a minimal left ${}^\perp(\tau V)$ -approximation, and that $\mathcal{E}_{\mathcal{V}}(X) = f_V(Y_X)[1]$, where there is an exact sequence

$$(45) \quad Y_X \rightarrow V_X \rightarrow X \rightarrow 0$$

where the first map is a minimal left add V -approximation, and the second map is a minimal right add V -approximation. Then

$$\mathcal{E}_{\mathcal{U}\amalg\mathcal{V}}(X) = \mathcal{E}_{P[1]\amalg V} = \mathcal{E}_{\mathcal{E}_{\mathcal{V}}(P[1])}^{J(V)} \mathcal{E}_V(X) = f_{f_V(Y_P)}^{J(V)} f_V(Y_X)[1].$$

Hence we need to prove that

$$(46) \quad f_V^{P^\perp}(Y_X^{P^\perp}) \simeq f_{f_V(Y_P)}^{J(V)} f_V(Y_X).$$

We first make the following observation.

Lemma 7.1. *Let P be a projective module in $\text{mod } \Lambda$. Then f_P is a right exact functor from $\text{mod } \Lambda$ to P^\perp , and f_P sends projective modules to projective modules in P^\perp .*

Proof. Let e be an idempotent such that $P \simeq \Lambda e$. We first note that $t_P(M) = \Lambda eM$, so $f_P(M) \simeq M/\Lambda eM \simeq \Lambda/\Lambda e\Lambda \otimes_{\Lambda} M$. It follows that f_P is right exact.

Moreover, since $\Lambda/\Lambda e\Lambda \otimes_{\Lambda} \Lambda \simeq \Lambda/\Lambda e\Lambda$, and the tensor-functor is additive, we have that the indecomposable projective $\Lambda/\Lambda e\Lambda$ -modules are exactly $f_P(T)$ for T indecomposable projective in $\text{mod } \Lambda$ with T not a summand in P . \square

We proceed to prove (46). The main steps in the proof are as follows.

- Claim 7.2.** (a) We have $f_P(Y_X) \simeq Y_X^{P^\perp}$.
 (b) The composition $Y_X \xrightarrow{\alpha} f_P(Y_X) \xrightarrow{\beta} f_{\bar{V}}^{P^\perp} f_P(Y_X)$ is a left $J(V) \cap P^\perp$ -approximation.
 (c) The composition $Y_X \xrightarrow{\gamma} f_V(Y_X) \xrightarrow{\phi} f_{f_V(Y_P)}^{J(V)} f_V(Y_X)$ is a left $J(V) \cap P^\perp$ -approximation.
 (d) We have $f_{\bar{V}}^{P^\perp} f_P(Y_X) \simeq f_{f_V(Y_P)}^{J(V)} f_V(Y_X)$.
 (e) We have $f_V^{P^\perp}(Y_X^{P^\perp}) \simeq f_{f_V(Y_P)}^{J(V)} f_V(Y_X)$.

Proof. (a): Let $\mathcal{T} = {}^\perp(\tau V)$ and let $\mathcal{T}' = {}^\perp(\tau_{P^\perp} \bar{V}) \cap P^\perp$. Then we have $f_P \mathcal{T} = \mathcal{T}'$ by Lemma 5.6. We have that $Y_X^{P^\perp}$ is in $P^\perp \cap {}^\perp(\tau V) = P^\perp \cap {}^\perp(\tau \bar{V})$.

Note that since $\text{Hom}(P, X) = 0 = \text{Hom}(P, V_X)$, we have $f_P(X) = X$ and $f_P(V_X) = V_X$. Hence, applying f_P to the right exact sequence (45) we obtain the right exact sequence

$$(47) \quad f_P(Y_X) \xrightarrow{f_P(a)} V_X \rightarrow X \rightarrow 0.$$

We claim that $f_P(a)$ is a minimal left $\bar{V} = V$ -approximation in P^\perp . Let $b': f_P(Y_X) \rightarrow f_P(V') = V'$ be a map with $V' \in \text{add } V \subseteq P^\perp$. By Lemma 5.2, there is a map $b: Y_X \rightarrow V'$ such that $b' = f_P(b)$. Since a is left add V -approximation, there is a map $c: V_X \rightarrow V'$ such that $b = ca$. So $f_P(c)f_P(a) = f_P(b)$. We have that $f_P(a)$ is minimal, since otherwise we would have X in $\text{add } V$.

Using now [4, Proposition 5.6], we have that $f_P(Y_X) \simeq Y_X^{P^\perp}$, and this concludes the proof of (a).

(b): Let Z be in $J(V) \cap P^\perp$ and let $g: Y_X \rightarrow Z$ be a map. Since α is a left P^\perp -approximation and Z is in P^\perp , there is a map $l: f_P(Y_X) \rightarrow Z$, such that $l\alpha = g$. Since β is a left $P^\perp \cap V^\perp$ -approximation and Z is in $P^\perp \cap V^\perp$, there is a map $r: f_{\bar{V}}^{P^\perp} f_P(Y_X) \rightarrow Z$, such that $l = r\beta$. Hence $g = l\alpha = r\beta\alpha$. Since Y_X is in ${}^\perp(\tau V)$, we have that also $f_{\bar{V}}^{P^\perp} f_P(Y_X)$ is in ${}^\perp(\tau V)$, and hence $f_{\bar{V}}^{P^\perp} f_P(Y_X)$ is in $P^\perp \cap V^\perp \cap {}^\perp(\tau V) = J(V) \cap P^\perp$. This proves the claim that $\beta\alpha$ is a left $J(V) \cap P^\perp$ -approximation.

(c): Let Z be in $J(V) \cap P^\perp$ and let $g: Y_X \rightarrow Z$ be a map. Since γ is a left V^\perp -approximation, and Z is in V^\perp , there is a map $s: f_V(Y_X) \rightarrow Z$ such that $g = s\gamma$. Note that we have that $f_V(Y_P)$ is in $\mathcal{P}(J(V))$, and so

$$(f_V(Y_P))^\perp \cap J(V) = J_{J(V)}(f_V(Y_P)[1]) = J_{J(V)}(\mathcal{E}_V(P[1])) = J(V \amalg P[1]) = J(V) \cap P^\perp.$$

The map ϕ is a left $(f_V(Y_P))^\perp \cap J(V) = J(V) \cap P^\perp$ -approximation. Hence, there is a map $t: f_{f_V(Y_P)}^{J(V)} f_V(Y_X) \rightarrow Z$ such that $s = t\phi$, and therefore $g = s\gamma = t\phi\gamma$. This proves the claim that $\phi\gamma$ is a left $J(V) \cap P^\perp$ -approximation.

(d): Since both $\beta\alpha$ and $\phi\gamma$ are epimorphisms, they are both minimal left $J(V) \cap P^\perp$ -approximations, and the claim follows.

(e): This follows directly from combining (a) and (d). \square

Equation (25) in this case now follows from (46).

Case III (c): We assume that $X = R[1]$, where R is an indecomposable projective module in $\text{mod } \Lambda$ and $\text{Hom}(R, V) = 0$.

We then have $\mathcal{E}_{\mathcal{U}}(X) = f_P(R)[1]$, which is in $\mathcal{P}(P^\perp)[1]$ and we have that

$$\mathcal{E}_{\mathcal{V}}^{J(\mathcal{U})} \mathcal{E}_{\mathcal{U}}(X) = \mathcal{E}_{\mathcal{V}}^{P^\perp}(f_P(R)[1]).$$

Note that $V = \bar{V}$ is τ -rigid in $J(\mathcal{U}) = P^\perp$. Therefore $\mathcal{E}_{\mathcal{V}}^{P^\perp}(f_P(R)[1]) = f_V^{P^\perp}(Y_0)[1]$, where $f_P(R) \rightarrow Y_0$ is a minimal left ${}^\perp(\tau_{P^\perp} V) \cap P^\perp$ -approximation.

We have that $\mathcal{E}_{\mathcal{V}}(P[1]) = f_V(Y_P)[1]$, where $P \rightarrow Y_P$ is a minimal left ${}^\perp(\tau V)$ -approximation and similarly $\mathcal{E}_{\mathcal{V}}(X) = f_V(Y_R)[1]$, where $R \rightarrow Y_R$ is a minimal left ${}^\perp(\tau V)$ -approximation. It follows that

$$(48) \quad \mathcal{E}_{\mathcal{U} \sqcup \mathcal{V}}(X) = \mathcal{E}_{P[1] \sqcup V}(X) = \mathcal{E}_{\mathcal{E}_{\mathcal{V}}(P[1])}^{J(V)} \mathcal{E}_{\mathcal{V}}(X) = \mathcal{E}_{f_V(Y_P)[1]}^{J(V)} \mathcal{E}_{\mathcal{V}}(X) = f_{f_V(Y_P)}^{J(V)} f_V(Y_R)[1].$$

So it will be sufficient to prove that

$$(49) \quad f_V^{P^\perp} Y_0 \simeq f_{f_V(Y_P)}^{J(V)} f_V(Y_R).$$

The main steps in the proof of this are as follows.

Claim 7.3. (a) *We have that Y_0 is a direct summand of $f_P(Y_R)$.*

(b) *We have that $f_V^{P^\perp}(Y_0)$ is a direct summand of $f_V^{P^\perp} f_P(Y_R)$.*

(c) *The composition*

$$Y_R \xrightarrow{\alpha} f_P(Y_R) \xrightarrow{\beta} f_V^{P^\perp} f_P(Y_R)$$

is a minimal left $J(V) \cap P^\perp$ -approximation.

(d) *The composition*

$$Y_R \xrightarrow{\gamma} f_V(Y_R) \xrightarrow{\phi} f_{f_V(Y_P)}^{J(V)} f_V(Y_R)$$

is a minimal left $J(V) \cap P^\perp$ -approximation.

(e) *We have $f_V^{P^\perp} f_P(Y_R) \simeq f_{f_V(Y_P)}^{J(V)} f_V(Y_R)$.*

(f) *We have $f_V^{P^\perp} Y_0 \simeq f_{f_V(Y_P)}^{J(P)} f_V(Y_R)$.*

Proof. (a): Let $\mathcal{T} = {}^\perp(\tau V)$ and $\mathcal{T}' = {}^\perp(\tau_{P^\perp} V) \cap P^\perp$. Note that the map $f_P(R) \rightarrow Y_0$ is a minimal left \mathcal{T}' -approximation so, for the claim, it is sufficient to prove that $f_P(R) \rightarrow f_P(Y_R)$ is a left \mathcal{T}' -approximation. By Lemma 5.6, we have that $f_P(\mathcal{T}) = \mathcal{T}'$, so by Lemma 5.3 we have that $f_P(R) \rightarrow f_P(Y_R)$ is a left \mathcal{T}' -approximation (noting that $\text{Hom}(P, \tau R) = 0$ as R is projective), giving the claim.

(b): This follows directly from (a).

(c): Note first that since Y_R is in ${}^\perp(\tau V)$, also the factor module $f_V^{P^\perp} f_P(Y_R)$ is in ${}^\perp(\tau V)$. This module also lies in $P^\perp \cap V^\perp$ by the definition of $f_V^{P^\perp}$, so it lies in $J(V) \cap P^\perp$.

Let Z be in $J(V) \cap P^\perp$, and consider a map $g: Y_R \rightarrow Z$. Since α is a left P^\perp -approximation, and Z is in P^\perp , so there is a map $l: f_P(Y_R) \rightarrow Z$, such that $l\alpha = g$. Since β is a left $P^\perp \cap V^\perp$ -approximation, and Z is in $P^\perp \cap V^\perp$, there is a map $r: f_V^{P^\perp} f_P(Y_R) \rightarrow Z$, such that $l = r\beta$. Hence we have $g = l\alpha = r\beta\alpha$, and this proves that $\beta\alpha$ is left $J(V) \cap P^\perp$ approximation. Since this composition is an epimorphism, it must also be left minimal, giving the claim.

(d): First note that $f_V(Y_P)$ is in $\mathcal{P}(J(V))$, and hence $(f_V(Y_P))^\perp \cap J(V) = J_{J(V)}(f_V(Y_P)[1]) = J_{J(V)}(\mathcal{E}_V(P[1])) = J(V \amalg P[1]) = J(V) \cap P^\perp$. Let $g: Y_R \rightarrow Z$ be a map, with Z in $J(V) \cap P^\perp$. Since γ is a V^\perp -approximation, and Z is in V^\perp , there is a map $t: f_V(Y_R) \rightarrow Z$ such that $g = t\gamma$. Since ϕ is a left $(f_V(Y_P))^\perp \cap J(V) = J(V) \cap P^\perp$ -approximation, there is a map $u: f_{f_V(Y_P)}^{J(V)} f_V(Y_R) \rightarrow Z$, such that $t = u\phi$. It follows that $g = t\gamma = u\phi\gamma$. This proves that $\phi\gamma$ is left $J(V) \cap P^\perp$ approximation. Since this composition is an epimorphism, it must also be left minimal, giving the claim.

(e): This follows directly from (c) and (d).

(f): Note that $f_V^{P^\perp}(Y_0)$ is indecomposable by [4, Proposition 5.6] (see the definition of Y_0 above). It is a direct summand of $f_{f_V(Y_P)}^{J(V)} f_V(Y_R)$ which is indecomposable by (48) and [4, Proposition 5.6]. The claim follows. \square

We have proved that (49) holds, and (25) in this case now follows.

8. PROOF OF THEOREM 5.9, CASE IV

We assume that $\mathcal{U} = P[1]$ and $\mathcal{V} = Q[1]$, where P, Q lie in $\mathcal{P}(\Lambda)$.

We set $\overline{V} = \overline{Q}[1]$. Then $\overline{Q} = f_P Q$ lies in $\mathcal{P}(J(\mathcal{U})) = \mathcal{P}(P^\perp)$.

We need in this case to consider two possible cases for X :

- (a) X is τ -rigid, X lies in $\text{ind}(\Lambda)$, and $\text{Hom}(P \amalg Q, X) = 0$.
- (b) X lies in $\text{ind } \mathcal{P}(\Lambda)[1]$.

Case IV (a): We assume that X is an indecomposable τ -rigid module with $\text{Hom}(P \amalg Q, X) = 0$.

We have $\mathcal{E}_{\mathcal{U}}(X) = \mathcal{E}_{P[1]}(X) = X$, and then $\mathcal{E}_{\mathcal{V}}^{J(\mathcal{U})} \mathcal{E}_{\mathcal{U}}(X) = \mathcal{E}_{\overline{Q}[1]}^{P^\perp} X = X$, since $\text{Hom}(\overline{Q}, X) = 0$ by Theorem 3.7.

We also have $\mathcal{E}_{\mathcal{U} \amalg \mathcal{V}}(X) = \mathcal{E}_{P[1] \amalg Q[1]}(X) = X$, so the claim that equation (25) holds follows also in this case.

Case IV (b): We assume that X is of the form $R[1]$, where R is an indecomposable module in $\mathcal{P}(\Lambda)$.

We then have that

$$\mathcal{E}_{\mathcal{V}}^{J(\mathcal{U})} \mathcal{E}_{\mathcal{U}}(X) = \mathcal{E}_{\overline{Q}[1]}^{P^\perp} \mathcal{E}_{P[1]}(X) = \mathcal{E}_{\overline{Q}[1]}^{P^\perp} (f_P(R)[1]) = f_Q^{P^\perp} f_P(R)[1].$$

On the other hand, we have $\mathcal{E}_{\mathcal{U} \amalg \mathcal{V}}(X) = \mathcal{E}_{P[1] \amalg Q[1]}(X) = f_{P \amalg Q}(R)[1]$.

So, it is sufficient to prove that $f_Q^{P^\perp} f_P(R) \simeq f_{P \amalg Q}(R)$. The main steps in the proof are as follows.

Claim 8.1. (a) We have $P^\perp \cap \overline{Q}^\perp = P^\perp \cap Q^\perp$.

(b) The composition $R \xrightarrow{\alpha} f_P(R) \xrightarrow{\beta} f_Q^{P^\perp} f_P(R)$ is a $(P \amalg Q)^\perp$ -approximation.

(c) We have $f_Q^{P^\perp} f_P(R) \simeq f_{P \amalg Q}(R)$.

Proof. (a): Note first that, for any module M , $\text{Hom}(P, M) = 0$ implies $\text{Hom}(\text{Gen } P, M) = 0$. Suppose M lies in $P^\perp \cap \overline{Q}^\perp$. We apply $\text{Hom}(_, M)$ to the canonical sequence

$$0 \rightarrow t_P(Q) \rightarrow Q \rightarrow \overline{Q} \rightarrow 0$$

for Q . We have $\text{Hom}(\overline{Q}, M) = 0$ and also $\text{Hom}(t_P(Q), M) = 0$, since $t_P(Q)$ is in $\text{Gen } P$. Hence $\text{Hom}(Q, M) = 0$ and we have $P^\perp \cap \overline{Q}^\perp \subseteq P^\perp \cap Q^\perp$. The reverse inclusion follows immediately from the fact that \overline{Q} is a factor of Q .

(b): By (a), we have that $f_Q^{P^\perp}(f_P(R))$ is in $P^\perp \cap \overline{Q}^\perp = P^\perp \cap Q^\perp$. Consider a map $g: R \rightarrow Z$ with Z in $(P \amalg Q)^\perp$. Since α is a left P^\perp -approximation and Z is in P^\perp , there is a map $t: f_P(R) \rightarrow Z$ such that $g = t\alpha$. Since $f_P(R) \rightarrow f_Q^{P^\perp} f_P(R)$ is a left $P^\perp \cap \overline{Q}^\perp = (P \amalg Q)^\perp$ -approximation and Z is in $P^\perp \cap Q^\perp$, there is a map $u: f_Q^{P^\perp} f_P(R) \rightarrow Z$ such that $t = u\beta$. We then have $g = t\alpha = u\beta\alpha$. This proves the claim.

(c): This follows directly from (b), noting that both $f_Q^{P^\perp} f_P(R)$ and $f_{P \amalg Q}(R)$ are indecomposable (since they are factors of the indecomposable projective module R). \square

This finishes the proof that (25) holds in this case.

9. END OF THE PROOF OF THEOREM 5.9: MIXED CASE

We have now proved that 25 holds for all of the cases I-IV. It remains to deal with the mixed cases, where we have support τ -rigid objects $\mathcal{U} = U \amalg P[1]$ and $\mathcal{V} = V \amalg Q[1]$ in $C(\Lambda)$, with no common direct summands, but where we allow indecomposable direct summands of \mathcal{U} and \mathcal{V} to lie both in $\text{mod } \Lambda$ and in $\mathcal{P}(\Lambda)[1]$.

Let us summarize the formulas we need to proceed. By Cases I-IV, we have that the formulas

$$(50) \quad \mathcal{E}_{\mathcal{E}_{\mathcal{U}}(\mathcal{V})}^{J(\mathcal{U})} \mathcal{E}_{\mathcal{U}} = \mathcal{E}_{\mathcal{U} \amalg \mathcal{V}} = \mathcal{E}_{\mathcal{E}_{\mathcal{V}}(\mathcal{U})}^{J(\mathcal{V})} \mathcal{E}_{\mathcal{V}}$$

hold when we have both of the following:

- $U = 0$ or $P = 0$, and
- $V = 0$ or $Q = 0$.

Note that a particular case is when $\mathcal{U} = U$ and $\mathcal{V} = Q[1]$, where U lies in $\text{mod } \Lambda$ and Q lies in $\mathcal{P}(\Lambda)$. We therefore have

$$(51) \quad \mathcal{E}_{U \amalg Q[1]} = \mathcal{E}_{\mathcal{E}_U(P[1])}^{J(U)} \mathcal{E}_U.$$

Recall also from Section 4 that we have

$$(52) \quad J_{J(\mathcal{U})}(\mathcal{E}_{\mathcal{U}}(\mathcal{V})) = J(\mathcal{U} \amalg \mathcal{V}),$$

for any pair of support τ -rigid objects \mathcal{U}, \mathcal{V} in $C(\Lambda)$.

Case A: We first discuss the case with $P = 0$, that is $\mathcal{U} = U \neq 0$, while $\mathcal{V} = V \amalg Q[1]$ is arbitrary. We work by induction on $n = r(\bmod \Lambda)$. We then have

$$\begin{aligned}
(53) \quad \mathcal{E}_{\mathcal{U} \amalg \mathcal{V}} &= \mathcal{E}_{U \amalg V \amalg Q[1]} \\
(54) \quad &= \mathcal{E}_{\mathcal{E}_{U \amalg V}(Q[1])}^{J(U \amalg V)} \mathcal{E}_{U \amalg V} \\
(55) \quad &= \mathcal{E}_{\mathcal{E}_{U \amalg V}(Q[1])}^{J(U \amalg V)} \mathcal{E}_{\mathcal{E}_U(V)}^{J(U)} \mathcal{E}_U \\
(56) \quad &= \mathcal{E}_{\mathcal{E}_{U \amalg V}(Q[1])}^{J_{J(U)}(\mathcal{E}_U(V))} \mathcal{E}_{\mathcal{E}_U(V)}^{J(U)} \mathcal{E}_U \\
(57) \quad &= \mathcal{E}_{\mathcal{E}_{\mathcal{E}_U(V) \amalg \mathcal{E}_U(Q[1])}}^{J(U)} \mathcal{E}_U \\
&= \mathcal{E}_{\mathcal{E}_U(V \amalg Q[1])}^{J(U)} \mathcal{E}_U \\
&= \mathcal{E}_{\mathcal{E}_U(\mathcal{V})}^{J(\mathcal{U})} \mathcal{E}_U
\end{aligned}$$

where equation (53) follows from (51), while (54) and (56) follows from (50) and (55) from (52). Furthermore, equation (57) follows from the induction assumption, since $r(J(U)) < n$. This concludes the proof of the case with $P = 0$, i.e. $\mathcal{U} = U$.

Case B: We next discuss the case with $U = 0$, that is $\mathcal{U} = P[1] \neq 0$, while $\mathcal{V} = V \amalg Q[1]$ is arbitrary. We also assume $V \neq 0$, note that we have already dealt with the case $V = 0$ (this is Case IV). We then have:

$$\begin{aligned}
(58) \quad \mathcal{E}_{\mathcal{E}_U(\mathcal{V})}^{J(\mathcal{U})} \mathcal{E}_U &= \mathcal{E}_{\mathcal{E}_{P[1]}(V \amalg Q[1])}^{J(P[1])} \mathcal{E}_{P[1]} \\
&= \mathcal{E}_{\mathcal{E}_{P[1]}(V) \amalg \mathcal{E}_{P[1]}(Q[1])}^{J(P[1])} \mathcal{E}_{P[1]} \\
(59) \quad &= \mathcal{E}_{\mathcal{E}_{P[1]}(V)}^{J_{J(P[1])}(\mathcal{E}_{P[1]}(V))} \mathcal{E}_{\mathcal{E}_{P[1]}(V)}^{J(P[1])} \mathcal{E}_{P[1]} \\
(60) \quad &= \mathcal{E}_{\mathcal{E}_{P[1]} \amalg V(Q[1])}^{J_{J(P[1])}(\mathcal{E}_{P[1]}(V))} \mathcal{E}_{\mathcal{E}_{P[1]}(V)}^{J(P[1])} \mathcal{E}_{P[1]} \\
(61) \quad &= \mathcal{E}_{\mathcal{E}_{P[1]} \amalg V(Q[1])}^{J(P[1] \amalg V)} \mathcal{E}_{P[1]}^{J(P[1])} \mathcal{E}_{P[1]} \\
(62) \quad &= \mathcal{E}_{\mathcal{E}_{P[1]} \amalg V(Q[1])}^{J(P[1] \amalg V)} \mathcal{E}_{\mathcal{E}_V(P[1])}^{J(V)} \mathcal{E}_V \\
(63) \quad &= \mathcal{E}_{\mathcal{E}_V(P[1])}^{J(P[1] \amalg V)} \mathcal{E}_{\mathcal{E}_V(Q[1])}^{J(V)} \mathcal{E}_V \\
(64) \quad &= \mathcal{E}_{\mathcal{E}_V(P[1])}^{J_{J(V)}(\mathcal{E}_V(P[1]))} \mathcal{E}_{\mathcal{E}_V(Q[1])}^{J(V)} \mathcal{E}_V
\end{aligned}$$

$$(65) \quad = \mathcal{E}_{\mathcal{E}_V(P[1] \amalg Q[1])}^{J(V)} \mathcal{E}_V$$

$$(66) \quad = \mathcal{E}_{V \amalg P[1] \amalg Q[1]}$$

$$= \mathcal{E}_{\mathcal{U} \amalg \mathcal{V}}$$

where for (58), we use that (25) holds in $J(P[1])$ by induction. For (65), we use that the (25) holds in $J(V)$ by induction. For (59) and (61) we apply (50), for (62), (63) and (66) we apply (51), while for (60) and (64) we apply (52).

The general case: We now discuss the general case with $\mathcal{U} = U \amalg P[1]$ and $\mathcal{V} = V \amalg Q[1]$.

We then have

$$(67) \quad \mathcal{E}_{\mathcal{E}_U(\mathcal{V})}^{J(\mathcal{U})} \mathcal{E}_U = \mathcal{E}_{\mathcal{E}_{U \amalg P[1]}(V \amalg Q[1])}^{J(U \amalg P[1])} \mathcal{E}_{U \amalg P[1]}$$

$$(68) \quad = \mathcal{E}_{\mathcal{E}_{U \amalg P[1]}(V \amalg Q[1])}^{J(U \amalg P[1])} \mathcal{E}_{\mathcal{E}_U(P[1])}^{J(U)} \mathcal{E}_U$$

$$(69) \quad = \mathcal{E}_{\mathcal{E}_{\mathcal{E}_U(P[1])} \mathcal{E}_U(V \amalg Q[1])}^{J(U \amalg P[1])} \mathcal{E}_{\mathcal{E}_U(P[1])}^{J(U)} \mathcal{E}_U$$

$$(70) \quad = \mathcal{E}_{\mathcal{E}_{\mathcal{E}_U(P[1])} \mathcal{E}_U(V \amalg Q[1])}^{J_{J(U)}(\mathcal{E}_U(P[1]))} \mathcal{E}_{\mathcal{E}_U(P[1])}^{J(U)} \mathcal{E}_U$$

$$(71) \quad = \mathcal{E}_{\mathcal{E}_U(P[1]) \amalg \mathcal{E}_U(V \amalg Q[1])}^{J(U)} \mathcal{E}_U$$

$$= \mathcal{E}_{\mathcal{E}_U(P[1] \amalg V \amalg Q[1])}^{J(U)} \mathcal{E}_U$$

$$= \mathcal{E}_{\mathcal{E}_U(V \amalg P[1] \amalg Q[1])}^{J(U)} \mathcal{E}_U$$

$$(71) \quad = \mathcal{E}_{U \amalg P[1] \amalg V \amalg Q[1]}$$

$$= \mathcal{E}_{\mathcal{U} \amalg \mathcal{V}}$$

where (67) and (68) hold by (51), while (69) holds by (52). For (70) we note that $\mathcal{E}_U(P[1])$ is in $\mathcal{P}(J(U))[1]$, so that we are in the situation of Case B in $J(U)$. For equation (71), we apply Case A.

This concludes the proof of Theorem 5.9. \square

10. IRREDUCIBLE MORPHISMS IN \mathfrak{B}_Λ

In this section we prove the following Theorem.

Theorem 10.1. *Let Λ be a τ -tilting finite algebra, and let $\mathcal{W}' \subseteq \mathcal{W}$ be wide subcategories of $\text{mod } \Lambda$, where $r(\mathcal{W}) - r(\mathcal{W}') = 1$ (i.e \mathcal{W}' is of corank 1 in \mathcal{W}). Then exactly one of the following occurs:*

- (a) *There is exactly one morphism in \mathfrak{B}_Λ from \mathcal{W} to \mathcal{W}' and $\mathcal{W}' = J_{\mathcal{W}}(U)$, where U is an indecomposable τ -rigid module which is non-projective in \mathcal{W} .*
- (b) *There are exactly two morphisms in \mathfrak{B}_Λ from \mathcal{W} to \mathcal{W}' and $\mathcal{W}' = J_{\mathcal{W}}(P) = J_{\mathcal{W}}(P[1])$, where P is an indecomposable module which is projective in \mathcal{W} .*

The main step in the proof is to show that if U and V are indecomposable τ -rigid Λ -modules satisfying $J(U) = J(V)$, then $U = V$.

Definition 10.2. A morphism g in \mathfrak{B}_Λ is said to be *irreducible* if, whenever g is expressed as a composition $g_1 \circ g_2$, we have that either g_1 or g_2 is an identity map.

Lemma 10.3. *Let \mathcal{W} be a wide subcategory of $\text{mod } \Lambda$ and let \mathcal{V} be a support τ -rigid object in $C(\mathcal{W})$. Then the following are equivalent.*

- (a) *The morphism $g = g_{\mathcal{V}}^{\mathcal{W}}: \mathcal{W} \rightarrow J_{\mathcal{W}}(\mathcal{V})$ is irreducible.*
- (b) *The object \mathcal{V} is indecomposable.*
- (c) *The subcategory $J_{\mathcal{W}}(\mathcal{V})$ is of corank 1 in \mathcal{W} .*

Proof. Suppose first that \mathcal{V} is indecomposable, and that $g = g_1 \circ g_2$ for maps g_1 and g_2 in \mathfrak{B} . Then we have $g_1 = g_{\mathcal{U}_1}^{\mathcal{W}_1}$ and $g_2 = g_{\mathcal{U}_2}^{\mathcal{W}_2}$ where \mathcal{U}_1 is a support τ -rigid object in $C(\mathcal{W}_1)$, \mathcal{U}_2 is a support τ -rigid object in $C(\mathcal{W}_2)$ and $J_{\mathcal{W}_2}(\mathcal{U}_2) = \mathcal{W}_1$. The composition is:

$$g_{\mathcal{V}}^{\mathcal{W}} = g_{\mathcal{U}_1}^{\mathcal{W}_1} \circ g_{\mathcal{U}_2}^{\mathcal{W}_2} = g_{\mathcal{F}_{\mathcal{U}_2}(\mathcal{U}_1) \amalg \mathcal{U}_2}^{\mathcal{W}_2}.$$

Hence $\mathcal{W}_2 = \mathcal{W}$ and $\mathcal{V} = \mathcal{F}_{\mathcal{U}_2}(\mathcal{U}_1) \amalg \mathcal{U}_2$. Since \mathcal{V} is indecomposable, we have $\mathcal{U}_2 = 0$ or $\mathcal{F}_{\mathcal{U}_2}(\mathcal{U}_1) = 0$. So $\mathcal{U}_1 = 0$ or $\mathcal{U}_2 = 0$, and g_1 or g_2 is an identity map. It follows that g is irreducible. This proves that (b) implies (a).

If \mathcal{V} is decomposable, it can be written in the form $\mathcal{V} = \mathcal{V}_1 \amalg \mathcal{V}_2$ where \mathcal{V}_1 and \mathcal{V}_2 are non-zero support τ -rigid objects in $C(\mathcal{W})$. Then we have:

$$g = g_{\mathcal{V}_1 \amalg \mathcal{V}_2}^{\mathcal{W}} = g_{\mathcal{E}_{\mathcal{V}_2}(\mathcal{V}_1)}^{\mathcal{W}_1} \circ g_{\mathcal{V}_2}^{\mathcal{W}_2},$$

where $\mathcal{W}_2 = \mathcal{W}$ and $\mathcal{W}_1 = J_{\mathcal{W}_2}(\mathcal{V}_2)$. Since \mathcal{V}_1 and $\mathcal{E}_{\mathcal{V}_2}(\mathcal{V}_1)$ are non-zero, $g_{\mathcal{E}_{\mathcal{V}_2}(\mathcal{V}_1)}^{\mathcal{W}_1}$ and $g_{\mathcal{V}_2}^{\mathcal{W}_2}$ are not identity maps, so g is not irreducible. This proves that (a) implies (b).

We have that (b) and (c) are equivalent by Proposition 4.1. \square

Recall that for any τ -rigid Λ -module U there is a unique basic module B_U , known as the *Bongartz complement* of U , such that $\text{add}(U \amalg B_U)$ is a τ -tilting module and $\text{add}(U \amalg B_U) = \mathcal{P}(^{\perp}\tau U)$. We also recall that a Λ module M is said to be *Gen-minimal* if, whenever $M = M' \oplus M''$, we have $M'' \notin \text{Gen } M'$ (see e.g. [2, VI.6]). We recall the following:

Lemma 10.4. [8, Lemma 2.8] *Let Λ be an algebra, and let \mathcal{T} be a finitely generated torsion class in $\text{mod } \Lambda$. Then \mathcal{T} has a unique Gen-minimal generator, \mathcal{T}_{\min} , consisting of the direct sum of the indecomposable split projective objects in \mathcal{T} .*

If T is a support τ -tilting module, then we denote the unique Gen-minimal generator of $\text{Gen } T$ by T_s . Note that T is an additive generator for $\mathcal{P}(\mathcal{T})$ by [1, Thm. 2.7], so T_s is a direct summand of T , and we write T_{ns} for a complement, the direct sum of the non-split projective objects in $\text{Gen } T$.

If Z is a minimal direct summand of T such that $\text{Gen } Z = \text{Gen } T$ then $T_s \in \text{Gen } Z$, so T_s is a direct summand of Z since it is split projective. Since $\text{Gen } T_s = \text{Gen } T$, we must have $T_s = Z$. In the light of this discussion, we also recall the following:

Theorem 10.5. [6, Thm. 3.34] *Let Λ be a τ -tilting finite algebra. Then there is a bijection between the set of τ -tilting pairs in $\text{mod } \Lambda$ and the set of wide subcategories of $\text{mod } \Lambda$ given by mapping a τ -tilting pair (T, P) to $\mathcal{W}(T, P) = J(T_{\text{ns}}) \cap P^{\perp}$.*

Lemma 10.6. *Let Λ be a τ -tilting finite algebra. Let U be a non-projective τ -rigid module in $\text{mod } \Lambda$. Let B_U be the Bongartz complement of U , and let $T_U = U \amalg B_U$. Then $(T_U)_s = B_U$ and $(T_U)_{ns} = U$.*

Proof. By the definition of Bongartz complement, we have that $\text{add}(T_U) = \mathcal{P}({}^\perp(\tau U))$. By [4, Lemma 4.12], the indecomposable direct summands of B_U are split projective in ${}^\perp(\tau U)$. Suppose that U was also split projective in ${}^\perp(\tau U)$. Then we would have $(T_U)_{ns} = 0$ and therefore $\mathbf{W}(T_U, 0) = \text{mod } \Lambda$ in Theorem 10.5. But $\mathbf{W}(P, 0) = \text{mod } \Lambda$, where P is an additive generator for $\mathcal{P}(\Lambda)$, so $T_U = P$ by Theorem 10.5, and U is projective, giving a contradiction. Hence U is not split projective in ${}^\perp(\tau U)$ and we are done. \square

Proposition 10.7. *Let Λ be a τ -tilting finite algebra. Let U and V be indecomposable τ -rigid Λ -modules and suppose that $J(U) = J(V)$. Then $U = V$.*

Proof. Let B_U (respectively, B_V) be the Bongartz complement of U (respectively, V), and set $T_U = U \amalg B_U$ and $T_V = V \amalg B_V$. Then, since T_U and T_V are τ -tilting modules, we have that $(T_U, 0)$ and $(T_V, 0)$ are τ -tilting pairs. We have $\mathbf{W}(T_U, 0) = J((T_U)_{ns}) = J(U)$ by Lemma 10.6, and similarly $\mathbf{W}(T_V, 0) = J(V)$. So, by Theorem 10.5, $U = V$. \square

We now finish the proof of Theorem 10.1.

Proof of Theorem 10.1: By Lemma 10.3 and Proposition 4.2, we have $\mathbf{W}' = J_{\mathbf{W}}(\mathcal{U})$ where \mathcal{U} is either an indecomposable τ -rigid module or $\mathcal{U} = P[1]$ for an indecomposable module P which is projective in \mathbf{W} . The result now follows from Proposition 10.7 and the fact that $J_{\mathbf{W}}(P) = J_{\mathbf{W}}(P[1])$. \square

11. MORPHISMS IN \mathfrak{B}_Λ AND SIGNED τ -EXCEPTIONAL SEQUENCES

The notion of signed τ -exceptional sequence was introduced in [4]. Such sequences can be interpreted as factorizations of morphisms in the category \mathbf{W}_Λ . Our aim in this section is to make a precise version of this statement.

Recall from [4] that an object $M \amalg P[1]$ in $C(\Lambda)$ is said to be support τ -rigid if M is a τ -rigid module in $\text{mod } \Lambda$, P lies in $\mathcal{P}(\Lambda)$ and $\text{Hom}(P, M) = 0$. Furthermore, a sequence

$$(72) \quad \mathcal{S} = (\mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_t)$$

of indecomposable objects in $C(\Lambda)$ is said to be a *signed τ -exceptional sequence* if \mathcal{U}_t is support τ -rigid in $C(\Lambda)$ and the subsequence $(\mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_{t-1})$ is a signed τ -exceptional sequence in $J(\mathcal{U}_t)$.

Theorem 11.1. [4, Thm. 5.4] *For each $t \in \{1, \dots, n\}$ there is a bijection φ_t from the set of signed τ -exceptional sequences of length t in $C(\Lambda)$ to the set of ordered support τ -rigid objects of length t in $C(\Lambda)$.*

We have the following, noting that if Λ is τ -tilting finite then every wide subcategory of $\text{mod } \Lambda$ is equivalent to a module category, by Proposition 4.2.

Corollary 11.2. *Suppose that Λ is τ -tilting finite, and let \mathbf{W} be a wide subcategory of $\text{mod } \Lambda$. Then for each $t \in \{1, \dots, n\}$ there is a bijection $\varphi_t^{\mathbf{W}}$ between the set of signed τ -exceptional sequences of length t in \mathbf{W} and the set of ordered support τ -rigid objects of length t in $C(\mathbf{W})$.*

Recall now the following fact from [4, Remark 5.12].

Proposition 11.3. [4] *Assume that Λ is τ -tilting finite. Let \mathcal{W} be a wide subcategory of $\text{mod } \Lambda$. Then the bijection $\varphi_t^{\mathcal{W}}$ in Corollary 11.2 is given by*

$$(\mathcal{U}_1, \dots, \mathcal{U}_t) \mapsto (\mathcal{F}_{\mathcal{U}_t}^{\mathcal{W}_t} \cdots \mathcal{F}_{\mathcal{U}_2}^{\mathcal{W}_2}(\mathcal{U}_1), \mathcal{F}_{\mathcal{U}_t}^{\mathcal{W}_t} \cdots \mathcal{F}_{\mathcal{U}_3}^{\mathcal{W}_3}(\mathcal{U}_2), \dots, \mathcal{U}_t)$$

where $\mathcal{W}_t = \mathcal{W}$ and $\mathcal{W}_i = J_{\mathcal{W}_{i+1}}(\mathcal{U}_{i+1})$ for all i .

To prepare for our main results in this section, we now state and prove the following three lemmas.

Lemma 11.4. *Let \mathcal{W} be a wide subcategory of $\text{mod } \Lambda$, and let $\mathcal{U}_1, \dots, \mathcal{U}_t$ be indecomposable objects in $C(\mathcal{W})$. Then the following are equivalent.*

- (a) *The sequence $(\mathcal{U}_1, \dots, \mathcal{U}_t)$ is a signed τ -exceptional sequence in \mathcal{W} ;*
- (b) *There are wide subcategories $\mathcal{W}_1, \dots, \mathcal{W}_t$ of $\text{mod } \Lambda$ with $\mathcal{W}_t = \mathcal{W}$, and maps $g_{\mathcal{U}_i}^{\mathcal{W}_i}$ for $i = 1, \dots, t$, such that the composition $g_{\mathcal{U}_1}^{\mathcal{W}_1} \cdots g_{\mathcal{U}_t}^{\mathcal{W}_t}$ is well-defined in \mathfrak{B} .*

Proof. We prove that (a) implies (b) by induction on t . If $t = 1$ then \mathcal{U}_1 is support τ -rigid in $C(\mathcal{W})$, so there is a corresponding map $g_{\mathcal{U}_1}^{\mathcal{W}_1}$, taking $\mathcal{W}_1 = \mathcal{W}$, and the result holds for this case. Suppose the result holds for $t - 1$, and let $(\mathcal{U}_1, \dots, \mathcal{U}_t)$ be a signed τ -exceptional sequence in \mathcal{W} of length t . Then $(\mathcal{U}_1, \dots, \mathcal{U}_{t-1})$ is a signed τ -exceptional sequence of length $t - 1$ in $J(\mathcal{U}_t)$. By the induction hypothesis, there are wide subcategories $\mathcal{W}_1, \dots, \mathcal{W}_{t-1}$ of $\text{mod } \Lambda$ with $\mathcal{W}_{t-1} = J_{\mathcal{W}}(\mathcal{U}_t)$, and maps $g_{\mathcal{U}_i}^{\mathcal{W}_i}$ for $i = 1, \dots, t - 1$, such that the composition $g_{\mathcal{U}_1}^{\mathcal{W}_1} \cdots g_{\mathcal{U}_{t-1}}^{\mathcal{W}_{t-1}}$ is well-defined. Since \mathcal{U}_t is support τ -rigid in $C(\mathcal{W})$, there is a map $g_{\mathcal{U}_t}^{\mathcal{W}} : \mathcal{W} \rightarrow J_{\mathcal{W}}(\mathcal{U}_t)$ in \mathfrak{B} . The result follows, taking $\mathcal{W}_t = \mathcal{W}$.

We prove that (b) implies (a) by induction on t . For $t = 1$ the result is clear, so suppose that the result holds for $t - 1$, and let \mathcal{W}_i and $g_{\mathcal{U}_i}^{\mathcal{W}_i}$ be as in (b). Since the composition $g_{\mathcal{U}_1}^{\mathcal{W}_1} \cdots g_{\mathcal{U}_{t-1}}^{\mathcal{W}_{t-1}}$ is well-defined, $(\mathcal{U}_1, \dots, \mathcal{U}_{t-1})$ is a signed τ -exceptional sequence in \mathcal{W}_{t-1} by the induction hypothesis. Since $g_{\mathcal{U}_t}^{\mathcal{W}_t} = g_{\mathcal{U}_t}^{\mathcal{W}}$ is a map, \mathcal{U}_t is support τ -rigid in $C(\mathcal{W})$, and since the composition $g_{\mathcal{U}_1}^{\mathcal{W}_1} \cdots g_{\mathcal{U}_t}^{\mathcal{W}_t}$ is well-defined, we have $J_{\mathcal{W}_t}(\mathcal{U}_t) = \mathcal{W}_{t-1}$, giving (a). \square

Let \mathcal{W} be a wide subcategory of $\text{mod } \Lambda$. For a signed τ -exceptional sequence $\mathcal{U}_1, \dots, \mathcal{U}_t$ in \mathcal{W} , we denote by $\bar{\varphi}_t^{\mathcal{W}}(\mathcal{U}_1, \dots, \mathcal{U}_t)$ the direct sum of the entries in $\varphi_t^{\mathcal{W}}(\mathcal{U}_1, \dots, \mathcal{U}_t)$.

Lemma 11.5. *Let \mathcal{W} be a wide subcategory of $\text{mod } \Lambda$, and suppose that the sequence $(\mathcal{U}_1, \dots, \mathcal{U}_t)$ is a signed τ -exceptional sequence in \mathcal{W} . Set $\mathcal{W}_t = \mathcal{W}$ and $\mathcal{W}_i = J_{\mathcal{W}_{i+1}}(\mathcal{U}_{i+1})$ for all i . Then*

$$g_{\mathcal{U}_1}^{\mathcal{W}_1} \cdots g_{\mathcal{U}_t}^{\mathcal{W}_t} = g_{\bar{\varphi}_t^{\mathcal{W}_t}(\mathcal{U}_1, \dots, \mathcal{U}_t)}^{\mathcal{W}_t}.$$

Proof. We prove the result by induction on t . The result is clear for $t = 1$, so suppose that the result holds for $t - 1$. We have, using Proposition 11.3:

$$\begin{aligned} g_{\mathcal{U}_1}^{\mathcal{W}_1} \cdots g_{\mathcal{U}_t}^{\mathcal{W}_t} &= (g_{\mathcal{U}_1}^{\mathcal{W}_1} \cdots g_{\mathcal{U}_{t-1}}^{\mathcal{W}_{t-1}}) g_{\mathcal{U}_t}^{\mathcal{W}_t} \\ &= g_{\bar{\varphi}_{t-1}^{\mathcal{W}_{t-1}}(\mathcal{U}_1, \dots, \mathcal{U}_{t-1})}^{\mathcal{W}_{t-1}} g_{\mathcal{U}_t}^{\mathcal{W}_t} \end{aligned}$$

$$\begin{aligned}
&= g_{\mathcal{F}_{\mathcal{U}_t}(\overline{\varphi}_{t-1}(\mathcal{U}_1, \dots, \mathcal{U}_{t-1}) \amalg \mathcal{U}_t)}^{\mathbb{W}_t} \\
&= g_{\overline{\varphi}_t}^{\mathbb{W}_t}(\mathcal{U}_1, \dots, \mathcal{U}_t),
\end{aligned}$$

as required. \square

Lemma 11.6. *Let \mathbb{W} be a wide subcategory of $\text{mod } \Lambda$, and let \mathcal{V} be a support τ -rigid object in $C(\mathbb{W})$. If*

$$g_{\mathcal{V}}^{\mathbb{W}} = g_{\mathcal{U}_1}^{\mathbb{W}_1} \cdots g_{\mathcal{U}_t}^{\mathbb{W}_t}$$

is a factorization of $g_{\mathcal{V}}^{\mathbb{W}}$ as a composition of t irreducible maps, then t is the number of indecomposable direct summands of \mathcal{V} .

Proof. By Lemma 10.3, the \mathcal{U}_i are indecomposable objects of $C(\mathbb{W})$. By Lemma 11.5, we have

$$g_{\mathcal{V}}^{\mathbb{W}} = g_{\overline{\varphi}^{\mathbb{W}_t}(\mathcal{U}_1, \dots, \mathcal{U}_t)}^{\mathbb{W}_t},$$

so

$$\mathcal{V} = \overline{\varphi}^{\mathbb{W}_t}(\mathcal{U}_1, \dots, \mathcal{U}_t),$$

and the result follows. \square

We now prove our first main result of this section.

Proposition 11.7. *Let \mathbb{W} be a wide subcategory of $\text{mod } \Lambda$, and let \mathcal{V} be a support τ -rigid object in $C(\mathbb{W})$ with t indecomposable direct summands. Then there is a bijection between:*

- (a) *The set of τ -exceptional sequences $(\mathcal{U}_1, \dots, \mathcal{U}_t)$ in \mathbb{W} such that $\overline{\varphi}_t(\mathcal{U}_1, \dots, \mathcal{U}_t) = \mathcal{V}$;*
- (b) *The set of factorizations of $g_{\mathcal{V}}^{\mathbb{W}}$ into compositions of irreducible maps in \mathfrak{B} .*

Proof. Given a sequence $(\mathcal{U}_1, \dots, \mathcal{U}_t)$ as in (a), set $\mathbb{W}_t = \mathbb{W}$ and $\mathbb{W}_i = J_{\mathbb{W}_{i+1}}(\mathcal{U}_{i+1})$ for all i . Then the composition $g_{\mathcal{U}_1}^{\mathbb{W}_1} \cdots g_{\mathcal{U}_t}^{\mathbb{W}_t}$ is well-defined by Lemma 11.4 and equals $g_{\mathcal{V}}^{\mathbb{W}}$ by the assumption in (a) and Lemma 11.5. By Lemma 10.3, each map $g_{\mathcal{U}_i}^{\mathbb{W}_i}$ is irreducible in \mathfrak{B} .

Any factorization as in (b) must have t factors by Lemma 11.6, so must have form $g_{\mathcal{U}_1}^{\mathbb{W}_1} \cdots g_{\mathcal{U}_t}^{\mathbb{W}_t} = g_{\mathcal{V}}^{\mathbb{W}}$. Given such a factorization, each \mathcal{U}_i is indecomposable by Lemma 10.3 and $\mathcal{V} = \overline{\varphi}_t(\mathcal{U}_1, \dots, \mathcal{U}_t)$ by Lemma 11.5. Furthermore, $(\mathcal{U}_1, \dots, \mathcal{U}_t)$ is a τ -exceptional sequence by Lemma 11.4.

It is clear that these two constructions are inverses of each other, and hence give bijections between the sets in (a) and (b) as required. \square

Recall, from [4], that an *ordered support τ -tilting object* in $C(\Lambda)$ is a sequence

$$(\mathcal{T}_1, \dots, \mathcal{T}_n)$$

of indecomposable support τ -rigid objects in $C(\Lambda)$ with the property that $\amalg_i \mathcal{T}_i$ is a support tilting object.

Theorem 11.8. *Let \mathbb{W} be a wide subcategory of $\text{mod } \Lambda$ and \mathcal{V} a support τ -rigid object in $C(\mathbb{W})$ with t indecomposable direct summands. Then the bijection φ_t induces a bijection between the following sets:*


- (a) *Factorisations of $g_{\mathcal{V}}^{\mathbb{W}}$ into compositions of irreducible maps in \mathfrak{B} ;*

(b) *Ordered decompositions of \mathcal{V} into direct sums of indecomposable objects in $C(W)$.*


Proof. By Proposition 11.7, there is a bijection between the set in (a) and the set of τ -exceptional sequences $(\mathcal{U}_1, \dots, \mathcal{U}_t)$ in W such that $\overline{\varphi}_t^W(\mathcal{U}_1, \dots, \mathcal{U}_t) = \mathcal{V}$. The result now follows from Theorem 11.1. \square

12. EXAMPLE

In this section we consider the following example. Let Q be the quiver



and consider the algebra $\Lambda = kQ/I$ where I is the ideal generated by the path $\beta\alpha$. The AR-quiver of $\text{mod } \Lambda$ is



where the notation indicates which simple modules occur in the radical layers of the module, so $N = \begin{smallmatrix} 2 \\ 1 \\ 3 \\ 2 \end{smallmatrix}$ is a module of length 4, of radical length 2, and with top isomorphic to the direct sum of the simple modules corresponding to vertices 1 and 2.

Figure 1 gives an illustration of the category W_Λ . The vertices are the sets of indecomposable objects in each wide subcategory. A non-identity morphism $g_{\mathcal{T}}^W : W \rightarrow W'$ (so that \mathcal{T} is an indecomposable support τ -rigid object in $C(W)$ and $J_W(\mathcal{T}) = W'$) is shown as an arrow between W and W' labelled by T . When P is projective in W we have $J_W(P) = J_W(P[1])$, and there are two corresponding maps, g_P^W and $g_{P[1]}^W$ from W to $J_W(P)$; in this case we draw a doubled arrow labelled only by P . Wide subcategories of rank 1 have generally been shown more than once in the figure, and the corresponding vertices should be identified.

REFERENCES

[1] T. Adachi, O. Iyama and I. Reiten, *τ -tilting theory*, *Compos. Math.* 150 (2015), no. 3, 415–452
 [2] I. Assem, D. Simson and A. Skowroński, *Elements of the representation theory of associative algebras. Vol. 1. Techniques of Representation Theory*, London Mathematical Society Student Texts, 65. Cambridge University Press, Cambridge, 2006.
 [3] M. Auslander and S. O. Smalø, *Almost split sequences in subcategories*, *J. Algebra* 69 (1981) 426–454. Addendum; *J. Algebra* 71 (1981), 592–594
 [4] A. B. Buan and R. J. Marsh, *Signed τ -exceptional sequences*, preprint, arXiv:1802.01169 [math.RT]
 [5] L. Demonet, O. Iyama and G. Jasso, *τ -tilting finite algebras, bricks and g -vectors*, preprint, arXiv:1503.00285 [math.RT], to appear in *International Mathematics Research Notices*




FIGURE 1. Illustration of the category W_Λ

- [6] L. Demonet, O. Iyama, N. Reading, I. Reiten and H. Thomas, *Lattice theory of torsion classes* preprint, arXiv:1711.01785 [math.RT]
- [7] K. Igusa and G. Todorov, *Signed exceptional sequences and the cluster morphism category*, preprint, arXiv:1706.02041 [math.RT]
- [8] C. Ingalls and H. Thomas, *Noncrossing partitions and representation of quivers*, Compos. Math. 145 (2009), no. 6, 1533–1562
- [9] G. Jasso, *Reduction of τ -tilting modules and torsion pairs*, Int. Math. Res. Not. IMRN (2015), no. 16, 7190–7237
- [10] Y. Mizuno, *Classifying τ -tilting modules over preprojective algebras of Dynkin type*, Math. Z, 277(3-4), (2014), 665–690
- [11] A. Neeman, *Triangulated categories*, Annals of Mathematics Studies, 148. Princeton University Press, Princeton, NJ, 2001

DEPARTMENT OF MATHEMATICAL SCIENCES, NORWEGIAN UNIVERSITY OF SCIENCE AND TECHNOLOGY, 7491 TRONDHEIM, NORWAY

E-mail address: aslak.buan@ntnu.no

SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS LS2 9JT UK

E-mail address: marsh@maths.leeds.ac.uk