

# MULTIPLICATIVE QUIVER VARIETIES AND GENERALISED RUIJSENAARS-SCHNEIDER MODELS

OLEG CHALYKH AND MAXIME FAIRON

ABSTRACT. We study some classical integrable systems naturally associated with multiplicative quiver varieties for the (extended) cyclic quiver with  $m$  vertices. The phase space of our integrable systems is obtained by quasi-Hamiltonian reduction from the space of representations of the quiver. Three families of Poisson-commuting functions are constructed and written explicitly in suitable Darboux coordinates. The case  $m = 1$  corresponds to the tadpole quiver and the Ruijsenaars–Schneider system and its variants, while for  $m > 1$  we obtain new integrable systems that generalise the Ruijsenaars–Schneider system. These systems and their quantum versions also appeared recently in the context of supersymmetric gauge theory and cyclotomic DAHAs [BEF, BFN1, BFN2, KN], as well as in the context of the Macdonald theory [CE].

## 1. INTRODUCTION

Among the most powerful geometric techniques in the theory of integrable systems is the method of Hamiltonian (or symplectic) reduction. Invented initially for reducing the degrees of freedom in Hamiltonian systems with symmetries, the concept of a moment map and symplectic reduction have since found a multitude of uses beyond their initial scope. One of the earlier examples of a Hamiltonian reduction was given by Kazhdan, Kostant and Sternberg [KKS], who demonstrated how to obtain the celebrated Calogero–Moser system from a very simple system on  $T^*\mathfrak{gl}_n$ . Since then, many integrable systems have been obtained or interpreted by similar methods. Among those is a remarkable generalisation of the Calogero–Moser system introduced by Ruijsenaars and Schneider [RS]. The latter system was interpreted in terms of an infinite-dimensional symplectic reduction by Nekrasov [N1], extending his earlier work with A. Gorsky [GN]. Hyperbolic Ruijsenaars–Schneider system can also be obtained by a finite-dimensional reduction in the spirit of [KKS], as was demonstrated by Fock and Rosly [FR]. Although Fock and Rosly employ Hamiltonian (or Poisson) reduction, their construction allows an interpretation in terms of *quasi-Hamiltonian* reduction. We recall that the method of quasi-Hamiltonian reduction was developed by Alekseev, Malkin and Meinrenken in [AMM], see also [AKSM]. The main difference is that the reduction is performed on a space which may not be symplectic, and the moment map takes values in the Lie group rather than the Lie algebra. Not attempting at a comprehensive review, we refer the reader to some of the more recent papers [P, FK1, FK2, FK3, M, FG], where the Ruijsenaars–Schneider model and its variants are treated by the method of (quasi-)Hamiltonian reduction, and where further references can be found. Let us also mention an alternative geometric approach to many-body problems by Krichever [Kr1], in which the Lax matrix structure play the central role instead, and the Hamiltonian picture is derived from that, cf. [KrP, Kr2, KrS].

From yet another perspective, a unified view onto the Calogero–Moser and Ruijsenaars–Schneider system can be achieved by noticing that in both cases the reduction is done on (the cotangent bundle to) the space of representations of a one-loop quiver. Such a view onto the (complexified) Calogero–Moser system was brought forward by G. Wilson’s work [W] relating the rational Calogero–Moser system, adelic Grassmannian, and the KP hierarchy, and it has been further deepened in [BW1], [BW2], cf. [BC, BP]. The present paper stems from a natural idea to look for a generalisation of Wilson’s results for more complicated quivers. We recall that with any quiver Nakajima associates in [N2] a class of symplectic quotients called *quiver varieties*. There exists also a multiplicative version of quiver varieties, introduced by Crawley-Boevey and Shaw [CBS] and interpreted via quasi-Hamiltonian reduction by Van den Bergh [VdB1]. Affine Dynkin quivers are a particularly well studied class, and a large part of Wilson’s (and Berest–Wilson’s) results have already been extended to this case by Ginzburg, Baranovsky and Kuznetsov [BGK1], [BGK2] (see also [Esh], [CS]). However, the multiplicative case has not been systematically looked at, apart from the

already mentioned case of a one-loop quiver. This was the main motivation behind our work. We will focus on the link to integrable particle dynamics; other aspects of the Calogero–Moser correspondence will be discussed elsewhere. Our main result is a construction of new generalisations of the Ruijsenaars–Schneider system, related to cyclic quivers. This is achieved by performing a quasi-Hamiltonian reduction on the space of representations of the associated multiplicative pre-projective algebras  $\Lambda^q$  of Crawley-Boevey and Shaw [CBS]. Our main tool is the formalism of double (quasi-)Poisson algebras due to Van den Bergh [VdB1, VdB2]. The constructed integrable systems come equipped with a complete phase space (represented by a suitable multiplicative quiver variety), and the associated Hamiltonian dynamics can be explicitly integrated. By constructing Darboux coordinates on the phase space, we express the new integrable Hamiltonians in coordinates, which then allows us to identify them as generalisations of the Ruijsenaars–Schneider system. For non-multiplicative quiver varieties, analogous integrable systems can be identified with the rational Calogero–Moser system for  $W = S_n \wr \mathbb{Z}_m$ , see [CS]. Thus, the systems constructed in the present work can be considered as  $q$ -analogues of the rational Calogero–Moser system for such  $W$ . The very fact that such  $q$ -analogues exist is somewhat surprising. Indeed, the group  $S_n \wr \mathbb{Z}_m$  is noncrystallographic for general  $m$ , while trigonometric or hyperbolic Calogero–Moser systems and their  $q$ -analogues are usually expected to have a crystallographic symmetry group.

Interestingly, quantum versions of some of these systems appeared in a seemingly unrelated context in [CE], where they were called *twisted Macdonald–Ruijsenaars systems*. By computing some of these quantum Hamiltonians explicitly, we are able to see this relationship in the case of the quiver with two vertices, although a direct comparison in the general case is more difficult. However, a recent work by Braverman, Etingof and Finkelberg [BEF], which appeared while we were finishing the present paper, clarifies this connection rather remarkably. It introduces a cyclotomic version of the double affine Hecke algebra (DAHA) in type  $A$ . Inside the cyclotomic DAHA there are three natural commutative subalgebras and they give rise to quantum integrable systems, in the same way as the usual DAHA can be used to produce the Macdonald–Ruijsenaars operators. The classical versions of these systems correspond to the  $q = 1$  limit of the cyclotomic DAHA (cf. [O] for the case of the usual DAHA), and this leads to the multiplicative quiver varieties for the cyclic quiver. Thus, the integrable systems constructed in [BEF] coincide (on the classical level) with those constructed by us. The interpretation of these integrable systems via the cyclotomic DAHA in [BEF] allows to explain their relationship to the twisted Macdonald–Ruijsenaars systems from [CE] in type  $A$ . Our methods are quite different in comparison, and they allow us to find explicit formulas for the corresponding classical Hamiltonians and integrate the Hamiltonian flows (the approach via the cyclotomic DAHA in [BEF] is less explicit). Curiously, these Hamiltonians become much simpler under the Cherednik–Fourier transform. In this form they appeared in the work of Braverman, Finkelberg, and Nakajima [BFN1, BFN2] on the quantized Coulomb branch of quiver gauge theories, see also a related work of Kodera and Nakajima [KN]. This can also be seen from our formulas at the classical level, when the Cherednik–Fourier transform becomes the angle-action transform studied by Ruijsenaars [R]. See Section §5 below for more details. Apart from being more explicit compared to [BEF], our approach also has an advantage of being better suited for studying spin versions of the Ruijsenaars–Schneider system and its generalisations; this will be a subject of a future work.

The structure of the paper is as follows. In Section §2 we first describe the general formalism of double Poisson brackets and quasi-Poisson algebras due to Van den Bergh [VdB1], and then exemplify it for the multiplicative quiver varieties. Section §3 looks at the tadpole quiver, explaining how to obtain the hyperbolic Ruijsenaars–Schneider system by quasi-Hamiltonian reduction. In Section §4 we consider the multiplicative quiver varieties (Calogero–Moser spaces) for the framed cyclic quiver with  $m$  vertices. We introduce three Poisson commuting families of functions on those Calogero–Moser spaces, and integrate the corresponding Hamiltonian flows. We then write these Hamiltonians in suitable Darboux coordinates, identifying them as generalisations of the hyperbolic Ruijsenaars–Schneider system. Finally, in Section §5 we discuss the relationship between our work and the results of [CE] and [BFN1, BFN2, KN, BEF]. In particular, we were able to write explicitly the integrable quantum Hamiltonians from [BEF] in the case of a quiver with two vertices. The paper finishes with three appendices containing some of the more technical proofs.

*Acknowledgement.* It is our pleasure to thank L. Fehér, J.F. van Diejen, P. Etingof, P. Iliev, I. Marshall, S. Ruijsenaars for useful comments and discussions. We are especially grateful to

P. Etingof who shared with us the results of [BEF] before their publication. This allowed us to streamline some of our considerations in Section §4. A part of this paper was written during the first author's stay in November 2016 at the Centro Internacional de Ciencias (CIC), Cuernavaca. He thanks the organisers of the programme "Integrable and quasi-integrable systems" and CIC for the hospitality and excellent working conditions. The first author was partially supported by EPSRC under grant EP/K004999/1. The second author is supported by a University of Leeds 110 Anniversary Research Scholarship.

## 2. PRELIMINARIES

In this section we first recap the theory of double Poisson brackets and double (quasi-)Poisson algebras due to Van den Bergh [VdB1]. We then describe a concrete example of this formalism, related to multiplicative preprojective algebras and multiplicative quiver varieties of Crawley-Boevey and Shaw [CBS]. We will follow the notation of the papers [VdB1, VdB2], where the reader can find many more details.

**2.1. Double brackets and double derivations.** We fix an algebra  $A$  over  $\mathbb{C}$ . For an element  $a \in A \otimes A$ , we will use a shorthand notation  $a' \otimes a''$  for  $\sum_i a'_i \otimes a''_i$ . We set  $a^\circ = a'' \otimes a'$ . More generally, for any  $s \in S_n$  we define  $\tau_s : A^{\otimes n} \rightarrow A^{\otimes n}$  by  $\tau_s(a_1 \otimes \dots \otimes a_n) = a_{s^{-1}(1)} \otimes \dots \otimes a_{s^{-1}(n)}$ . Thus,  $a^\circ = \tau_{(12)}a$ . The multiplication map  $m : A^{\otimes n} \rightarrow A$  is  $m(a_1 \otimes \dots \otimes a_n) = a_1 \dots a_n$ . We view  $A^{\otimes n}$  as an  $A$ -bimodule via the *outer* bimodule structure  $b(a_1 \otimes \dots \otimes a_n)c = ba_1 \otimes \dots \otimes a_n c$ . If  $B$  is a  $\mathbb{C}$ -algebra, then we say that  $A$  is a  $B$ -algebra if it is equipped with a  $\mathbb{C}$ -algebra map  $B \rightarrow A$ .

An  $n$ -*bracket* is a linear map  $\{\{-, \dots, -\}\} : A^{\otimes n} \rightarrow A^{\otimes n}$  which is a derivation in its last argument for the outer bimodule structure on  $A^{\otimes n}$ , and which is cyclically anti-symmetric:

$$\tau_{(1\dots n)} \circ \{\{-, \dots, -\}\} \circ \tau_{(1\dots n)}^{-1} = (-1)^{n+1} \{\{-, \dots, -\}\}.$$

If  $A$  is a  $B$ -algebra then we assume that the bracket is  $B$ -linear, i.e. it vanishes when its last argument is in the image of  $B$ . We call a 2- and a 3-bracket respectively a *double* and a *triple bracket*. In particular, a double bracket satisfies  $\{\{a, b\}\} = -\{\{b, a\}\}^\circ$  and  $\{\{a, bc\}\} = b\{\{a, c\}\} + \{\{a, b\}\}c$ . Any double bracket  $\{\{-, -\}\}$  defines an induced triple bracket  $\{\{-, -, -\}\}$  given by

$$\{\{a, b, c\}\} = \{\{a, \{\{b, c\}\}'\}\} \otimes \{\{b, c\}\}'' + \tau_{(123)} \{\{b, \{\{c, a\}\}'\}\} \otimes \{\{c, a\}\}'' + \tau_{(123)}^2 \{\{c, \{\{a, b\}\}'\}\} \otimes \{\{a, b\}\}''.$$

A double bracket on  $A$  is called a *double Poisson* bracket if the associated triple bracket vanishes. For any double (respectively, double Poisson) bracket  $\{\{-, -\}\}$ , the bracket  $\{-, -\} := m \circ \{\{-, -\}\}$  descends to an antisymmetric biderivation (respectively, a Lie bracket) on  $A/[A, A]$  [VdB1, 2.4.1, 2.4.6].

Following [CB], we call the elements of  $D_{A/B} := \text{Der}_B(A, A \otimes A)$  *double derivations*, and we make  $D_{A/B}$  into an  $A$ -bimodule by using the *inner* bimodule structure on  $A \otimes A$ : if  $\delta \in D_{A/B}$  and  $a, b, c \in A$ , then  $(b\delta c)(a) = \delta(a)'c \otimes b\delta(a)''$ . Let  $D_B A := T_A D_{A/B}$  be the tensor algebra of this bimodule; this is a graded algebra, with  $A$  placed in degree 0 and  $D_{A/B}$  in degree 1. The elements of degree  $n$  in  $D_B A$  can be used to define  $n$ -brackets on  $A$ :

**Proposition 2.1.** ([VdB1, 4.1.1]) *There is a well-defined linear map  $\mu : (D_B A)_n \rightarrow \{B\text{-linear } n\text{-brackets on } A\}$ ,  $Q \mapsto \{\{-, \dots, -\}\}_Q$  which on  $Q = \delta_1 \dots \delta_n$  is given by*

$$\begin{aligned} \{\{-, \dots, -\}\}_Q &= \sum_{i=0}^{n-1} (-1)^{(n-1)i} \tau_{(1\dots n)}^i \circ \{\{-, \dots, -\}\}_Q \circ \tau_{(1\dots n)}^{-i}, \\ \{\{a_1, \dots, a_n\}\}_Q &= \delta_n(a_n)' \delta_1(a_1)'' \otimes \delta_1(a_1)' \delta_2(a_2)'' \otimes \dots \otimes \delta_{n-1}(a_{n-1})' \delta_n(a_n)'' . \end{aligned}$$

*The map  $\mu$  factors through  $D_B A/[D_B A, D_B A]$  (for the graded commutator).*

The algebra  $D_B A$  may be viewed as a noncommutative version of the algebra of polyvector fields: according to [VdB1, 3.2.2]  $D_B A$  admits a canonical *double Schouten–Nijenhuis bracket*, which makes  $D_B A$  into a double Gerstenhaber algebra. We denote this bracket as  $\{\{-, -\}\}_{\text{SN}}$ , and we compose it with the multiplication on  $D_B A$  to obtain  $\{-, -\}_{\text{SN}} := m \circ \{\{-, -\}\}_{\text{SN}}$ .

**2.2. Double quasi-Poisson algebras.** We now assume that  $B$  is commutative and semi-simple, i.e.  $B = \mathbb{C}e_1 \oplus \dots \oplus \mathbb{C}e_n$  with  $e_i e_j = \delta_{ij} e_i$ . We define for all  $i$  a double derivation  $E_i \in D_{A/B}$  such that  $E_i(a) = ae_i \otimes e_i - e_i \otimes ae_i$ . A *double quasi-Poisson bracket* on  $A$  is a  $B$ -linear bracket  $\{\{-, -\}\}$ , such that the induced triple bracket satisfies  $\{\{-, -, -\}\} = \frac{1}{12} \sum_i \{\{-, -, -\}\}_{E_i^3}$ , where the brackets in the right-hand side are defined in Proposition 2.1. In this case, we say that  $A$  is a *double quasi-Poisson algebra*.

A *multiplicative moment map* for a double quasi-Poisson algebra  $(A, \{\{-, -\}\})$  is an element  $\Phi = \sum_i \Phi_i$  with  $\Phi_i \in e_i A e_i$  such that we have  $\{\{\Phi_i, a\}\} = \frac{1}{2}(\Phi_i E_i + E_i \Phi_i)(a)$  for all  $a \in A$ . When a double quasi-Poisson algebra is equipped with a multiplicative moment map, we say that it is a *quasi-Hamiltonian algebra*.

Assume that there is an element  $P \in (D_B A)_2$  such that  $\{P, P\}_{\text{SN}} = \frac{1}{6} \sum_i E_i^3 \text{ mod } [D_B A, D_B A]$  (for the graded commutator). Then we say that  $A$  is a *differential double quasi-Poisson algebra* with the *differential double quasi-Poisson bracket*  $\{\{-, -\}\}_P$ . This implies that  $\{\{-, -\}\}_P$  is a double quasi-Poisson bracket [VdB1, 4.2.3].

**2.3. Representation spaces.** For a  $\mathbb{C}$ -algebra  $A$  and any  $N \in \mathbb{N}$ , the representation space  $\text{Rep}(A, N)$  is the affine scheme that parametrises algebra homomorphisms  $\varrho : A \rightarrow \text{Mat}_N(\mathbb{C})$ . The coordinate ring  $\mathcal{O}(\text{Rep}(A, N))$  is generated by the functions  $a_{ij}$  for  $a \in A$ ,  $i, j = 1, \dots, N$  defined by  $a_{ij}(\varrho) = \varrho(a)_{ij}$  at any point  $\varrho \in \text{Rep}(A, N)$ . The functions  $a_{ij}$  are linear in  $a$  and satisfy the relations  $(ab)_{ij} = \sum_k a_{ik} b_{kj}$ . We can therefore associate with any  $a \in A$  a *matrix-valued function*  $\mathcal{X}(a) := (a_{ij})_{i,j=1,\dots,N}$  on  $\text{Rep}(A, N)$ . Similarly, any double derivation  $\delta \in \text{Der}(A, A \otimes A)$  gives rise to a matrix-valued vector field  $\mathcal{X}(\delta) = (\delta_{ij})_{i,j=1,\dots,N}$  on  $\text{Rep}(A, N)$ , where  $\delta_{ij}$  is a derivation of  $\mathcal{O}(\text{Rep}(A, N))$  defined by the rule  $\delta_{ij}(a_{uv}) = \delta'(a)_{uj} \delta''(a)_{iv}$ .

Everything can also be defined in a relative setting, i.e. for a  $B$ -algebra  $A$ , where  $B$  is of the form  $B = \mathbb{C}e_1 \oplus \dots \oplus \mathbb{C}e_n$  with  $e_i^2 = e_i$ . Representation spaces are now indexed by  $n$ -tuples  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$ . Given  $\alpha$  with  $\alpha_1 + \dots + \alpha_n = N$ , we embed  $B$  diagonally into  $\text{Mat}_N(\mathbb{C})$  so that  $\text{Id}_N$  is split into a sum of  $n$  diagonal blocks of size  $\alpha_1, \dots, \alpha_n$ , representing the idempotents  $e_i$ . By definition,  $\text{Rep}_B(A, \alpha) = \text{Hom}_B(A, \text{Mat}_N(\mathbb{C}))$ , and it can be viewed as an affine scheme in the same way as  $\text{Rep}(A, N)$ . Note in particular that for any  $\Phi \in \oplus_i e_i A e_i$ , the matrix-valued function  $\mathcal{X}(\Phi)$  on  $\text{Rep}_B(A, \alpha)$  takes values in block matrices  $\prod_i \text{Mat}_{\alpha_i}(\mathbb{C})$ .

On  $\text{Rep}(A, N)$  we have a natural action of  $\text{GL}_N$ , induced by conjugation action on  $\text{Mat}_N(\mathbb{C})$ . Similarly, we have an action of  $\text{GL}_\alpha = \prod_i \text{GL}_{\alpha_i}$  on  $\text{Rep}_B(A, \alpha)$ .

**2.4. Quasi-Poisson manifolds.** A double quasi-Poisson bracket on an algebra  $A$  makes its representation space into a quasi-Poisson manifold. Let us first recall the geometric setup of [AKSM], following the notation of [VdB1, 7.13]. Let  $M$  be a  $G$ -manifold, for  $G$  a Lie group whose Lie algebra  $\mathfrak{g}$  admits a non-degenerate  $G$ -invariant bilinear form  $(-, -)$ . If  $(e_a)$  is a basis of  $\mathfrak{g}$  and  $(e^a)$  the dual basis with respect to  $(-, -)$ , the Cartan 3-tensor is defined as  $\phi = \frac{1}{12} C^{abc} e_a \wedge e_b \wedge e_c$ , for  $C^{abc} = (e^a, [e^b, e^c])$  the tensor of structure constants. Write  $\xi^L$  and  $\xi^R$  respectively to denote the left and right invariant vector fields on  $G$ , for  $\xi \in \mathfrak{g}$ .

The  $G$ -action on  $M$  gives rise to a Lie algebra homomorphism  $(-)_M : \mathfrak{g} \rightarrow \text{Der } \mathcal{O}(M)$ , which can be extended to polyvector fields to define a 3-tensor  $\phi_M$ . We say that  $M$  is a quasi-Poisson manifold if there exists an invariant bivector field  $P$  on  $M$  such that its Schouten-Nijenhuis bracket with itself satisfies  $[P, P] = \phi_M$ . One associates with  $P$  a bracket on  $\mathcal{O}(M)$  defined by  $\{f, g\} = P(df, dg)$ . A *multiplicative moment map* is an  $\text{Ad}$ -equivariant map  $\Phi : M \rightarrow G$  satisfying

$$\{g \circ \Phi, -\} = \frac{1}{2} (e^a)_M ((e_a^L + e_a^R)(g) \circ \Phi), \quad (2.1)$$

for all functions  $g \in \mathcal{O}(G)$ . A *Hamiltonian quasi-Poisson manifold* is such a triple  $(M, P, \Phi)$ . In the case where the action of  $G$  on  $M$  is free and proper, for each conjugacy class  $\mathcal{C}_g$  of  $g \in G$  the subset  $\Phi^{-1}(\mathcal{C}_g)/G$  is a Poisson manifold, and this process is called *quasi-Hamiltonian reduction*.

Now let us turn to geometric structures induced on representation spaces of a double quasi-Poisson algebra  $A$ . Assume that  $\{\{-, -\}\} : A \times A \rightarrow A \otimes A$  is a  $B$ -linear double bracket on  $A$ .

**Proposition 2.2.** ([VdB1, 7.5.1, 7.5.2, 7.8, 7.12.2]) *There is a unique antisymmetric biderivation (bivector)  $\{\{-, -\}\} : \mathcal{O}(\text{Rep}_B(A, \alpha)) \times \mathcal{O}(\text{Rep}_B(A, \alpha)) \rightarrow \mathcal{O}(\text{Rep}_B(A, \alpha))$  such that for all  $a, b \in A$ ,*

$$\{a_{ij}, b_{uv}\} = \{\{a, b\}'_{uj}\} \{\{a, b\}''_{iv}\}. \quad (2.2)$$

If  $\{\{-, -\}\}$  is a double Poisson bracket, then this bivector is Poisson so  $\mathcal{O}(\text{Rep}_B(A, \alpha))$  is a Poisson algebra.

**Theorem 2.3.** [VdB1, 7.8, 7.13.2] *Assume that  $(A, P)$  is a differential double quasi-Poisson algebra, which is quasi-Hamiltonian for the multiplicative moment map  $\Phi \in \oplus_i e_i A e_i$ . We have that  $\text{Rep}_B(A, \alpha)$  is a  $\text{GL}_\alpha$ -space with a quasi-Poisson bracket  $\{-, -\}$  determined from  $\{\{-, -\}\}_P$  by (2.2). Then the matrix-valued function  $\mathcal{X}(\Phi) : \text{Rep}_B(A, \alpha) \rightarrow \prod_i \text{Mat}_{\alpha_i}(k)$  is a (geometric) multiplicative moment map for  $\text{Rep}_B(A, \alpha)$ . Therefore,  $\text{Rep}_B(A, \alpha)$  (if smooth) admits a structure of a Hamiltonian quasi-Poisson manifold.*

**2.5. Multiplicative preprojective algebras.** Let  $Q = (Q, I)$  be a quiver with vertex set  $I$  and arrow set  $Q$ . Let  $\bar{Q}$  denote the double of  $Q$ , obtained by adjoining to every arrow  $a \in Q$  its opposite,  $a^*$ . Define the maps  $t, h : \bar{Q} \rightarrow I$  that associate to every arrow  $a$  its tail and head,  $t(a)$  and  $h(a)$ . In particular,  $t(a) = h(a^*)$  and  $h(a) = t(a^*)$ . We define  $\epsilon : \bar{Q} \rightarrow \{\pm 1\}$  the sign function which associates 1 to every arrow of  $Q$  and  $-1$  to each arrow of  $\bar{Q} \setminus Q$ . We write  $\mathbb{C}\bar{Q}$  for the path algebra of  $\bar{Q}$ ; it is generated by the idempotents  $e_i$  (zero paths) associated to the vertices  $i \in I$ , and arrows  $a \in \bar{Q}$ , with multiplication given by concatenation of paths. We view  $\mathbb{C}\bar{Q}$  as a  $B$ -algebra, with  $B = \oplus_{i \in I} \mathbb{C}e_i$ . Finally, we extend  $*$  to an involution on  $\mathbb{C}\bar{Q}$  by setting  $(a^*)^* = a$ .

**Remark 2.4.** *We will use the convention that the composition of paths in  $\mathbb{C}\bar{Q}$  is written from left to right, i.e.,  $ab$  means “ $a$  followed by  $b$ ”, with  $ab = 0$  if  $h(a) \neq t(b)$ .*

Let  $A$  be obtained from  $\mathbb{C}\bar{Q}$  by inverting all elements  $(1 + aa^*)_{a \in \bar{Q}}$ . For all  $a \in \bar{Q}$ , define the element  $\frac{\partial b}{\partial a}$  of  $D_B A$  which on  $b \in \bar{Q}$  acts as

$$\frac{\partial b}{\partial a} = \begin{cases} e_{t(a)} \otimes e_{h(a)} & \text{if } a = b \\ 0 & \text{otherwise} \end{cases} \quad (2.3)$$

Fix an arbitrary ordering  $<$  on  $\bar{Q}$  and consider the following element  $\Phi \in A$ :

$$\Phi = \prod_{a \in \bar{Q}} (1 + aa^*)^{\epsilon(a)}, \quad (2.4)$$

where the product is taken with respect to the chosen ordering  $<$ . Following [CBS], given  $q = \sum_{i \in I} q_i e_i$  with  $q_i \in \mathbb{C}^\times$ , we define the deformed *multiplicative preprojective algebra* as the quotient  $\Lambda^q = A/(\Phi - q)$ . Up to isomorphism, the algebra  $\Lambda^q$  is independent of the ordering [CBS, Theorem 1.4].

**Theorem 2.5.** ([VdB1, 6.7.1.]) *The algebra  $A$  has a quasi-Hamiltonian structure given by*

$$P = \frac{1}{2} \sum_{a \in \bar{Q}} \epsilon(a) (1 + a^* a) \frac{\partial}{\partial a} \frac{\partial}{\partial a^*} - \frac{1}{2} \sum_{a < b \in \bar{Q}} \left( \frac{\partial}{\partial a^*} a^* - a \frac{\partial}{\partial a} \right) \left( \frac{\partial}{\partial b^*} b^* - b \frac{\partial}{\partial b} \right), \quad (2.5)$$

and the multiplicative moment map given by (2.4).

Note that the corresponding double-Poisson bracket is defined also on  $\mathbb{C}\bar{Q}$ , but the elements  $1 + aa^*$  need to be invertible to define the moment map  $\Phi$ .

The corresponding double quasi-Poisson bracket on  $\mathbb{C}\bar{Q}$  is calculated as follows.

**Proposition 2.6.** *Suppose that the arrows of  $\bar{Q}$  are ordered in such a way that  $a < a^* < b < b^*$  for any  $a < b \in Q$ . Let  $\{\{-, -\}\}$  be the double bracket associated to the bivector (2.5). Then one has:*

$$\{\{a, a\}\} = \frac{1}{2} \epsilon(a) (a^2 \otimes e_{t(a)} - e_{h(a)} \otimes a^2) \quad (a \in \bar{Q}), \quad (2.6a)$$

$$\begin{aligned} \{\{a, a^*\}\} &= e_{h(a)} \otimes e_{t(a)} + \frac{1}{2} a^* a \otimes e_{t(a)} \\ &\quad + \frac{1}{2} e_{h(a)} \otimes aa^* + \frac{1}{2} (a^* \otimes a - a \otimes a^*) \delta_{h(a), t(a)} \quad (a \in Q), \end{aligned} \quad (2.6b)$$

$$\begin{aligned} \{\{a, b\}\} &= \frac{1}{2} e_{h(a)} \otimes ab + \frac{1}{2} ba \otimes e_{t(a)} \\ &\quad - \frac{1}{2} (b \otimes a) \delta_{h(a), h(b)} - \frac{1}{2} (a \otimes b) \delta_{t(a), t(b)} \quad (a, b \in \bar{Q}, a < b, b \neq a^*). \end{aligned} \quad (2.6c)$$

All other brackets are obtained by using  $\{\{a, b\}\} = -\{\{b, a\}\}^\circ$ , with  $(c' \otimes c'')^\circ = c'' \otimes c'$ . Note that (2.6a) is zero unless  $t(a) = h(a)$  (i.e.  $a$  is a loop), and (2.6c) is zero unless  $a$  and  $b$  share a vertex.

*Proof.* We give a proof for (2.6a) with  $a \in Q$ , other formulas are similar. First, recall that if  $\delta_1, \delta_2 \in D_{A/B}$ , then the double bracket associated with  $\delta_1 \delta_2 \in (D_B A)_2$  is given in Proposition 2.1:

$$\{\{a, b\}\}_{\delta_1 \delta_2} = \delta_2(b)' \delta_1(a)'' \otimes \delta_1(a)' \delta_2(b)'' - \delta_1(b)' \delta_2(a)'' \otimes \delta_2(a)' \delta_1(b)'' . \quad (2.7)$$

When calculating  $\{\{a, a\}\}_P$ , the only nonzero contribution comes from the bivector

$$P_a = \frac{1}{2} \left( a \frac{\partial}{\partial a} \right) \left( \frac{\partial}{\partial a} a \right) ,$$

which we write as

$$P_a = \frac{1}{2} U^- U^+ , \quad U^+ = \frac{\partial}{\partial a} a, \quad U^- = a \frac{\partial}{\partial a} . \quad (2.8)$$

Using the inner bimodule structure on  $D_{A/B}$ , we get

$$\begin{aligned} U^+(a) &= e_{t(a)} a \otimes e_{h(a)} = a \otimes e_{h(a)} , \\ U^-(a) &= e_{t(a)} \otimes a e_{h(a)} = e_{t(a)} \otimes a . \end{aligned} \quad (2.9)$$

Hence, combining this with (2.7) we find:

$$\begin{aligned} \{\{a, a\}\}_P &= \{\{a, a\}\}_{P_a} = \frac{1}{2} U^+(a)' U^-(a)'' \otimes U^-(a)' U^+(a)'' - \frac{1}{2} U^-(a)' U^+(a)'' \otimes U^+(a)' U^-(a)'' \\ &= \frac{1}{2} a^2 \otimes e_{t(a)} e_{h(a)} - \frac{1}{2} e_{t(a)} e_{h(a)} \otimes a^2 . \end{aligned}$$

Since  $e_{t(a)} e_{h(a)} = e_{t(a)} = e_{h(a)}$  if  $a$  is a loop and zero otherwise, we obtain (2.6a) for  $a \in Q$ . Similar calculation for  $\{\{a^*, a^*\}\}_P$  leads to the same result with the overall minus, which explains  $\epsilon(a)$  in (2.6a).  $\square$

For any  $a \in A$ , define  $\text{tr}(a) = \sum_{i=1}^{|\alpha|} a_{ii}$ ; this is a  $\text{GL}_\alpha$ -invariant function on  $\text{Rep}_B(A, \alpha)$ . We have the following useful formula [VdB1, Proposition 7.7.3]:

$$\{\text{tr}(a), \text{tr}(b)\} = \text{tr}\{a, b\} . \quad (2.10)$$

Here the bracket on the left is the one induced on  $\mathcal{O}(\text{Rep}_B(A, \alpha))$  by (2.2), while  $\{a, b\}$  in the right-hand side stands for the bracket on  $A$  obtained from the double bracket:

$$\{a, b\} = m \circ \{\{a, b\}\} = \{\{a, b\}\}' \{\{a, b\}\}'' . \quad (2.11)$$

We finish this subsection with two remarks. The first remark is that a total ordering on  $\bar{Q}$  is not necessary to define a quasi-Hamiltonian structure on  $A$ . Indeed, a bivector  $P$  can be obtained by fusion, see [VdB1, Proof of Theorem 6.7.1]. The construction only requires to order arrows that start at any given vertex  $i$ , and the resulting bivector  $P$  can be written as

$$P = \frac{1}{2} \sum_{a \in \bar{Q}} \epsilon(a) (1 + a^* a) \frac{\partial}{\partial a} \frac{\partial}{\partial a^*} - \frac{1}{2} \sum_{i \in I} \sum_{\substack{a < b \in \bar{Q} \\ t(a) = t(b) = i}} F_a F_b , \quad (2.12)$$

where

$$F_a = \frac{\partial}{\partial a^*} a^* - a \frac{\partial}{\partial a} .$$

The moment map  $\Phi$  is also well-defined, since each  $\Phi_i = e_i \Phi e_i$  can be written as

$$\Phi_i = \prod_{a \in \bar{Q}, t(a)=i} (e_i + a a^*)^{\epsilon(a)} , \quad (2.13)$$

so the order of the factors is only needed to be prescribed at each vertex.

Another remark is about a slight modification of Theorem 2.5 which will be useful later. Let  $S \subset \bar{Q}$  such that for all  $a \in S$  we have  $a^* \in S$ . Write  $1_S : \bar{Q} \rightarrow \{0, 1\}$  for the characteristic function of the subset  $S$ ; we have  $1_S(a) = 1_S(a^*)$  for all  $a \in \bar{Q}$ .



**Theorem 2.7.** *Let  $A_S$  be obtained from  $\mathbb{C}\bar{Q}$  by inverting all elements  $(1_S(a) + aa^*)_{a \in \bar{Q}}$ . The algebra  $A_S$  has a quasi-Hamiltonian structure given by*

$$P_S = \frac{1}{2} \sum_{a \in \bar{Q}} \epsilon(a)(1_S(a) + a^*a) \frac{\partial}{\partial a} \frac{\partial}{\partial a^*} - \frac{1}{2} \sum_{a < b \in \bar{Q}} \left( \frac{\partial}{\partial a^*} a^* - a \frac{\partial}{\partial a} \right) \left( \frac{\partial}{\partial b^*} b^* - b \frac{\partial}{\partial b} \right), \quad (2.14)$$

and the multiplicative moment map  $\Phi_S = (\Phi_{S,i})_{i \in I}$  given by

$$\Phi_{S,i} = \prod_{a \in \bar{Q}, t(a)=i} (1_S(a)e_i + aa^*)^{\epsilon(a)}. \quad (2.15)$$

*Proof.* In the case  $S = Q$  this is Theorem 2.5, and the general case can be proved by the same method. Alternatively,  $A_S$  can be obtained from  $A$  by inverting arrows in  $Q$ . Indeed, take  $a : i \rightarrow j$  in  $Q \setminus S$  and adjoin to  $A$  the formal inverse of  $a$ , i.e. an element  $a^{-1}$  such that  $e_j a^{-1} = a^{-1} e_i = a^{-1}$ ,  $aa^{-1} = e_i$ ,  $a^{-1}a = e_j$ . Set  $\hat{a} = a^{-1} + a^*$ , then  $e_i + aa^* = a\hat{a}$  and  $e_j + a^*a = \hat{a}a$ . Define  $\hat{a} = a$  for  $a \in Q \setminus S$ , while for  $a \in S$  set  $\hat{a} := a^*$ . Therefore, for all  $a \in Q$  the elements  $1_S(a)e_{t(a)} + a\hat{a}$  are invertible in  $A_S$ . Inside  $A_S$  we have a subalgebra (over  $B$ ), isomorphic to  $\mathbb{C}\bar{Q}$  and generated by all  $a, \hat{a}$  with  $a \in Q$ . One can now define double derivations  $\frac{\partial}{\partial a}$  and  $\frac{\partial}{\partial \hat{a}}$  on this subalgebra, in the same way as we define  $\frac{\partial}{\partial a}, \frac{\partial}{\partial a^*}$  on  $\mathbb{C}\bar{Q}$  in (2.3). One can think of this as a change of variables from  $a, a^*$  to  $a, \hat{a}$ ; then a simple calculation shows that under this change of variables,

$$\frac{\partial}{\partial a} \mapsto \frac{\partial}{\partial a} - a^{-1} \frac{\partial}{\partial \hat{a}} a^{-1}, \quad \frac{\partial}{\partial a^*} \mapsto \frac{\partial}{\partial \hat{a}}.$$

Substituting this into (2.5) and then renaming  $\hat{a}$  as  $a^*$ , we obtain (modulo graded commutators) the bivector (2.14), as claimed.  $\square$

Explicit formulas for the double bracket on  $A_S$  are almost the same, the only difference is (2.6b) which gets replaced by

$$\{\!\{a, a^*\}\!\} = 1_S(a)e_{h(a)} \otimes e_{t(a)} + \frac{1}{2}a^*a \otimes e_{t(a)} + \frac{1}{2}e_{h(a)} \otimes aa^* + \frac{1}{2}(a^* \otimes a - a \otimes a^*)\delta_{h(a),t(a)}. \quad (2.16)$$

**2.6. Multiplicative quiver varieties.** Let us turn now to representation spaces. Below we will always work in a relative setting, and from now on we will drop the subscript  $B$  from the notation. For instance, given  $\alpha \in \mathbb{N}^I$ , a representation  $\text{Rep}(\mathbb{C}\bar{Q}, \alpha)$  will always mean  $\text{Rep}_B(\mathbb{C}\bar{Q}, \alpha)$ , where  $B = \bigoplus_i \mathbb{C}e_i$ , with  $e_i$  acting as the identity on  $V_i = \mathbb{C}^{\alpha_i}$ . For each arrow  $a \in \bar{Q}$ , we have  $a = e_{t(a)}ae_{h(a)}$ , therefore,  $a$  is represented by a matrix with at most one non-zero block of size  $\alpha_{t(a)} \times \alpha_{h(a)}$ . Therefore, this can be viewed as a *quiver representation*, consisting of vector spaces  $\mathcal{V}_i = \mathbb{C}^{\alpha_i}$ ,  $i \in I$  and linear maps  $X_a : \mathcal{V}_{h(a)} \rightarrow \mathcal{V}_{t(a)}$  for each  $a \in \bar{Q}$ .<sup>1</sup> With this interpretation, we have

$$X_a \in \text{Mat}_{\alpha_{t(a), \alpha_{h(a)}}}(\mathbb{C}), \quad \text{Rep}(\mathbb{C}\bar{Q}, \alpha) \cong \prod_{a \in \bar{Q}} \text{Mat}_{\alpha_{t(a), \alpha_{h(a)}}}(\mathbb{C}). \quad (2.17)$$

Next,  $\text{Rep}(A, \alpha)$  is an affine open subset of  $\text{Rep}(\mathbb{C}\bar{Q}, \alpha)$ , so it is smooth. By Theorems 2.3, 2.5, this is a quasi-Hamiltonian manifold, with the quasi-Poisson bracket determined by  $P$  and with a multiplicative moment map  $\mathcal{X}(\Phi)$ . The representation space  $\text{Rep}(\Lambda^q, \alpha)$  is a level set  $\{\Phi = q\}$  of the momentum map, so it is a closed affine subvariety in  $\text{Rep}(A, \alpha)$ . Let  $q^\alpha = \prod_{i \in I} q^{\alpha_i}$ . Then  $\text{Rep}(\Lambda^q, \alpha)$  is empty unless  $q^\alpha = 1$  [CBS, Lemma 1.5]. Isomorphism classes of representations correspond to orbits under the group

$$G(\alpha) = \left( \prod_{i \in I} \text{GL}_{\alpha_i} \right) / \mathbb{C}^\times, \quad (2.18)$$

acting by conjugation. Here  $\mathbb{C}^\times$  denotes the diagonal subgroup of scalar matrices. Semi-simple representations correspond to closed orbits.

Consider the affine variety  $\mathcal{S}_{\alpha, q} := \text{Rep}(\Lambda^q, \alpha) // G(\alpha)$ , whose points correspond to semi-simple representations of  $\Lambda^q$  of dimension  $\alpha$ . We will mostly deal with the situation when  $q$  and  $\alpha$  are such that all representations in  $\text{Rep}(\Lambda^q, \alpha)$  are simple. In this case, we have the following result which is a combination of [CBS, Theorems 1.8 & 1.10] and [VdB1, Proposition 1.7].

<sup>1</sup>If this looks to the reader as a representation of the opposite quiver, that is because of our convention for composing arrows, see Remark 2.4.

**Theorem 2.8.** *Let  $p(\alpha) = 1 + \sum_{a \in Q} \alpha_{t(a)} \alpha_{h(a)} - \alpha \cdot \alpha$ , where  $\alpha \cdot \alpha = \sum_{i \in I} \alpha_i^2$ . Suppose that  $\text{Rep}(\Lambda^q, \alpha)$  is non-empty and all representations in  $\text{Rep}(\Lambda^q, \alpha)$  are simple. Then  $\alpha$  is a positive root of  $Q$  and  $\text{Rep}(\Lambda^q, \alpha)$  is a smooth affine variety of dimension  $g + 2p(\alpha)$ , with  $g = \dim G(\alpha) = \alpha \cdot \alpha - 1$ . The group  $G(\alpha)$  acts freely on  $\text{Rep}(\Lambda^q, \alpha)$ , so  $\mathcal{S}_{\alpha, q} = \text{Rep}(\Lambda^q, \alpha)/G(\alpha)$  is a Poisson manifold of dimension  $2p(\alpha)$ , obtained by quasi-Hamiltonian reduction.*

The Poisson bracket on  $\mathcal{O}(\mathcal{S}_{\alpha, q}) = \mathcal{O}(\text{Rep}(\Lambda^q, \alpha))^{G(\alpha)}$  is obtained from 2.2, 2.5. Moreover, it follows from [VdB2, 8.3.1] that this Poisson bracket is non-degenerate, so the variety  $\mathcal{S}_{\alpha, q}$  is, in fact, symplectic. The varieties  $\mathcal{S}_{\alpha, q}$  are sometimes referred to as *multiplicative quiver varieties*, though we will reserve this name for a special case related to framed quivers, described below.

Let  $Q$  be an arbitrary quiver with the vertex set  $I$ . A *framing* of  $Q$  is a quiver  $\tilde{Q}$  that has one additional vertex, denoted  $\infty$ , and a number of arrows  $i \rightarrow \infty$  from the vertices of  $Q$  (multiple arrows are allowed). Given  $\alpha \in \mathbb{N}^I$  and  $q \in (\mathbb{C}^\times)^I$ , we extend them from  $Q$  to  $\tilde{Q}$  by putting  $\alpha_\infty = 1$  and  $q_\infty = q^{-\alpha}$ . Thus, we put

$$\tilde{\alpha} = (1, \alpha), \quad \tilde{q} = q^{-\alpha} e_\infty + \sum_{i \in I} q_i e_i. \quad (2.19)$$

Consider the representation space  $\text{Rep}(\Lambda^{\tilde{q}}, \tilde{\alpha})$  for the multiplicative preprojective algebra of  $\tilde{Q}$ . The quotients

$$\mathcal{M}_{\alpha, q}(Q) := \text{Rep}(\Lambda^{\tilde{q}}, \tilde{\alpha}) // G(\tilde{\alpha}) \quad (2.20)$$

are called multiplicative quiver varieties. Note that since  $\alpha_\infty = 1$ , we have

$$G(\tilde{\alpha}) = \left( \prod_{i \in I \sqcup \{\infty\}} \text{GL}_{\alpha_i} \right) / \mathbb{C}^\times \cong \prod_{i \in I} \text{GL}_{\alpha_i} = \text{GL}_\alpha. \quad (2.21)$$

We say that  $q = \sum_{i \in I} q_i e_i$  is *regular* if  $q^\alpha \neq 1$  for any root  $\alpha$  of the quiver  $Q$ . We have the following result, cf. [N2, Theorem 2.8], [BCE, Proposition 3].

**Proposition 2.9.** *Choose an arbitrary framing  $\tilde{Q}$  of  $Q$  and let  $\tilde{\alpha}$  and  $\tilde{q}$  be as in (2.19). If  $q$  is regular, then every module of dimension  $\tilde{\alpha}$  over the multiplicative preprojective algebra  $\Lambda^{\tilde{q}}$  is simple. Hence, the group  $\text{GL}_\alpha$  acts freely on  $\text{Rep}(\Lambda^{\tilde{q}}, \tilde{\alpha})$  and the multiplicative quiver variety  $\mathcal{M}_{\alpha, q}(Q)$  is smooth.*

*Proof.* Let  $V$  be a  $\Lambda^{\tilde{q}}$ -module of dimension  $\tilde{\alpha} = (1, \alpha)$ . If  $V$  is non-simple, then it has a proper submodule  $U \subset V$ . The dimension vector of  $U$  is either of the form  $(1, \beta)$  or  $(0, \beta)$ , for some  $\beta \in \mathbb{N}^I$ . In the latter case, by passing to a submodule, we may assume that  $U$  is simple. But then  $\beta$  must be a positive root of  $Q$  and  $q^\beta = 1$ , by [CBS, Lemma 1.5 & Theorem 1.8]. Therefore,  $q$  cannot be regular. In the case when  $\dim U = (1, \beta)$  we consider the quotient module  $V/U$  and repeat the argument.  $\square$

It follows that if  $q$  is regular and  $\mathcal{M}_{\alpha, q}(Q) \neq \emptyset$ , then  $\tilde{\alpha} = (1, \alpha)$  is a positive root of  $\tilde{Q}$  and  $\mathcal{M}_{\alpha, q}(Q)$  is a smooth affine variety of dimension  $2p(\tilde{\alpha})$ .

**Remark 2.10.** *The varieties  $\mathcal{M}_{\alpha, q}(Q)$  are the same as framed multiplicative quiver varieties studied by Yamakawa [Y] (with the zero stability parameter), see also the Appendix by Nakajima and Yamakawa in [BEF].*

### 3. TADPOLE QUIVER

In this section we describe the way to obtain the hyperbolic Ruijsenaars–Schneider system by a quasi-Hamiltonian reduction. The main results in this section are not new, see e.g. [R], [FR], [O], [FK3], but we provide self-contained proofs which will serve as a preparation for the later sections. We should stress that we focus on algebraic and geometric aspects, working over  $\mathbb{C}$ . Therefore, we do not address more subtle questions about various real, compact and non-compact forms of the complexified system, see [R], [FK3] and references therein. Note that the choice of a real form is crucial for studying the particle dynamics and scattering, cf. [R].

Let  $Q$  be a tadpole quiver with vertices  $\{\infty, 0\}$  and two arrows,  $x : 0 \rightarrow 0$  and  $v : 0 \rightarrow \infty$ . Let us write  $y = x^*$ ,  $w = v^*$  for the opposite arrows. We choose an ordering  $x < y < v < w$  on



$\bar{Q}$ . As before, form an algebra  $A$  by adjoining  $\{(1 + aa^*)^{-1}\}_{a \in \bar{Q}}$  to the path algebra  $\mathbb{C}\bar{Q}$ . The quasi-Poisson bracket on  $A$  is given in Proposition 2.6:

$$\{\{x, x\}\} = \frac{1}{2}(x^2 \otimes e_0 - e_0 \otimes x^2), \quad \{\{y, y\}\} = \frac{1}{2}(e_0 \otimes y^2 - y^2 \otimes e_0), \quad (3.1a)$$

$$\{\{x, y\}\} = e_0 \otimes e_0 + \frac{1}{2}yx \otimes e_0 + \frac{1}{2}e_0 \otimes xy + \frac{1}{2}(y \otimes x - x \otimes y), \quad (3.1b)$$

$$\{\{v, v\}\} = \{\{w, w\}\} = 0, \quad \{\{v, w\}\} = e_\infty \otimes e_0 + \frac{1}{2}e_\infty \otimes vw + \frac{1}{2}vw \otimes e_0, \quad (3.1c)$$

$$\{\{x, v\}\} = \frac{1}{2}e_0 \otimes xv - \frac{1}{2}x \otimes v, \quad \{\{x, w\}\} = \frac{1}{2}wx \otimes e_0 - \frac{1}{2}w \otimes x, \quad (3.1d)$$

$$\{\{y, v\}\} = \frac{1}{2}e_0 \otimes yv - \frac{1}{2}y \otimes v, \quad \{\{y, w\}\} = \frac{1}{2}wy \otimes e_0 - \frac{1}{2}w \otimes y. \quad (3.1e)$$

If we further localise  $A$  by adding  $x^{-1}$ , then we can replace  $y$  by  $z = y + x^{-1}$ , and the brackets between  $x, z$  are very similar, cf. Proposition 2.7 and (2.16):

$$\{\{z, z\}\} = \frac{1}{2}(e_0 \otimes z^2 - z^2 \otimes e_0), \quad \{\{x, z\}\} = \frac{1}{2}zx \otimes e_0 + \frac{1}{2}e_0 \otimes xz + \frac{1}{2}(z \otimes x - x \otimes z). \quad (3.2)$$

Let us calculate some further brackets. Let  $\{-, -\}$  denote the bracket  $A \times A \rightarrow A$  defined by (2.11). This bracket is not anti-symmetric in general, but it satisfies Leibniz's rule in the second argument, and by [VdB1, 2.4 & Proposition 5.1.2]  $A$  is a left Loday algebra. I.e., we have the following identities:

$$\{a, bc\} = \{a, b\}c + b\{a, c\}, \quad \{a, \{b, c\}\} = \{\{a, b\}, c\} + \{b, \{a, c\}\}. \quad (3.3)$$

**Proposition 3.1.** *We have the following identities in  $A$  for all  $a, b \geq 0$ :*

$$\{x^a, x^b\} = 0, \quad \{y^a, y^b\} = 0, \quad \{(xy)^a, (xy)^b\} = 0.$$

*If we further localise  $A$  on  $x$ , then we also have*

$$\{z^a, z^b\} = 0, \quad z = y + x^{-1}.$$

**Proposition 3.2.** *Let  $A'$  denote the algebra  $A$  localised on  $x$ , and  $z = y + x^{-1}$ . For any  $a, b \geq 0$  we have*

$$\{x^a, zx^b\} = axz^{a+b} \pmod{[A', A']}, \quad (3.4a)$$

$$\{zx^a, zx^b\} = \sum_{r=1}^a zx^r zx^{a+b-r} - \sum_{r=1}^b zx^r zx^{a+b-r} \pmod{[A', A']}. \quad (3.4b)$$

Proofs can be found in Appendix §A.  $\square$

For  $q = (q_\infty, q_0)$ , the multiplicative preprojective algebra  $\Lambda^q$  is the quotient of  $A$  by the relation

$$(1 + xy)(1 + yx)^{-1}(1 + vw)(1 + wv)^{-1} = q_0 e_0 + q_\infty e_\infty. \quad (3.5)$$

Multiplication by the idempotents  $e_\infty, e_0$  turns this into two relations:

$$(e_0 + xy)(e_0 + yx)^{-1}(e_0 + vw) = q_0 e_0, \quad (3.6a)$$

$$(e_\infty + wv)^{-1} = q_\infty e_\infty. \quad (3.6b)$$

Choose a dimension vector  $\alpha = (1, n)$  and set  $q_\infty = q_0^{-n}$  to satisfy  $q^\alpha = 1$ . A representation of  $\Lambda^q$  of dimension  $\alpha$  is a pair  $(\mathcal{V}_\infty, \mathcal{V}_0) = (\mathbb{C}, \mathbb{C}^n)$  together with linear maps representing arrows of  $\bar{Q}$  and satisfying (3.6a), (3.6b). Denote the matrices representing the arrows as  $X, Y, V, W$ . Therefore, points of  $\text{Rep}(\Lambda^q, \alpha)$  are represented by quadruples  $(X, Y, V, W)$ ,

$$X, Y \in \text{Mat}_{n \times n}(\mathbb{C}), \quad V \in \text{Mat}_{n \times 1}(\mathbb{C}), \quad W \in \text{Mat}_{1 \times n}(\mathbb{C}),$$

satisfying

$$(\text{Id}_n + XY)(\text{Id}_n + YX)^{-1}(\text{Id}_n + VW) = q_0 \text{Id}_n, \quad (3.7a)$$

$$(1 + WV)^{-1} = q_\infty \quad (q_\infty = q_0^{-n}). \quad (3.7b)$$

The group  $\text{GL}_n(\mathbb{C})$  acts on these quadruples by

$$g \cdot (X, Y, V, W) = (gXg^{-1}, gYg^{-1}, gV, Wg^{-1}), \quad g \in \text{GL}_n, \quad (3.8)$$

and the orbits in  $\text{Rep}(\Lambda^q, \alpha) // \text{GL}_n$  correspond to isomorphism classes of semisimple representations. Introduce the *Calogero–Moser space*  $\mathcal{C}_{n,q_0}$  as

$$\mathcal{C}_{n,q_0} = \text{Rep}(\Lambda^q, \alpha) // \text{GL}_n .$$

This is a multiplicative quiver variety for a framed one-loop quiver, and applying the results of §2.6, we have

**Proposition 3.3.** *Suppose  $q_0$  is not a root of unity. Then the group  $\text{GL}_n$  acts on  $\text{Rep}(\Lambda^q, \alpha)$  freely, and  $\mathcal{C}_{n,q_0}$  is a smooth symplectic variety of dimension  $2n$ .*

The variety  $\mathcal{C}_{n,q_0}$  admits a description in terms of pairs of matrices as follows:

$$\mathcal{C}_{n,q_0} = \{X, Y \in \text{Mat}_{n \times n}(\mathbb{C}) \mid \text{rank}((\text{Id}_n + XY)(\text{Id}_n + YX)^{-1} - q_0 \text{Id}_n) = 1\} // \text{GL}_n .$$

We may also consider the open subset  $\mathcal{C}_{n,q_0}^0 \subset \mathcal{C}_{n,q_0}$  on which  $X$  is invertible. Introducing  $Z := Y + X^{-1}$ , we have  $\text{Id}_n + XY = XZ$ ,  $\text{Id}_n + YX = ZX$  and therefore

$$\mathcal{C}_{n,q_0}^0 = \{X, Z \in \text{GL}_n \mid \text{rank}(XZX^{-1}Z^{-1} - q_0 \text{Id}_n) = 1\} // \text{GL}_n .$$

The Poisson bracket on  $\mathcal{O}(\mathcal{C}_{n,q_0}) = \mathcal{O}(\text{Rep}(\Lambda^q, \alpha))^{\text{GL}_n(\mathbb{C})}$  is induced by the double bracket on  $A$ . Proposition 3.1 together with (2.10) give us

**Theorem 3.4.** *The following families of functions on  $\mathcal{C}_{n,q_0}$  are Poisson commuting:*

$$\{\text{tr} X^j \mid j \in \mathbb{N}\}, \quad \{\text{tr} Y^j \mid j \in \mathbb{N}\}, \quad \{\text{tr}(1 + XY)^j \mid j \in \mathbb{Z}\}, \quad \{\text{tr}(Y + X^{-1})^j \mid j \in \mathbb{Z}\},$$

where the last family is viewed on  $\mathcal{C}_{n,q_0}^0 \subset \mathcal{C}_{n,q_0}$ .

**Remark 3.5.** *If we assume  $Y$  invertible, we can get another commuting family  $\{\text{tr}(X + Y^{-1})^j \mid j \in \mathbb{Z}\}$ .*

To interpret these families as integrable particle systems, we next write them down in suitable canonical (Darboux) coordinates.

**3.1. Darboux coordinates.** We take  $\mathfrak{h} = \mathbb{C}^n$  with coordinates  $x_1, \dots, x_n$ , and define

$$\mathfrak{h}_{\text{reg}} = \{x \in \mathfrak{h} \mid x_i \neq 0, x_i \neq x_j, x_i \neq q_0 x_j \text{ for all } i \neq j\} .$$

Let  $\mathfrak{h}^\times = (\mathbb{C}^\times)^n$  with coordinates  $\nu_1, \dots, \nu_n \in \mathbb{C}^\times$ . We are going to define a map

$$\xi : \mathfrak{h}_{\text{reg}} \times \mathfrak{h}^\times \rightarrow \mathcal{C}_{n,q_0}^0 .$$

Given  $x \in \mathfrak{h}_{\text{reg}}$ ,  $\nu \in \mathfrak{h}^\times$ , we set  $\xi(x, \nu) = (X, Z)$  where

$$X = \text{diag}(x_1, \dots, x_n), \quad Z = (Z_{ij}), \quad \text{with } Z_{ij} = \frac{(q_0 - 1)\nu_j}{q_0 - x_i/x_j} .$$

One can check directly, using Cauchy’s determinant formula, that  $Z$  is invertible. We also have

$$(XZ - q_0 ZX)_{ij} = x_i Z_{ij} - q_0 Z_{ij} x_j = (1 - q_0)\nu_j x_j ,$$

which shows that the matrix  $XZX^{-1}Z^{-1} - q_0 \text{Id}_n$  has rank one, and so the pair  $(X, Z)$  determines a point of  $\mathcal{C}_{n,q_0}^0$ . Moreover, if one simultaneously permutes  $x_i$  and  $\nu_i$ , the resulting  $(X, Z)$  get conjugated by the matrix of that permutation. Therefore, we have in fact a map

$$\xi : \mathfrak{h}_{\text{reg}} \times \mathfrak{h}^\times / S_n \rightarrow \mathcal{C}_{n,q_0}^0 .$$

It is easy to see that  $\xi$  is injective, and since  $\dim \mathcal{C}_{n,q_0} = 2n$ , we can use  $(x_1, \dots, x_n, \nu_1, \dots, \nu_n)$  as local coordinates on  $\mathcal{C}_{n,q_0}^0$ . Note that by [O], the variety  $\mathcal{C}_{n,q_0}^0$  is connected, hence the image of  $\xi$  is dense.

**Proposition 3.6** ([FR], [O]). *The local diffeomorphism  $\xi : \mathfrak{h}_{\text{reg}} \times \mathfrak{h}^\times / S_n \rightarrow \mathcal{C}_{n,q_0}^0$  becomes a Poisson map if we equip  $\mathfrak{h}_{\text{reg}} \times \mathfrak{h}^\times$  with the following  $S_n$ -invariant Poisson bracket  $\{-, -\}'$ :*

$$\{x_i, x_j\}' = 0, \tag{3.9a}$$

$$\{x_i, \nu_j\}' = \delta_{ij} x_i \nu_j, \tag{3.9b}$$

$$\{\nu_i, \nu_j\}' = \frac{(1 - q_0)^2 (x_i + x_j) x_i x_j \nu_i \nu_j}{(x_i - x_j)(x_i - q_0 x_j)(x_j - q_0 x_i)}. \tag{3.9c}$$

*Proof.* We need to check that the map  $\xi$  satisfies  $\xi^*\{f, g\} = \{\xi^*f, \xi^*g\}'$  for any  $f, g \in \mathcal{O}(\mathcal{C}_{n, q_0}^0)$ . Since the functions  $f_a := \text{tr}(X^a)$  and  $g_b := \text{tr}(ZX^b)$  with  $a, b = 1, \dots, n$  form local coordinates at a generic point, it suffices to check the Poisson property for these functions (cf. the proof of Proposition 2.7 in [Et]). From (2.10) and Propositions 3.1, 3.2 we have:

$$\{f_a, f_b\} = 0, \quad \{f_a, g_b\} = ag_{a+b}, \quad \{g_b, g_c\} = \sum_{r=b+1}^c h_{r, b+c-r}, \quad (3.10)$$

where  $h_{r,s} := \text{tr}(ZX^r ZX^s)$  (we assume  $b < c$  in the last formula). Next,

$$\xi^*f_a = \sum_i x_i^a, \quad \xi^*g_b = \sum_i \nu_i x_i^b, \quad \xi^*h_{r, a+b-r} = \sum_{i,j} \frac{(q_0 - 1)^2 \nu_i \nu_j}{(q_0 - x_i/x_j)(q_0 - x_j/x_i)} x_j^r x_i^{a+b-r}. \quad (3.11)$$

Therefore,

$$\sum_{r=b+1}^a \xi^*h_{r, a+b-r} = (a-b) \sum_i \nu_i^2 x_i^{a+b} + \sum_{i \neq j} \frac{(q_0 - 1)^2 \nu_i \nu_j}{(q_0 - x_i/x_j)(q_0 - x_j/x_i)} \sum_{r=b+1}^a x_j^r x_i^{a+b-r}. \quad (3.12)$$

Now notice that (3.9a) and (3.9b) give

$$\{\xi^*f_a, \xi^*f_b\}' = \sum_{i,j} \{x_i^a, x_j^b\}' = 0, \quad \{\xi^*f_a, \xi^*g_b\}' = \sum_{i,j} \{x_i^a, \nu_j x_j^b\}' = a \sum_i x_i^{a+b} \nu_i,$$

which agrees with the first two formulas in (3.10). Next, we use (3.9c) to find that

$$\{\xi^*g_a, \xi^*g_b\}' = \sum_{i,j} \{\nu_i x_i^a, \nu_j x_j^b\}' = \sum_{i \neq j} \frac{(1 - q_0)^2 (x_i + x_j) \nu_i \nu_j x_i^a x_j^b}{(x_i - x_j)(x_i/x_j - q_0)(x_j/x_i - q_0)} + (a-b) \sum_i \nu_i^2 x_i^{a+b}.$$

It is easy to see that this expression coincides with (3.12).  $\square$

The coordinates  $(x_i, \nu_i)$  are not yet canonical since  $\{\nu_i, \nu_j\}' \neq 0$ . A set of log-canonical coordinates can be constructed analogously to [FR]. Namely, introduce

$$\sigma_i = \nu_i \prod_{j: j \neq i} \frac{1 - x_i x_j^{-1}}{1 - q_0 x_i x_j^{-1}} \quad (i = 1, \dots, n). \quad (3.13)$$

Then one checks directly that

$$\{x_i, x_j\}' = 0, \quad \{x_i, \sigma_j\}' = \delta_{ij} x_i \sigma_j, \quad \{\sigma_i, \sigma_j\}' = 0.$$

In these coordinates, we can identify some of the known integrable particle systems among those in Theorem 3.4. The best known is the one corresponding to the Hamiltonians  $\text{tr}(Y + X^{-1})^j$ . Namely, after writing  $Y + X^{-1}$  in coordinates  $x_i, \sigma_i$ , we have

$$Y + X^{-1} = Z, \quad \text{where } Z_{ij} = \sigma_j \frac{q_0 - 1}{q_0 - x_i x_j^{-1}} \prod_{k \neq j} \frac{1 - q_0 x_j x_k^{-1}}{1 - x_j x_k^{-1}}. \quad (3.14)$$

This gives

$$\text{tr } Z = \sum_{i=1}^n \sigma_i \prod_{j \neq i} \frac{1 - q_0 x_i x_j^{-1}}{1 - x_i x_j^{-1}}, \quad (3.15)$$

which is equivalent to the classical hyperbolic Ruijsenaars–Schneider Hamiltonian. In fact,  $Z$  after conjugation by  $\text{diag}(\sqrt{\nu_1}, \dots, \sqrt{\nu_n})$  and a change of notation coincides with the Lax matrix from [RS, Section 4].

**Remark 3.7.** *The Hamiltonian (3.15) is written in the so-called Macdonald form related to the original formulation from [RS] by a canonical change of variables. Namely, let us replace the variables  $\sigma_i$  with*

$$s_i = \sigma_i \prod_{j \neq i} \left( \frac{q_0 x_i - x_j}{x_i - q_0 x_j} \right)^{1/2}.$$

It is easy to see that the transformation from  $(x_i, \sigma_i)$  to  $(x_i, s_i)$  is canonical. In these new coordinates the Hamiltonian (3.15) becomes

$$\mathrm{tr} Z = \sum_{i=1}^n s_i \left( \prod_{j \neq i} \frac{(x_i - q_0 x_j)(q_0 x_i - x_j)}{(x_i - x_j)^2} \right)^{1/2},$$

which is easily seen to be equivalent to  $S_1$  in [RS, (2.27), (2.29)].

Other families in Theorem 3.4 lead to closely related integrable systems. The family  $\{\mathrm{tr} X^j\}_{j \in \mathbb{N}}$  is trivial, while the simplest Hamiltonians for the other two are:

$$\mathrm{tr} Y = \sum_{i=1}^n \sigma_i \prod_{j \neq i} \frac{1 - q_0 x_i x_j^{-1}}{1 - x_i x_j^{-1}} - \sum_{i=1}^n x_i^{-1}, \quad \mathrm{tr}(1 + XY) = \sum_{i=1}^n \sigma_i x_i \prod_{j \neq i} \frac{1 - q_0 x_i x_j^{-1}}{1 - x_i x_j^{-1}}.$$

The second Hamiltonian  $\mathrm{tr}(1 + XY)$  reduces to the Ruijsenaars–Schneider Hamiltonian by a canonical change of variables  $\tilde{x}_i = x_i$ ,  $\tilde{\sigma}_i = \sigma_i x_i$ . The first Hamiltonian  $\mathrm{tr} Y$ , as explained in [BEF, Remark 3.25], is related to the quantum system introduced by J.F. van Diejen [VD, VDE] and a closely related system considered by Baker and Forrester [BF]. In slightly different canonical coordinates, the classical Hamiltonian system described by  $\mathrm{tr} Y$  was also considered by P. Iliev [I]. See also recent works [M, FG, FM] devoted to the study of some special cases of the van Diejen’s system.

Note that the interpretation of these systems through quasi-Hamiltonian reduction achieves two things at once: a completed phase space allowing particles to coalesce, and explicit dynamics on the affine space  $\mathrm{Rep}(A, \alpha)$ . Since this is mostly well-familiar (cf. [RS, FR, CF, I]), we skip the details (see Proposition 4.7 below for the more general systems related to the cyclic quivers). Another benefit is that we can see rather easily the *self-duality* for this system, originally established by Ruijsenaars.

**Proposition 3.8** ([R], [FK3]). *The transformation  $(X, Z) \mapsto (Z, X)$  induces a Poisson map  $\varphi : \mathcal{C}_{n, q_0}^0 \rightarrow \mathcal{C}_{n, q_0^{-1}}^0$  (more precisely,  $\varphi$  changes the sign of the Poisson bracket).*

*Proof.* From the proof of Proposition 3.6, we know that the Poisson bracket on  $\mathcal{C}_{n, q_0}^0$  is completely determined by the brackets between the functions  $\mathrm{tr}(X^a Z^b)$  with  $a, b \geq 0$ . These brackets, on the other hand, are determined by the antisymmetric biderivation (bivector) on  $\mathrm{Rep}(\mathbb{C}\bar{Q}, \alpha)$ , defined according to (2.2). Using this and (3.1a), (3.2), we find that

$$\begin{aligned} \{X_{ij}, X_{uv}\} &= \frac{1}{2}(X^2)_{uj} \delta_{iv} - \frac{1}{2} \delta_{uj} (X^2)_{iv}, & \{Z_{ij}, Z_{uv}\} &= \frac{1}{2} \delta_{uj} (Z^2)_{iv} - \frac{1}{2} (Z^2)_{uj} \delta_{iv}, \\ \{X_{ij}, Z_{uv}\} &= \frac{1}{2} (ZX)_{uj} \delta_{iv} + \frac{1}{2} \delta_{uj} (XZ)_{iv} + \frac{1}{2} Z_{uj} X_{iv} - \frac{1}{2} X_{uj} Z_{iv}. \end{aligned}$$

It is now obvious that swapping  $X$  with  $Z$  in these formulas leads to a change of sign. Therefore,  $\{\mathrm{tr} X^a Z^b, \mathrm{tr} X^c Z^d\} = -\{\mathrm{tr} Z^a X^b, \mathrm{tr} Z^c X^d\}$  for all  $a, b, c, d \geq 0$ , as needed.  $\square$

For the later use, let us formulate the above proposition in coordinates. We already have log-canonical coordinates  $x = (x_1, \dots, x_n)$ ,  $\sigma = (\sigma_1, \dots, \sigma_n)$  on  $\mathcal{C}_{n, q_0}^0$ , obtained by taking  $X = \mathrm{diag}(x_1, \dots, x_n)$  and  $Z$  as in (3.14). The second set of coordinates  $z = (z_1, \dots, z_n)$ ,  $\theta = (\theta_1, \dots, \theta_n)$  on the same space is given by

$$Z = \mathrm{diag}(z_1, \dots, z_n), \quad X_{ij} = \theta_j \frac{q_0^{-1} - 1}{q_0^{-1} - z_i z_j^{-1}} \prod_{k \neq j} \frac{1 - q_0^{-1} z_j z_k^{-1}}{1 - z_j z_k^{-1}}. \quad (3.16)$$

Then the Proposition 3.8 can be reformulated by saying that the transformation  $(x, \sigma) \rightarrow (z, \theta)$  is anti-symplectic, i.e.

$$\{z_i, z_j\} = 0, \quad \{z_i, \theta_j\} = -\delta_{ij} z_i \theta_j, \quad \{\theta_i, \theta_j\} = 0. \quad (3.17)$$

**Remark 3.9.** *From the construction, it is obvious that the transformation  $(x, \sigma) \rightarrow (z, \theta)$  is an involution. In [R] the canonicity of that involution was established by a rather roundabout method. For a more natural geometric approach, see [FK3, FGNR]. This can also be deduced from the results of [O], where the space  $\mathcal{C}_{n, \tau}^0$  was related to the centre  $\mathcal{Z}(H_{1, \tau})$  of the double affine Hecke algebra  $H_{q, \tau}$ . Indeed, we have the Cherednik–Fourier transform  $\varepsilon : H_{q, \tau} \rightarrow H_{q^{-1}, \tau^{-1}}$  as in [O,*

3.5]. It is an algebra homomorphism, so in the classical limit  $q \rightarrow 1$  it gives a Poisson map  $\mathcal{Z}(H_{1,\tau}) \rightarrow \mathcal{Z}(H_{1,\tau^{-1}})$  which interchanges  $X$  and  $Z$  (the bracket changes sign because  $q$  goes to  $q^{-1}$  under  $\epsilon$ ).

**Remark 3.10.** The Calogero–Moser spaces  $\mathcal{C}_{n,q_0}$  and  $\mathcal{C}_{n,q_0}^0$  are  $q$ -analogues of Wilson’s Calogero–Moser spaces [W]. In [1] and [CN] they appeared in the context of the bispectral problem. They have also been studied from the point of view of non-commutative geometry, as moduli spaces of non-commutative instantons [KKO] and ideals of the quantum torus algebra (see [BRT] and references therein).

#### 4. CYCLIC QUIVERS

For  $m \geq 2$ , let  $Q$  be a framed cyclic quiver with the arrows  $x_i : i \rightarrow i+1$ ,  $i \in I = \mathbb{Z}/m\mathbb{Z}$ , and  $v : 0 \rightarrow \infty$ . Write  $y_i = x_i^* : i+1 \rightarrow i$  and  $w = v^* : \infty \rightarrow 0$  for the opposite arrows. We choose the following ordering of arrows around each vertex:

$$x_{i-1} < y_{i-1} < x_i < y_i \quad \text{for } i \neq 0, \quad x_{m-1} < y_{m-1} < x_0 < y_0 < v < w \quad \text{at } i = 0.$$

As before, we form an algebra  $A$  by adjoining to  $\mathbb{C}\bar{Q}$  the elements  $(1 + aa^*)^{-1}$  for all  $a \in \bar{Q}$ . Let  $\{\{-, -\}\}$  be a double bracket on  $A$ , associated to the bivector (2.12). It gives the following brackets between the arrows  $x_i, y_j$ :

$$\{\{x_i, x_j\}\} = \frac{1}{2}e_{i+1} \otimes x_i x_{i+1} \delta_{j,i+1} - \frac{1}{2}x_{i-1} x_i \otimes e_i \delta_{j,i-1}, \quad (4.1a)$$

$$\{\{y_i, y_j\}\} = \frac{1}{2}y_{i+1} y_i \otimes e_{i+1} \delta_{j,i+1} - \frac{1}{2}e_i \otimes y_i y_{i-1} \delta_{j,i-1}, \quad (4.1b)$$

$$\begin{aligned} \{\{x_i, y_j\}\} &= \delta_{i,j} \left( e_{i+1} \otimes e_i + \frac{1}{2}y_i x_i \otimes e_i + \frac{1}{2}e_{i+1} \otimes x_i y_i \right) \\ &\quad - \frac{1}{2}y_{i+1} \otimes x_i \delta_{j,i+1} + \frac{1}{2}x_i \otimes y_{i-1} \delta_{j,i-1}, \end{aligned} \quad (4.1c)$$

$$\begin{aligned} \{\{y_i, x_j\}\} &= -\delta_{i,j} \left( e_i \otimes e_{i+1} + \frac{1}{2}x_i y_i \otimes e_{i+1} + \frac{1}{2}e_i \otimes y_i x_i \right) \\ &\quad - \frac{1}{2}y_i \otimes x_{i+1} \delta_{j,i+1} + \frac{1}{2}x_{i-1} \otimes y_i \delta_{j,i-1}. \end{aligned} \quad (4.1d)$$

Set  $x = x_0 + \dots + x_{m-1}$ ,  $y = y_0 + \dots + y_{m-1}$ , so then  $x_i = e_i x = x e_{i+1}$ ,  $y_i = e_{i+1} y = y e_i$ . If we formally invert all  $x_i$ , then in the localised algebra we have  $x^{-1} x = x x^{-1} = 1$  with  $x^{-1} = \sum_{i \in I} x_i^{-1}$ . Introduce the following elements  $E_{\pm 1} \in A \otimes A$ :

$$E_1 = \sum_{i \in I} e_{i+1} \otimes e_i, \quad E_{-1} = \sum_{i \in I} e_{i-1} \otimes e_i.$$

With this notation, we have the following easily verified formulas:

**Lemma 4.1.**

$$\{\{x, x\}\} = \frac{1}{2}(E_1 x^2 - x^2 E_1), \quad \{\{y, y\}\} = \frac{1}{2}(y^2 E_{-1} - E_{-1} y^2), \quad (4.2a)$$

$$\{\{x, y\}\} = E_1 + \frac{1}{2}y x E_1 + \frac{1}{2}E_1 x y - \frac{1}{2}y E_1 x + \frac{1}{2}x E_1 y, \quad (4.2b)$$

$$\{\{y, x\}\} = -E_{-1} - \frac{1}{2}E_{-1} y x - \frac{1}{2}x y E_{-1} + \frac{1}{2}x E_{-1} y - \frac{1}{2}y E_{-1} x. \quad (4.2c)$$

**Remark 4.2.** In the above formulas we use the outer bimodule structure on  $A \otimes A$ . For example,  $x E_1 = \sum_{i,j} x_j (e_{i+1} \otimes e_i) = \sum_i x_i \otimes e_i$ .

Let  $\{-, -\}$  denote the bracket  $A \times A \rightarrow A$  defined by (2.11).

**Proposition 4.3.** We have the following identities in  $A$  for all integers  $a, b \geq 0$ :

$$\{x^a, x^b\} = 0, \quad \{y^a, y^b\} = 0, \quad \{(xy)^a, (xy)^b\} = 0.$$

If we further localise  $A$  by inverting  $x$ , then we also have

$$\{z^a, z^b\} = 0, \quad z = y + x^{-1}.$$

**Proposition 4.4.** *For any  $a, b, c \geq 0$  with  $a, b-1, c-1 \equiv 0 \pmod{m}$ , we have*

$$\{x^a, yx^b\} = ax^{a+b-1} + ayx^{a+b} \pmod{[A, A]}, \quad (4.3a)$$

$$\{yx^b, yx^c\} = (b-c)yx^{b+c-1} + \sum_{t=1}^b yx^t yx^{b+c-t} - \sum_{t=1}^c yx^t yx^{b+c-t} \pmod{[A, A]}. \quad (4.3b)$$

Proofs can be found in Appendix §A.

For  $\tilde{q} = q_\infty e_\infty + \sum_{i \in I} q_i e_i$ , the multiplicative preprojective algebra  $\Lambda^{\tilde{q}}$  is the quotient of  $A$  by the relations:

$$(e_i + y_{i-1}x_{i-1})^{-1}(e_i + x_i y_i) = q_i e_i \quad (i \neq 0), \quad (4.4a)$$

$$(e_0 + y_{m-1}x_{m-1})^{-1}(e_0 + x_0 y_0)(e_0 + vw) = q_0 e_0, \quad (4.4b)$$

$$(e_\infty + vw)^{-1} = q_\infty e_\infty. \quad (4.4c)$$

Choose a dimension vector  $\tilde{\alpha} = (1, \alpha)$  where  $\alpha \in (\mathbb{N}^\times)^I$  and set  $q_\infty = q^{-\alpha} := \prod_{i \in I} q_i^{-\alpha_i}$ . Recall that  $q \in (\mathbb{C}^\times)^I$  is regular if  $q^\beta \neq 1$  for any root of the (unframed) cyclic quiver. The roots for the cyclic quiver form the affine root system of type  $\tilde{A}_{m-1}$ , therefore the regularity is equivalent to the following conditions:

$$\prod_{i \leq k \leq j-1} q_k \neq t^n \text{ for any } n \in \mathbb{Z} \text{ and } 1 \leq i \leq j \leq m, \text{ where } t := \prod_{i \in I} q_i.$$

We use the convention that the product in the left-hand side is empty when  $i = j$ , so in particular  $t$  must not be a root of unity. Applying the results of §2.6, we have

**Proposition 4.5.** *For regular  $q$ , the variety  $\mathcal{M}_{\alpha, q} = \text{Rep}(\Lambda^{\tilde{q}}, \tilde{\alpha}) // G(\tilde{\alpha})$ , if non-empty, is smooth symplectic, of dimension  $2p(\tilde{\alpha}) = 2\alpha_0 + 2 \sum_{i \in I} \alpha_i(\alpha_{i+1} - \alpha_i)$ .*

A representation of  $\Lambda^{\tilde{q}}$  of dimension  $\alpha$  is a collection of vector spaces  $\mathcal{V}_\infty = \mathbb{C}$  and  $\mathcal{V}_i = \mathbb{C}^{\alpha_i}$ , together with linear maps representing the arrows and satisfying the relations of (4.4a)–(4.4c). Denote the matrices representing the arrows as  $X_i, Y_i, V, W$ , and set  $X = X_0 + \cdots + X_{m-1}$ ,  $Y = Y_0 + \cdots + Y_{m-1}$ . We can view  $X, Y$  as linear endomorphisms of  $\mathcal{V} := \oplus_{i \in I} \mathcal{V}_i$ . Proposition 4.3 together with (2.10) gives us

**Theorem 4.6.** *The following families of functions on  $\mathcal{M}_{\alpha, q}$  are Poisson commuting:*

$$\{\text{tr } X^{jm} \mid j \in \mathbb{N}\}, \quad \{\text{tr } Y^{jm} \mid j \in \mathbb{N}\}, \quad \{\text{tr}(1 + XY)^j \mid j \in \mathbb{Z}\}, \quad \{\text{tr}(Y + X^{-1})^{jm} \mid j \in \mathbb{Z}\},$$

where the last family is viewed on the open subset  $\mathcal{M}_{\alpha, q}^0 \subset \mathcal{M}_{\alpha, q}$  on which  $X$  is invertible.

We can integrate explicitly the associated Hamiltonian flows in the most interesting cases:

**Proposition 4.7.** *Let  $t$  denote the time flow associated to  $H_k := \frac{1}{k} \text{tr } Y^k$ ,  $k \in m\mathbb{N}$ . Given an initial position  $X(0), Y(0), V(0), W(0)$  on  $\mathcal{M}_{\alpha, q}$ , the solution  $X, Y, V, W$  at time  $t$  is given by*

$$X(t) = e^{-tY^k} X(0) + Y^{-1}(e^{-tY^k} - 1), \quad Y(t) = Y(0), \quad V(t) = V(0), \quad W(t) = W(0).$$

Similarly, if  $t$  denotes the time flow associated to  $G_k := \frac{1}{k} \text{tr } Z^k$  with  $Z = Y + X^{-1}$ , then the solution at time  $t$  is given by

$$X(t) = e^{-tZ^k} X(0), \quad Z(t) = Z(0), \quad V(t) = V(0), \quad W(t) = W(0).$$

In both cases the flows are complete: in the first case on the whole of  $\mathcal{M}_{\alpha, q}$ , and in the second case on the open subset  $\mathcal{M}_{\alpha, q}^0 \subset \mathcal{M}_{\alpha, q}$  where  $X$  is invertible.

The proof of the proposition is given in Appendix §B.

We will be particularly interested in the special choice of the dimension vector  $\alpha_i = n$  for all  $i \in I$ , because only in this case  $\mathcal{M}_{\alpha, q}^0$  is nonempty. Accordingly, we choose  $q \in (\mathbb{C}^\times)^I$  and set  $q_\infty = t^{-n}$ , where  $t = \prod_{i \in I} q_i$ . With this choice, points of  $\text{Rep}(\Lambda^{\tilde{q}}, \tilde{\alpha})$  are represented by a collection of  $X_i, Y_i, V, W$ ,

$$X_i, Y_i \in \text{Mat}_{n \times n}(\mathbb{C}), \quad V \in \text{Mat}_{n \times 1}(\mathbb{C}), \quad W \in \text{Mat}_{1 \times n}(\mathbb{C}),$$



satisfying

$$(\mathrm{Id}_n + Y_{i-1}X_{i-1})^{-1}(\mathrm{Id}_n + X_iY_i) = q_i \mathrm{Id}_n \quad (i \neq 0), \quad (4.5a)$$

$$(\mathrm{Id}_n + Y_{m-1}X_{m-1})^{-1}(\mathrm{Id}_n + X_0Y_0)(\mathrm{Id}_n + VW) = q_0 \mathrm{Id}_n, \quad (4.5b)$$

$$(1 + WV)^{-1} = t^{-n}. \quad (4.5c)$$

Here all factors are assumed invertible, so the last relation can be obtained from the others after taking determinants. The group  $G := \mathrm{GL}_n^m$  acts on these linear data by

$$X_i \mapsto g_i X_i g_{i+1}^{-1}, \quad Y_i \mapsto g_{i+1} Y_i g_i^{-1}, \quad V \mapsto g_0 V, \quad W \mapsto W g_0^{-1},$$

for any  $g = (g_0, \dots, g_{m-1}) \in G$ . The space of equivalence classes of such linear data will be referred to as the *Calogero–Moser space*  $\mathcal{C}_{n,q}(m)$ , or simply  $\mathcal{C}_{n,q}$  when it does not lead to a confusion. This is a special case of a multiplicative quiver variety, so we can apply Theorem 2.8 and Proposition 2.9.

**Proposition 4.8.** *For regular  $q$ , the Calogero–Moser space  $\mathcal{C}_{n,q}(m)$  is a smooth symplectic variety of dimension  $2n$ .*

In the next subsection we will introduce local coordinates on this space and calculate the Poisson bracket explicitly.

**4.1. Coordinates and Poisson bracket.** We choose a regular set of parameters  $q \in (\mathbb{C}^\times)^I$ ; it will be convenient to introduce a separate notation for the quantities

$$t_s := \prod_{0 \leq i \leq s} q_i \quad (s = 0, \dots, m-1).$$

We will keep using the symbol  $t$  for  $t_{m-1} = q_0 \dots q_{m-1}$ . Let us define a family of representations of  $\Lambda^{\tilde{q}}$  with  $\tilde{q} = (t^{-n}, q_0, \dots, q_{m-1})$  and of dimension  $\tilde{\alpha} = (1, n, \dots, n)$ . We will assume that  $X_i$  are invertible, so we can use  $Z_i = Y_i + X_i^{-1}$  to rewrite the relations (4.5a)–(4.5c) as

$$(Z_{i-1}X_{i-1})^{-1}X_iZ_i = q_i \mathrm{Id}_n \quad (i \neq 0), \quad (4.6a)$$

$$(Z_{m-1}X_{m-1})^{-1}X_0Z_0(\mathrm{Id}_n + VW) = q_0 \mathrm{Id}_n, \quad (4.6b)$$

$$(1 + WV)^{-1} = t^{-n}. \quad (4.6c)$$

Then by changing bases we can achieve that  $X_s = \mathrm{Id}_n$  for  $s = 0, \dots, m-2$ . If we introduce  $A := X_{m-1}$  and  $B := q_0^{-1}Z_0$ , then from (4.6a) we find that  $Z_s = t_s B$  for  $s = 0, \dots, m-2$  and  $Z_{m-1} = tA^{-1}B$ . Using this in (4.6b) gives that  $A^{-1}B^{-1}AB(1 + VW) = t \mathrm{Id}_n$ , which we can rewrite as

$$ABA^{-1}B^{-1}(1 + \tilde{V}\tilde{W}) = t \mathrm{Id}_n, \quad (4.7)$$

where  $\tilde{V} = BAV$  and  $\tilde{W} = WB^{-1}A^{-1}$ . It is easy to see that isomorphic representation of  $\Lambda^{\tilde{q}}$  produce isomorphic quadruples  $(A, B, \tilde{V}, \tilde{W})$ , up to the equivalence (3.8). Therefore, we obtain

**Proposition 4.9** (cf. [BEF], Section 5.2). *Let  $\mathcal{C}_{n,q}^0(m)$  and  $\mathcal{C}_{n,t}^0$  be the Calogero–Moser spaces of isomorphism classes of linear data (4.6a)–(4.6c) and (4.7), respectively. We assume that the parameters are regular, so both spaces are smooth varieties. Then the map  $\xi$  sending  $(A, B, \tilde{V}, \tilde{W})$  to  $X_s = \mathrm{Id}_n$ ,  $Z_s = t_s B$  for  $s = 0, \dots, m-2$  and  $X_{m-1} = A$ ,  $Z_{m-1} = tA^{-1}B$ ,  $V = A^{-1}B^{-1}\tilde{V}$ ,  $W = \tilde{W}AB$  defines an isomorphism of these varieties. In particular,  $\mathcal{C}_{n,q}^0(m)$  is connected because so is  $\mathcal{C}_{n,t}^0$ .*

In fact, a stronger claim is true. Recall, that both spaces are Poisson varieties.

**Proposition 4.10.** *The isomorphism  $\xi : \mathcal{C}_{n,q}^0 \rightarrow \mathcal{C}_{n,q}^0(m)$  from the above proposition is a Poisson map.*

A proof can be found in Appendix §C.

As a result, we can construct canonical Darboux coordinates on  $\mathcal{C}_{n,q}^0(m)$  by transferring them from  $\mathcal{C}_{n,t}^0$ . Namely, let us choose  $A, B$  as suggested by (3.13), (3.14):

$$A = \mathrm{diag}(x_1, \dots, x_n), \quad B_{ij} = \sigma_j \frac{t-1}{t-x_i x_j^{-1}} \prod_{k \neq j} \frac{1-tx_j x_k^{-1}}{1-x_j x_k^{-1}}. \quad (4.8)$$

Then (4.7) determines  $\widetilde{V}, \widetilde{W}$  (uniquely, up to a simultaneous rescaling). By mapping these  $A, B, \widetilde{V}, \widetilde{W}$  to  $X_s, Z_s, V, W$  as described in Proposition 4.9, we obtain local coordinates  $x = (x_1, \dots, x_n)$ ,  $\sigma = (\sigma_1, \dots, \sigma_n)$  on  $\mathcal{C}_{n,q}^0(m)$ . Then the results of Section §3 combined with Proposition 4.10 tell us that

$$\{x_i, x_j\} = 0, \quad \{x_i, \sigma_j\} = \delta_{ij} x_i \sigma_j, \quad \{\sigma_i, \sigma_j\} = 0.$$

**Remark 4.11.** By [BEF, Theorem 5.17], the Calogero–Moser space  $\mathcal{C}_{n,q}(m)$  is connected. Therefore,  $\mathcal{C}_{n,q}^0(m)$  is a dense open subset, so the above coordinates  $x, \sigma$  can be used as local coordinates on both spaces.

**4.2. Generalised Ruijsenaars–Schneider models.** Having obtained canonical coordinates on the Calogero–Moser space  $\mathcal{C}_{n,q}(m)$ , we can now turn to the Hamiltonians from Theorem 4.6. The Hamiltonians  $E_{j,m} := \text{tr}(X^{j,m}) = m \text{tr}(X_0 \dots X_{m-1})^j = m \text{tr} A^j = m \sum_{i=1}^n x_i^j$  are trivial. Next, we have

$$F_{m,j} := \text{tr}(1 + XY)^j = \sum_{i \in I} \text{tr}(\text{Id}_n + X_i Y_i)^j = \sum_{i \in I} \text{tr}(X_i Z_i)^j = \sum_{i \in I} (t_s)^j \text{tr} B^j.$$

Since  $B$  is the Lax matrix for the Ruijsenaars–Schneider model, we do not find anything new here. Now let us look at

$$G_{m,j} := \text{tr}(Y + X^{-1})^{j,m} = m \text{tr}(Z_{m-1} \dots Z_0)^j = (t_0 \dots t_{m-1})^j \text{tr}(A^{-1} B^m)^j.$$

Ignoring the constant factor, the simplest Hamiltonian is  $G_{m,1} := \text{tr}(A^{-1} B^m)$ , with  $A, B$  given by (4.8); this formula also makes sense for  $m = 0, 1$ . Here are explicit formulas for  $G_{m,1}$  with  $m \leq 3$ ,

where we use the shorthand notation  $\Upsilon_{ij} = \frac{1 - tx_i x_j^{-1}}{1 - x_i x_j^{-1}} = \frac{t - x_j x_i^{-1}}{1 - x_j x_i^{-1}}$ :

$$G_{0,1} = \sum_{i=1}^n x_i^{-1}, \quad G_{1,1} = \sum_{i=1}^n \sigma_i x_i^{-1} \prod_{a \neq i} \Upsilon_{ia}, \quad (4.9a)$$

$$G_{2,1} = \sum_{i=1}^n \sigma_i^2 x_i^{-1} \prod_{a \neq i} \Upsilon_{ia}^2 + \sum_{i < j} \sigma_i \sigma_j \frac{(t-1)^2 (x_i^{-1} + x_j^{-1})}{(1 - x_i x_j^{-1})(1 - x_j x_i^{-1})} \prod_{a \neq i, j} \Upsilon_{ia} \Upsilon_{ja}, \quad (4.9b)$$

$$G_{3,1} = \sum_{i=1}^n \sigma_i^3 x_i^{-1} \prod_{a \neq i} \Upsilon_{ia}^3 + \sum_{i \neq j} \sigma_i \sigma_j^2 \frac{(t-1)^2 (x_i^{-1} + 2x_j^{-1})}{(t - x_i x_j^{-1})(t - x_j x_i^{-1})} \prod_{a \neq i} \Upsilon_{ia} \prod_{b \neq j} \Upsilon_{jb}^2 \\ + \sum_{i \neq j \neq k} \sigma_i \sigma_j \sigma_k \frac{(t-1)^3 x_i^{-1}}{(t - x_i x_j^{-1})(t - x_j x_k^{-1})(t - x_k x_i^{-1})} \prod_{a \neq i} \Upsilon_{ia} \prod_{b \neq j} \Upsilon_{jb} \prod_{c \neq k} \Upsilon_{kc}. \quad (4.9c)$$

The general formula is

$$G_{m,1} = \sum_{1 \leq j_0, \dots, j_{m-1} \leq n} (\sigma_{j_0} \dots \sigma_{j_{m-1}}) x_{j_0}^{-1} \prod_{s=0}^{m-1} \frac{t-1}{t - x_{j_s} x_{j_{s+1}}^{-1}} \prod_{s=0}^{m-1} \prod_{a \neq j_s} \Upsilon_{j_s a}. \quad (4.10)$$

Finally, let us look at

$$H_{m,j} := \text{tr} Y^{j,m} = m \text{tr}(Y_{m-1} \dots Y_0)^j.$$

We have  $Y_s = Z_s - X_s^{-1}$ , so in terms of  $A, B$  we have

$$Y_s = t_s B - \text{Id}_n \quad (s = 0, \dots, m-2), \quad Y_{m-1} = A^{-1}(t_{m-1} B - \text{Id}_n).$$

After rescaling,

$$H_{m,j} = \text{tr}(A^{-1} P(B))^j, \quad \text{where } P(B) := \prod_{i \in I} (B - t_i^{-1} \text{Id}_n).$$

We see that in the limit  $t_i \rightarrow \infty$ , each  $H_{m,j}$  tends to  $G_{j,m} = \text{tr}(A^{-1} B^m)^j$ . Thus,  $\{H_{m,j}\}$  is a more general  $m$ -parametric family of integrable systems. For a given  $m$ , the simplest Hamiltonian is  $H_{m,1} = \text{tr}(A^{-1} P(B))$ , which is nothing but a general linear combination of  $G_{m,1}$  with smaller  $m$ . For example,

$$H_{2,1} = \text{tr}(A^{-1} B^2) - (t_0^{-1} + t_1^{-1}) \text{tr}(A^{-1} B) + (t_0 t_1)^{-1} \text{tr} A^{-1} \quad (4.11)$$

is a linear combination of  $G_{2,1}, G_{1,1}$  and  $G_{0,1}$ . Therefore,  $H_{m,1}$  can be written explicitly using the expressions for  $G_{l,1}$  with  $l \leq m$ .

## 5. QUANTIZATION AND FURTHER LINKS

Integrable systems closely related to those constructed above appeared recently in a different context, so below we indicate these connections.

**5.1. Twisted Macdonald–Ruijsenaars systems.** In [CE, Appendix], certain generalisations of the quantum Macdonald–Ruijsenaars system were proposed. Such generalisations depend on an integer  $\ell \geq 2$  and they exist for any root system  $R$ . Their construction given in [CE] is very implicit: at the first step, the eigenfunctions  $\psi_\ell(\lambda, z)$  for the twisted system are constructed by integrating certain products of the Gaussian and the eigenfunctions  $\psi(\lambda, z)$  of the usual Macdonald–Ruijsenaars system. By analysing the properties of  $\psi_\ell(\lambda, z)$ , the existence of a complete family of commuting quantum Hamiltonians is then deduced (see [CE, Appendix] for more details). To compare these integrable systems with the ones constructed above, let us write the corresponding quantum Hamiltonian explicitly in the case  $R = A_{n-1}$  and  $\ell = 2$ .

We consider the algebra of difference operators in  $n$  variables  $z_1, \dots, z_n$  and denote by  $T_i$  the shift operator in the  $i$ th variable, acting by  $T_i f(z_1, \dots, z_n) = f(z_1, \dots, z_i + 1, \dots, z_n)$ . It will be convenient to introduce  $q \in \mathbb{C}^\times$ , not a root of unity, and work with exponential coordinates  $x_i = q^{z_i}$ , also allowing  $x_i^{1/2} = q^{z_i/2}$ . We have  $T_i(x_i^{1/2}) = q^{1/2}x_i^{1/2}$ . In the formulas below we will also use  $t \neq 0$  as a coupling parameter of the system.

**Proposition 5.1.** *For  $R = A_{n-1}$  and  $\ell = 2$ , the Hamiltonian of the twisted Macdonald–Ruijsenaars system [CE] is given by the following difference operator  $D_{2,1}$ :*

$$D_{2,1} = \sum_{i=1}^n a_i T_i^2 + \sum_{i < j} b_{ij} T_i T_j, \quad (5.1)$$

where the coefficients  $a_i, b_{ij}$  are given by

$$a_i = \prod_{j \neq i} \frac{(1 - tx_i x_j^{-1})(1 - qtx_i x_j^{-1})}{(1 - x_i x_j^{-1})(1 - qx_i x_j^{-1})}, \quad (5.2)$$

$$b_{ij} = q^{1/2}(t-1)(t-q) \frac{(x_i^{1/2} x_j^{-1/2} + x_i^{-1/2} x_j^{1/2})}{(1 - qx_i x_j^{-1})(1 - qx_j x_i^{-1})} \prod_{l \neq i, j} \frac{(1 - tx_i x_l^{-1})(1 - tx_j x_l^{-1})}{(1 - x_i x_l^{-1})(1 - x_j x_l^{-1})}. \quad (5.3)$$

*Proof.* According to [CE, Theorem 7.1(1)], for any (reduced) root system  $R \subset V$  of a Weyl group  $W$ ,  $W$ -invariant multiplicities  $m_\alpha \in \mathbb{Z}_{\geq 0}$ , and any  $\ell \in \mathbb{N}$ , there exists a twisted BA function  $\psi_\ell(\lambda, z)$  of the form

$$\psi_\ell(\lambda, z) = q^{\langle \lambda, z \rangle / \ell} \sum_{\nu \in \mathcal{N} \cap \ell^{-1}P} \psi_\nu(\lambda) q^{\langle \nu, z \rangle}, \quad \lambda, z \in V, \quad (5.4)$$

where  $\mathcal{N}$  denotes the convex hull of the  $W$ -orbit of a vector  $\rho = \sum_{\alpha \in R_+} m_\alpha \alpha$ , and  $P$  is the weight lattice of  $R$ . The function  $\psi_\ell$  is characterised by the following properties: for each  $\alpha \in R$ ,  $j = 1, \dots, m_\alpha$  and any  $\epsilon$  with  $\epsilon^\ell = 1$  one has

$$\psi_\ell \left( \lambda, z - \frac{1}{2} j \alpha \right) = \epsilon^j \psi_\ell \left( \lambda, z + \frac{1}{2} j \alpha \right) \quad \text{for } q^{\langle \alpha, z \rangle / \ell} = \epsilon. \quad (5.5)$$

Properties (5.5) determine  $\psi_\ell$  uniquely, up to an arbitrary  $\lambda$ -dependent factor. Moreover, by [CE, Theorem 7.1(3)], the function  $\psi_\ell$  is a common eigenfunction of a family of commuting  $W$ -invariant difference operators  $D_\ell^\pi$  in the  $x$ -variable, so that  $D_\ell^\pi \psi_\ell = \mathbf{m}_\pi(\lambda) \psi_\ell$ . Here  $\pi$  is any dominant weight and  $\mathbf{m}_\pi(\lambda) = \sum_{\tau \in W\pi} q^{\langle \tau, \lambda \rangle}$  is the corresponding orbitsum.

Let us consider the case  $R = A_{n-1}$ ,  $m_\alpha = m$  and  $\ell = 2$ . In this case properties (5.5) can be rewritten as

$$\psi_\ell(\lambda, z + j e_a) = \epsilon^j \psi_\ell(\lambda, z + j e_b) \quad \text{for } q^{z_a/2} = \epsilon q^{z_b/2}, \quad (5.6)$$

which should hold for all  $j = 1, \dots, m$ , distinct  $a, b \in \{1, \dots, n\}$  and  $\epsilon = \pm 1$ .

To identify the operator  $D_{2,1}$  with one of the twisted Macdonald–Ruijsenaars Hamiltonian from [CE], it suffices to show that

$$D_{2,1} \psi_\ell(\lambda, z) = \left( \sum_{i=1}^n q^{\lambda_i} \right) \psi_\ell(\lambda, z). \quad (5.7)$$

Using the approach of [C, Section 3], this would follow once we establish that the operator  $D_{2,1}$  preserves the properties (5.6) of  $\psi_\ell$ . Note that if we multiply  $\psi_\ell$  by the function  $g(z) = q^{\langle z, z \rangle/4}$ , where  $\langle z, z \rangle = z_1^2 + \dots + z_n^2$ , then  $\widetilde{\psi}_\ell(\lambda, z) = g(z)\psi_\ell(\lambda, z)$  will satisfy the following conditions:

$$\widetilde{\psi}_\ell(\lambda, z + je_a) = \widetilde{\psi}_\ell(\lambda, z + je_b) \quad \text{for } q^{z_a/2} = \epsilon q^{z_b/2}, \quad (5.8)$$

which means that

$$\widetilde{\psi}_\ell(\lambda, z + je_a) = \widetilde{\psi}_\ell(\lambda, z + je_b) \quad \text{for } q^{z_a} = q^{z_b}. \quad (5.9)$$

Therefore, it remains to check that the operator  $\widetilde{D}_{2,1} := g \circ D_{2,1} \circ g^{-1}$  in the case  $t = q^{-m}$  preserves the properties (5.9) for all  $j = 1, \dots, m$ . Explicitly, we have

$$\widetilde{D}_{2,1} = \sum_{i=1}^n \widetilde{a}_i T_i^2 + \sum_{i < j} \widetilde{b}_{ij} T_i T_j, \quad (5.10)$$

where the coefficients  $\widetilde{a}_i, \widetilde{b}_{ij}$  are given by

$$\widetilde{a}_i = (qx_i)^{-1} \prod_{j \neq i} \frac{(1 - tx_i x_j^{-1})(1 - qtx_i x_j^{-1})}{(1 - x_i x_j^{-1})(1 - qx_i x_j^{-1})}, \quad (5.11)$$

$$\widetilde{b}_{ij} = \frac{(t-1)(t-q)(x_i^{-1} + x_j^{-1})}{(1 - qx_i x_j^{-1})(1 - qx_j x_i^{-1})} \prod_{l \neq i, j} \frac{(1 - tx_i x_l^{-1})(1 - tx_j x_l^{-1})}{(1 - x_i x_l^{-1})(1 - x_j x_l^{-1})}. \quad (5.12)$$

For this operator we have the following result.

**Lemma 5.2.** *Let  $\mathcal{Q}_m$  denote the space of functions  $f(x_1, \dots, x_n)$  holomorphic on  $(\mathbb{C}^\times)^n$  and such that for any  $j = 1, \dots, m$  and  $1 \leq a < b \leq n$  we have  $(T_a)^j f = (T_b)^j f$  for  $x_a = x_b$ . Then  $\widetilde{D}_{2,1}(\mathcal{Q}_m) \subseteq \mathcal{Q}_m$ .*

This lemma is proved analogously to [C, Proposition 2.1], using [C, Lemma 2.5]. It implies that the operator  $\widetilde{D}_{2,1}$  preserves the properties (5.9), so we are done.  $\square$

To see a link with the Hamiltonian  $G_{2,1}$  (4.9b), consider the classical limit of  $\widetilde{D}_{2,1}$ . On the quantum level we have the algebra of  $q$ -difference operators, with  $[T_i, x_j] = \delta_{ij}(q-1)x_i T_i$ . In the classical limit  $q = e^{-\hbar} \rightarrow 1$  we obtain  $2n$  commuting variables  $\bar{x}_i^{\pm 1}, \bar{T}_i^{\pm 1}$  with the Poisson bracket  $\{\bar{x}_i, \bar{T}_j\} = \delta_{ij} \bar{x}_i \bar{T}_i$ . The classical limit of  $\widetilde{D}_{2,1}$  is, therefore, the following function  $\bar{D}_{2,1}$ :

$$\bar{D}_{2,1} = \sum_{i=1}^n \bar{a}_i \bar{T}_i^2 + \sum_{i < j} \bar{b}_{ij} \bar{T}_i \bar{T}_j, \quad \bar{a}_i = \bar{x}_i^{-1} \prod_{j \neq i} \frac{(1 - t\bar{x}_i \bar{x}_j^{-1})^2}{(1 - \bar{x}_i \bar{x}_j^{-1})^2}, \quad (5.13)$$

$$\bar{b}_{ij} = \frac{(t-1)^2(\bar{x}_i^{-1} + \bar{x}_j^{-1})}{(1 - \bar{x}_i \bar{x}_j^{-1})(1 - \bar{x}_j \bar{x}_i^{-1})} \prod_{l \neq i, j} \frac{(1 - t\bar{x}_i \bar{x}_l^{-1})(1 - t\bar{x}_j \bar{x}_l^{-1})}{(1 - \bar{x}_i \bar{x}_l^{-1})(1 - \bar{x}_j \bar{x}_l^{-1})}. \quad (5.14)$$

A substitution  $\bar{x}_i = x_i, \bar{T}_i = \sigma_i$  matches it to the formula (4.9b).

**Remark 5.3.** *More generally, it follows from the results of [BEF] that for any  $\ell$ , the twisted Macdonald–Ruijsenaars system from [CE, Appendix] in type A coincides, in the classical limit, with the system defined by  $G_{\ell,j} = \text{tr}(A^{-1}B^\ell)^j$ ,  $j \in \mathbb{N}$ . See [BEF, Remark 3.26].*

**Remark 5.4.** *According to [CE, Theorem 7.3], for  $t = q^{-m}$  with  $m \in \mathbb{Z}_+$  the operator  $D_{2,1}$  is algebraically integrable. Moreover, in that case its eigenfunctions  $\psi_\ell(\lambda, z)$  are given by [CE, (7.2)], with  $\ell = 2$ . The operator  $D_{2,1}$  is bispectrally self-dual, namely, one has  $\psi_\ell(\lambda, z) = \psi_\ell(z, \lambda)$ .*

**5.2. Cyclotomic DAHA and quiver gauge theory.** In the process of writing this paper we became aware of the work of Braverman, Finkelberg and Nakajima [BFN1, BFN2], and of Kodera and Nakajima [KN], where some operators generalising the Macdonald–Ruijsenaars operators appeared in the context of quiver gauge theory. In particular, in [KN] an isomorphism is established between the quantized Coulomb branch of a  $3d \mathcal{N} = 4$  supersymmetric gauge theory and the spherical subalgebra of the Cherednik algebra for  $W = \mathbb{Z}_l \wr S_N$ . The form of this isomorphism (see [KN, Theorem 1.5]) suggested to us that there should be a relation to the integrable systems constructed in this paper. Indeed, on the classical level this can be seen as follows. Recall that for each  $m$ , the Calogero–Moser space  $\mathcal{C}_{n,q}(m)$  carries three families of Poisson-commuting functions

$E_{m,j} = \text{tr } X^{mj}$ ,  $F_{m,j} = \text{tr}(1 + XY)^j$ , and  $H_{m,j} = \text{tr } Y^{jm}$ . We have an open dense subspace of  $\mathcal{C}_{n,q}(m)$  where  $X$  is invertible and everything can be written in terms of the matrices  $A, B$  that parametrise the Calogero–Moser space  $\mathcal{C}_{n,t}^0$  for the tadpole quiver, resulting in

$$E_{m,j} = \text{tr } A^j, \quad F_{m,j} = \text{tr } B^j, \quad H_{m,j} = \text{tr}(A^{-1} \prod_{i \in I} (B - t_i^{-1} \text{Id}_n))^j. \quad (5.15)$$

Choosing the coordinates  $x, \sigma$  as in (4.8), we obtain the expressions for these Hamiltonians as in section §4.2. That coordinate system was chosen so to make  $A$  diagonal. Instead, we can choose to diagonalise  $B$ , paramtrising  $A, B$  by  $w = (w_1, \dots, w_n)$  and  $u = (u_1, \dots, u_n)$  as follows:

$$B = \text{diag}(w_1, \dots, w_n), \quad A_{ij} = u_j \frac{t^{-1} - 1}{t^{-1} - w_i w_j^{-1}} \prod_{k \neq j} \frac{1 - t^{-1} w_j w_k^{-1}}{1 - w_j w_k^{-1}}. \quad (5.16)$$

This corresponds to the canonical transformation  $\Phi : (A, B) \mapsto (B, A)$  (which changes the sign of the bracket), so we have:

$$\{w_i, w_j\} = 0, \quad \{u_i, w_j\} = \delta_{ij} u_i w_j, \quad \{u_i, u_j\} = 0.$$

To write the Hamiltonians (5.15) in these new coordinates, one needs to calculate  $A^{-1}$ . This is done with the help of the Cauchy’s determinant formula; the result is

$$(A^{-1})_{ij} = u_i^{-1} \frac{t - 1}{t - w_i w_j^{-1}} \prod_{k \neq j} \frac{1 - t w_j w_k^{-1}}{1 - w_j w_k^{-1}}.$$

As a result, we have

$$E_{m,1} = \sum_{i=1}^n \prod_{j \neq i} \frac{1 - t^{-1} w_i w_j^{-1}}{1 - w_i w_j^{-1}} u_i, \quad F_{m,j} = \sum_{i=1}^n w_i^j, \quad H_{m,1} = \sum_{i=1}^n \prod_{j \neq i} \frac{1 - t w_i w_j^{-1}}{1 - w_i w_j^{-1}} \prod_{k=0}^{m-1} (w_i - t_k^{-1}) u_i^{-1}. \quad (5.17)$$

Note that in this form the integrability of the Hamiltonians  $E_{m,1}$  and  $H_{m,1}$  becomes obvious:  $E_{m,1}$  is the Ruijsenaars–Schneider Hamiltonian, and  $H_{m,1}$  reduces to such by a canonical transformation  $w_i \mapsto w_i$ ,  $u_i \mapsto \prod_{k=0}^{m-1} (w_i - t_k^{-1}) u_i$  ( $i = 1, \dots, n$ ). By contrast, in coordinates  $x_i, \sigma_i$  as in Section §4.2, the Hamiltonian  $H_{m,1}$  looks complicated and its integrability is not obvious without knowing that change of variables that reduces it to the Ruijsenaars–Schneider system. We should emphasize that the Hamiltonian flows defined by  $H_{m,j}$  are *not* complete when viewed on  $\mathcal{C}_{n,q}^0(m) \simeq \mathcal{C}_{n,t}^0$ , so one does need a cyclic-quiver interpretation in order to get a completed phase space and integrate the flows.

Now, there is an obvious parallel between (5.17) and the generators  $E_1[1]$ ,  $F_1[1]$  and  $\sum_i w_i^j$  of the quantized Coulomb branch as in [KN, Theorem 1.5]. Therefore, one should expect the above Hamiltonians to be connected with the  $K$ -theoretic Coulomb branch of a quiver gauge theory, cf. [BFN2, A(ii) and Remark A.6]. Then the quantized  $K$ -theoretic Coulomb branch [BFN1, BFN2] should also provide quantization of these integrable Hamiltonians.

In fact, a recent paper of Braverman, Etingof and Finkelberg (with an appendix by Nakajima and Yamakawa) [BEF] greatly clarifies this connection. In particular, it introduces a cyclotomic version of the double affine Hecke algebra for  $\text{GL}_n$ , giving a construction of both the quantum and the classical versions of the integrable systems considered in the present paper. Below we indicate the relationship between our results and those from [BEF].

The Calogero–Moser space  $\mathcal{C}_{n,q}(m)$  is the same as the space  $\mathcal{M}_N^l(Z, t)$  from [BEF, Section 5] with  $l = m$  and  $N = n$ . Our parameter  $t$  is the same as in [BEF], while their parameters  $Z_i$  are related to our  $q_i$  by  $Z_{i-1}/Z_i = q_i$ . By [BEF, Theorems 5.16 & 5.17], the coordinate ring  $\mathcal{O}(\mathcal{M}_N^l(Z, t))$  is identified with the spherical subalgebra of the cyclotomic DAHA  $\text{HH}_N^l(Z, 1, t)$ . This ring admits a noncommutative deformation, a spherical subalgebra  $e_N \text{HH}_N^l(Z, q, t) e_N$ , where  $q$  is a quantization parameter. The cyclotomic DAHA  $\text{HH}_N^l(Z, q, t)$ , in its turn, is defined as a subalgebra of the usual DAHA for  $\text{GL}_N$ , see [BEF, Sec. 3.4]. By [BEF, Theorem 3.28], the algebra  $\text{HH}_N^l(Z, q, t)$  contains three commutative subalgebras generated by  $X_i$ ,  $Y_i^{\pm 1}$  and  $D_i^{(l)}$  with  $i = 1, \dots, N$ . By symmetrisation, one obtains three commutative subalgebras in the spherical subalgebra of  $\text{HH}_N^l(Z, q, t)$ , see [BEF, Section 3.6]. The elements of the spherical subalgebra can be realised as difference operators; in this way one obtains three commuting subalgebras of difference operators that quantize the Poisson-commuting families  $\{E_{l,j} \mid j \in \mathbb{N}\}$ ,  $\{F_{l,j} \mid j \in \mathbb{Z}\}$  and

$\{H_{i,j} \mid j \in \mathbb{N}\}$  from Section §4.2 (the first family consists of functions of  $x_i$  so its quantization is obvious).

Let us note that writing down these quantum integrable Hamiltonians explicitly does not seem easy. In the case of a quiver with two vertices, we have the following explicit formula for the quantization of  $H_{2,1}$  (4.11).

**Proposition 5.5.** *The quantum Hamiltonian*

$$\widetilde{H}_{2,1} = \sum_{i=1}^n \widetilde{a}_i T_i^2 + \sum_{i < j}^n \widetilde{b}_{ij} T_i T_j + \alpha \sum_{i=1}^n \prod_{k \neq i}^n \frac{1 - tx_i x_k^{-1}}{1 - x_i x_k^{-1}} x_i^{-1} T_i + \beta \sum_{i=1}^n x_i^{-1} \quad (5.18)$$

is completely integrable for any values of the parameters  $\alpha, \beta$ . Here the coefficients  $\widetilde{a}_i, \widetilde{b}_{ij}$  are given by (5.11)–(5.12).

This formula is obtained by combining Proposition 5.1, (5.10) and [BEF, Corollary 3.22, Example 3.24, Remarks 3.25 & 3.26].  $\square$

There is also a “dual” realisation of these quantum Hamiltonians, obtained by applying the Cherednik–Fourier transform, see [BEF, Section 3.6]. These dual families are quantum versions of (5.17), and they are much easier to write down. Namely, one has the same formulas as in (5.17), but with  $u_i$  replaced by the multiplicative shift  $T_i$ . For instance, a quantum version of  $H_{m,1}$  is

$$H_{m,1} = \sum_{i=1}^n \prod_{j \neq i}^n \frac{1 - tw_i w_j^{-1}}{1 - w_i w_j^{-1}} \prod_{k=0}^{m-1} (w_i - t_k^{-1}) T_i^{-1}.$$

In yet another context, similar operators appeared in a recent work by Di Francesco and Kedem [DFK]. Namely, the limit  $t_k \rightarrow \infty$  gives

$$G_{m,1} = \sum_{i=1}^n \prod_{j \neq i}^n \frac{1 - tw_i w_j^{-1}}{1 - w_i w_j^{-1}} w_i^m T_i^{-1},$$

which coincides with the generalised Macdonald operator  $\mathcal{M}_{1;m}$  in [DFK, (1.5)].

## APPENDIX A. CALCULATIONS WITH THE BRACKETS

In this section we prove Propositions 3.1, 3.2, 4.3, 4.4. We will start with the case of the cyclic quiver. Given  $m \geq 2$ , we set  $I := \mathbb{Z}/m\mathbb{Z}$ . For  $r \in \mathbb{Z}$ , we set

$$E_r := \sum_{i \in I} e_{i+r} \otimes e_i \in A \otimes A. \quad (A.1)$$

Recall that  $x := \sum_{i \in I} x_i$ ,  $y := \sum_{i \in I} y_i$ , with  $e_i x = x e_{i+1}$ ,  $e_{i+1} y = y e_i$ . The element  $e = \sum_i e_i$  commutes with  $x, y$ , so we will identify it with the identity. If one introduces a  $\mathbb{Z}$ -grading on  $k\langle x, y \rangle$  by setting  $\deg x = 1$ ,  $\deg y = -1$ , then

$$(u E_r v)^\circ = v E_{r'} u, \quad r' = -r + \deg u + \deg v, \quad (A.2)$$

for any homogeneous elements  $u, v \in k\langle x, y \rangle$ .

**Lemma A.1.** *We have:*

$$\{\{xy, xy\}\} = xy E_0 - E_0 xy - \frac{1}{2} E_0 (xy)^2 + \frac{1}{2} (xy)^2 E_0. \quad (A.3)$$

*Proof.*

$$\{\{xy, xy\}\} = \{\{xy, x\}\} y + x \{\{xy, y\}\} = -\{\{x, xy\}\}^\circ y - x \{\{y, xy\}\}^\circ. \quad (A.4)$$

Using the formulas (4.2a)–(4.2c), we have

$$\{\{x, xy\}\} = \{\{x, x\}\} y + x \{\{x, y\}\} = \frac{1}{2} (E_1 x^2 - x^2 E_1) y + x \left( E_1 + \frac{1}{2} (y x E_1 + E_1 x y - y E_1 x + x E_1 y) \right),$$

and so

$$\{\{x, xy\}\}^\circ = \frac{1}{2} (x^2 y E_0 - y E_0 x^2) + E_0 x + \frac{1}{2} (E_0 x y x + x y E_0 x - x E_0 x y + y E_0 x^2).$$



Similarly, we find that

$$\begin{aligned}\{\{y, xy\}\} &= \frac{1}{2}(xy^2E_{-1} - xE_{-1}y^2) - E_{-1}y - \frac{1}{2}(E_{-1}yxy + xyE_{-1}y - xE_{-1}y^2 + yE_{-1}xy), \\ \{\{y, xy\}\}^\circ &= \frac{1}{2}(E_0xy^2 - y^2E_0x) - yE_0 - \frac{1}{2}(yxyE_0 + yE_0xy - y^2E_0x + xyE_0y).\end{aligned}$$

Substituting these expressions into (A.4) leads, after cancellations, to the expression (A.3).  $\square$

**Lemma A.2.** *Let us adjoin to  $A$  the elements  $x_i^{-1}$ , so that  $x^{-1} = \sum_{i \in I} x_i^{-1}$ . Then for  $z = y + x^{-1}$  one has:*

$$\{\{z, z\}\} = \frac{1}{2}(z^2E_{-1} - E_{-1}z^2). \quad (\text{A.5})$$

*Proof.* This formula can be checked directly, using that  $0 = \{\{y, xx^{-1}\}\} = \{\{y, x\}\}x^{-1} + x\{\{y, x^{-1}\}\}$  and  $0 = \{\{x, xx^{-1}\}\} = \{\{x, x\}\}x^{-1} + x\{\{x, x^{-1}\}\}$ . Alternatively, we can use the idea from the proof of Theorem 2.7. Namely, one can rewrite the bivector  $P$  in terms of the generators  $x_i, z_i = y_i + x_i^{-1}$ . The resulting double brackets are almost identical (with  $y_i$  replaced by  $z_i$ , which is now thought of as  $x_i^*$ ), the only difference appears in the brackets  $\{\{a, a^*\}\}$ , see (2.16). Therefore, we have

$$\{\{x, x\}\} = \frac{1}{2}(E_1x^2 - x^2E_1), \quad \{\{z, z\}\} = \frac{1}{2}(z^2E_{-1} - E_{-1}z^2), \quad (\text{A.6a})$$

$$\{\{x, z\}\} = \frac{1}{2}(zxE_1 + E_1xz - zE_1x + xE_1z). \quad (\text{A.6b})$$

This gives the needed formula for  $\{\{z, z\}\}$ .  $\square$

Now we need the following general lemma. Let  $A$  be an associative  $k$ -algebra with a double bracket  $\{\{-, -\}\}$  and the associated ordinary bracket  $\{-, -\} = m \circ \{\{-, -\}\}$ .

**Lemma A.3.** *Given  $a \in A$ , suppose that  $\mathcal{E} \subset A \otimes A$  is a subset such that for any  $E \in \mathcal{E}$  and any  $r, s \geq 0$  we have  $(a^r E a^s)^\circ = a^s E' a^r$  for some  $E' \in \mathcal{E}$  which depends on  $r + s$  but not on  $r, s$  individually. Moreover, assume that  $a$  commutes with any element in  $m(\mathcal{E}) \subset A$ . If  $\{\{a, a\}\} = \sum_i (a^i E_i - E_i a^i)$  with  $E_i \in \mathcal{E}$ , then  $\{a^k, a^l\} = 0$  for all  $k, l$ .*

*Proof.* Since  $\{-, -\}$  satisfies Leibniz's rule in the second argument, it is enough to prove that  $\{a^k, a\} = 0$  for all  $a$ . Now,  $\{\{a^k, a\}\} = -\{\{a, a^k\}\}^\circ$ , while

$$\{\{a, a^k\}\} = \sum_{\substack{r, s \geq 0 \\ r+s=k-1}} a^r \{\{a, a\}\} a^s = \sum_{\substack{r, s \geq 0 \\ r+s=k-1}} \sum_i a^r (a^i E_i - E_i a^i) a^s,$$

for some  $E_i \in \mathcal{E}$ . Using that  $(a^{r+i} E_i a^s - a^r E_i a^{s+i})^\circ = a^s E'_i a^{r+i} - a^{s+i} E'_i a^r$ , for some  $E'_i \in \mathcal{E}$ , we obtain:

$$m(\{\{a^k, a\}\}) = - \sum_{\substack{r, s \geq 0 \\ r+s=k-1}} \sum_i m(a^s E'_i a^{r+i} - a^{s+i} E'_i a^r) = \sum_{r, s} \sum_i (a^s m(E'_i) a^{r+i} - a^{s+i} m(E'_i) a^r) = 0,$$

since  $m(E'_i)$  commutes with  $a$ . Therefore,  $\{a^k, a\} = 0$ , as needed.  $\square$

*Proof of Proposition 4.3.* We now use this lemma with  $\mathcal{E} = \bigoplus_{r \in I} \mathbb{C}E_r$  with  $E_r$  as in (A.1), and with  $a = x, y, xy$  or  $z = y + x^{-1}$ . We have  $m(\mathcal{E}) = \mathbb{C}e$ , where  $e := \sum_i e_i$  commutes with any of the above  $a$ . The assumptions of the lemma are satisfied due to (A.2). The rest follows from (4.2a), (A.3) and (A.5).  $\square$

*Proof of Proposition 4.4.* We start by calculating some double brackets. First,

$$\begin{aligned}
\{\{x, x^a\}\} &= \sum_{\substack{r, s \geq 0 \\ r+s=a-1}} x^r \{\{x, x\}\} x^s = \sum_{\substack{r, s \geq 0 \\ r+s=a-1}} \left( \frac{1}{2} x^r E_1 x^{s+2} - \frac{1}{2} x^{r+2} E_1 x^s \right), \\
\{\{x^a, x\}\} &= -\{\{x, x^a\}\}^\circ = \sum_{\substack{r, s \geq 0 \\ r+s=a-1}} \left( -\frac{1}{2} x^{s+2} E_a x^r + \frac{1}{2} x^s E_a x^{r+2} \right), \\
\{\{x^a, x^b\}\} &= \sum_{\substack{r, s \geq 0 \\ r+s=b-1}} x^r \{\{x^a, x\}\} x^s \\
&= \sum_{\substack{r, s \geq 0 \\ r+s=b-1}} \sum_{\substack{r', s' \geq 0 \\ r'+s'=a-1}} \left( -\frac{1}{2} x^{r+s'+2} E_a x^{r'+s} + \frac{1}{2} x^{r+s'} E_a x^{r'+s+2} \right), \\
&= \sum_{\substack{r, s \geq 0 \\ r+s=b-1}} \sum_{\substack{r', s' \geq 0 \\ r'+s'=a-1}} \left( -\frac{1}{2} x^{r+s'+2} E_a x^{r'+s} + \frac{1}{2} x^{r'+s} E_a x^{r+s'+2} \right).
\end{aligned}$$

Next, we have

$$\begin{aligned}
\{\{y, x^a\}\} &= \sum_{\substack{r, s \geq 0 \\ r+s=a-1}} x^r \{\{y, x\}\} x^s \\
&= \sum_{\substack{r, s \geq 0 \\ r+s=a-1}} \left( -x^r E_{-1} x^s - \frac{1}{2} x^r E_{-1} y x^{s+1} - \frac{1}{2} x^{r+1} y E_{-1} x^s + \frac{1}{2} x^{r+1} E_{-1} y x^s - \frac{1}{2} x^r y E_{-1} x^{s+1} \right).
\end{aligned}$$

For  $\{\{x^a, y\}\} = -\{\{y, x^a\}\}^\circ$  we, therefore, obtain:

$$\{\{x^a, y\}\} = \sum_{\substack{r, s \geq 0 \\ r+s=a-1}} \left( x^s E_a x^r + \frac{1}{2} y x^{s+1} E_a x^r + \frac{1}{2} x^s E_a x^{r+1} y - \frac{1}{2} y x^s E_a x^{r+1} + \frac{1}{2} x^{s+1} E_a x^r y \right).$$

From this,

$$\begin{aligned}
\{\{x^a, yx^b\}\} &= \{\{x^a, y\}\} x^b + y \{\{x^a, x^b\}\} \\
&= \sum_{\substack{r, s \geq 0 \\ r+s=a-1}} \left( x^s E_a x^{r+b} + \frac{1}{2} y x^{s+1} E_a x^{r+b} + \frac{1}{2} x^s E_a x^{r+1} y x^b - \frac{1}{2} y x^s E_a x^{r+1+b} + \frac{1}{2} x^{s+1} E_a x^r y x^b \right) \\
&+ \sum_{\substack{r, s \geq 0 \\ r+s=b-1}} \sum_{\substack{r', s' \geq 0 \\ r'+s'=a-1}} \left( -\frac{1}{2} y x^{r+s'+2} E_a x^{r'+s} + \frac{1}{2} y x^{r'+s} E_a x^{r+s'+2} \right).
\end{aligned}$$

As a result, for  $\{\{yx^b, x^a\}\} = -\{\{x^a, yx^b\}\}^\circ$  we get:

$$\begin{aligned}
\{\{yx^b, x^a\}\} &= \sum_{\substack{r, s \geq 0 \\ r+s=a-1}} \left( -x^{r+b} E_{b-1} x^s - \frac{1}{2} x^{r+b} E_{b-1} y x^{s+1} - \frac{1}{2} x^{r+1} y x^b E_{b-1} x^s + \frac{1}{2} x^{r+1+b} E_{b-1} y x^s - \frac{1}{2} x^r y x^b E_{b-1} x^{s+1} \right) \\
&+ \sum_{\substack{r, s \geq 0 \\ r+s=b-1}} \sum_{\substack{r', s' \geq 0 \\ r'+s'=a-1}} \left( \frac{1}{2} x^{r'+s} E_{b-1} y x^{r+s'+2} - \frac{1}{2} x^{r+s'+2} E_{b-1} y x^{r'+s} \right).
\end{aligned}$$

Now we calculate

$$\begin{aligned}
\{\{y, yx^a\}\} &= \{\{y, y\}\} x^a + y \{\{y, x^a\}\} \\
&= \frac{1}{2} y^2 E_{-1} x^a - \frac{1}{2} E_{-1} y^2 x^a \\
&+ \sum_{\substack{r, s \geq 0 \\ r+s=a-1}} \left( -y x^r E_{-1} x^s - \frac{1}{2} y x^r E_{-1} y x^{s+1} - \frac{1}{2} y x^{r+1} y E_{-1} x^s + \frac{1}{2} y x^{r+1} E_{-1} y x^s - \frac{1}{2} y x^r y E_{-1} x^{s+1} \right).
\end{aligned}$$

Therefore, for  $\{\{yx^a, y\} = -\{yx^a, y\}^\circ$  we obtain:

$$\begin{aligned} \{\{yx^a, y\} &= -\frac{1}{2}x^a E_{a-1}y^2 + \frac{1}{2}y^2 x^a E_{a-1} \\ &+ \sum_{\substack{r,s \geq 0 \\ r+s=a-1}} (x^s E_{a-1}yx^r + \frac{1}{2}yx^{s+1}E_{a-1}yx^r + \frac{1}{2}x^s E_{a-1}yx^{r+1}y - \frac{1}{2}yx^s E_{a-1}yx^{r+1} + \frac{1}{2}x^{s+1}E_{a-1}yx^r y). \end{aligned}$$

All this allows us to find the ordinary brackets  $\{u, v\} = \{\{u, v\}' \{u, v\}''$ . First, we have for  $a = 0 \pmod m$ :

$$\{x^a, y\} = \sum_{\substack{r,s \geq 0 \\ r+s=a-1}} (x^{s+r} + x^{s+1+r}y) = ax^{a-1} + ax^a y, \quad (\text{A.7})$$

from which it follows that

$$\{x^a, yx^b\} = \{x^a, y\}x^b = ax^{a+b-1} + ax^a yx^b = ax^{a+b-1} + ayx^{a+b} \pmod [A, A],$$

which is the first relation in Proposition 4.4.

Next, for  $b = 1 \pmod m$

$$\begin{aligned} \{yx^b, x^a\} &= \sum_{\substack{r,s \geq 0 \\ r+s=a-1}} \left( -x^{r+b+s} - \frac{1}{2}x^{r+b}yx^{s+1} - \frac{1}{2}x^{r+1}yx^{b+s} + \frac{1}{2}x^{r+1+b}yx^s - \frac{1}{2}x^r yx^{b+s+1} \right) \\ &+ \sum_{\substack{r,s \geq 0 \\ r+s=b-1}} \sum_{\substack{r',s' \geq 0 \\ r'+s'=a-1}} \left( \frac{1}{2}x^{r'+s}yx^{r+s'+2} - \frac{1}{2}x^{r'+s'+2}yx^{r'+s} \right), \end{aligned}$$

from which we get that

$$\begin{aligned} y\{yx^b, x^a\} &= \sum_{\substack{r,s \geq 0 \\ r+s=a-1}} \left( -yx^{r+b+s} - \frac{1}{2}yx^{r+b}yx^{s+1} - \frac{1}{2}yx^{s+1}yx^{b+r} + \frac{1}{2}yx^{r+1+b}yx^s - \frac{1}{2}yx^s yx^{b+r+1} \right) \\ &+ \sum_{\substack{r,s \geq 0 \\ r+s=b-1}} \sum_{\substack{r',s' \geq 0 \\ r'+s'=a-1}} \frac{1}{2}[yx^{r'+s}, yx^{r'+s'+2}], \end{aligned}$$

which, modulo commutators, gives

$$y\{yx^b, x^a\} = -ayx^{a+b-1} - \sum_{\substack{r,s \geq 0 \\ r+s=a-1}} yx^{s+1}yx^{r+b} \pmod [A, A]. \quad (\text{A.8})$$

Finally, for  $a = 1 \pmod m$ :

$$\begin{aligned} \{yx^a, y\} &= -\frac{1}{2}x^a y^2 + \frac{1}{2}y^2 x^a \\ &+ \sum_{\substack{r,s \geq 0 \\ r+s=a-1}} \left( x^s yx^r + \frac{1}{2}yx^{s+1}yx^r + \frac{1}{2}x^s yx^{r+1}y - \frac{1}{2}yx^s yx^{r+1} + \frac{1}{2}x^{s+1}yx^r y \right), \end{aligned}$$

and therefore

$$\begin{aligned} \{yx^a, y\}x^b &= \frac{1}{2}[y^2 x^b, x^a] \\ &+ \sum_{\substack{r,s \geq 0 \\ r+s=a-1}} \left( x^s yx^{r+b} + \frac{1}{2}yx^{s+1}yx^{r+b} + \frac{1}{2}x^s yx^{r+1}yx^b - \frac{1}{2}yx^s yx^{r+1+b} + \frac{1}{2}x^{s+1}yx^r yx^b \right). \end{aligned}$$

Modulo commutators, this gives

$$\sum_{\substack{r,s \geq 0 \\ r+s=a-1}} \left( yx^{r+s+b} + \frac{1}{2}yx^{s+1}yx^{r+b} + \frac{1}{2}yx^{r+1}yx^{s+b} - \frac{1}{2}yx^s yx^{r+1+b} + \frac{1}{2}yx^r yx^{s+1+b} \right).$$

Therefore,

$$\{yx^a, y\}x^b = ayx^{a+b-1} + \sum_{\substack{r,s \geq 0 \\ r+s=a-1}} yx^{s+1}yx^{r+b} \pmod [A, A]. \quad (\text{A.9})$$

Now, using (A.8), (A.9), we obtain modulo commutators:

$$\{yx^b, yx^c\} = \{yx^b, y\}x^c + y\{yx^b, x^c\} = (b-c)yx^{b+c+1} + \sum_{\substack{r,s \geq 0 \\ r+s=b-1}} yx^{s+1}yx^{r+c} - \sum_{\substack{r,s \geq 0 \\ r+s=c-1}} yx^{s+1}yx^{r+b},$$

which gives (4.3b). This finishes the proof of Proposition 4.4.  $\square$

For the tadpole quiver, the brackets in (3.1a)–(3.1e) look entirely similar (with small sign differences), and all the above proofs carry over with  $E_r = E_0 = e_0 \otimes e_0$  for all  $r$ , leading to

**Proposition A.4.** *For any  $a, b \geq 0$  we have*

$$\{x^a, x^b\} = \{y^a, y^b\} = \{(xy)^a, (xy)^b\} = \{z^a, z^b\} = 0, \quad \text{where } z = y + x^{-1}, \quad (\text{A.10a})$$

$$\{x^a, yx^b\} = ax^{a+b-1} + ayx^{a+b} \pmod{[A, A]}, \quad (\text{A.10b})$$

$$\{yx^a, yx^b\} = (a-b)yx^{a+b-1} + \sum_{t=1}^a yx^t yx^{a+b-t} - \sum_{t=1}^b yx^t yx^{a+b-t} \pmod{[A, A]}. \quad (\text{A.10c})$$

Finally, replacing  $y$  by  $z - x^{-1}$  in the last two formulas, one gets after a simple rearrangement the formulas from Propositions 3.2.  $\square$

## APPENDIX B. HAMILTONIAN DYNAMICS

In this section we prove Propositions 4.7. We start with a simple formula, immediate from (2.2): for any  $a, b \in A$ ,

$$\{\text{tr } a, b_{ij}\} = \{a, b\}_{ij}.$$

It is convenient to rewrite this relation in matrix form, using the notation  $\mathcal{X}(a) = (a_{ij})$  for the matrix-valued function on the representation space, associated to  $a \in A$ :

$$\{\text{tr } \mathcal{X}(a), \mathcal{X}(b)\} = \mathcal{X}(\{a, b\}). \quad (\text{B.1})$$

Let now  $\bar{Q}$  be the doubled framed quiver as in Section §4, with the double bracket defined by the bivector (2.5), and  $\{-, -\}$  be the associated bracket  $\{-, -\} = m \circ \{\{-, -\}\}$ . Recall that  $\{y^k, y\} = 0$  for all  $k$ .

**Lemma B.1.** *For any  $k \in m\mathbb{N}$ ,*

$$\{y^k, v\} = \{y^k, w\} = 0, \quad \{y^k, x\} = -ky^{k-1} - ky^k x.$$

*Proof.* From (2.6c) we have

$$\begin{aligned} \{\{y_i, v\}\} &= -\frac{1}{2}(y_{m-1} \otimes v)\delta_{i,m-1} + \frac{1}{2}(e_0 \otimes y_0 v)\delta_{i,0}, \\ \{\{y_i, w\}\} &= \frac{1}{2}(wy_{m-1} \otimes e_0)\delta_{i,m-1} - \frac{1}{2}(w \otimes y_0)\delta_{i,0}. \end{aligned}$$

Using this, one performs a calculation as in Appendix §A to find that

$$\{\{v, y^k\}\} = \sum_{r+s=k-1} y^r \{\{v, y\}\} y^s = \sum_{r+s=k-1} \left( \frac{1}{2}y^r v \otimes e_0 y^{s+1} - \frac{1}{2}y^{r+1} v \otimes e_0 y^s \right),$$

and  $\{y^k, v\} = 0$  as a result. The relation  $\{y^k, w\} = 0$  is checked similarly. The last formula in the lemma is analogous to (A.7).  $\square$

Now consider a representation space  $\text{Rep}(\mathbb{C}\bar{Q}, \tilde{\alpha})$  for a dimension vector  $\tilde{\alpha} = (1, \alpha) \in \mathbb{N}^{m+1}$ , and let as before  $V, W, X, Y$  be the matrices representing the arrows  $v, w$  and  $x = \sum_i x_i, y = \sum_i y_i$ . The space  $\text{Rep}(\mathbb{C}\bar{Q}, \tilde{\alpha})$  is equipped with an anti-symmetric biderivation  $\{-, -\}$  defined by (2.2). We put  $H_k = \frac{1}{k} \text{tr } Y^k$ , where  $Y$  is the matrix representing  $y = \sum_{i \in I} y_i$  and  $k \in m\mathbb{N}$ . Then  $\{H_k, -\}$  defines a derivation (vector field) on the representation space. Since  $\{H_k, H_l\} = 0$  for all  $k, l$  and  $\{-, -\}$  is a Loday bracket (see (3.3)), the vector fields associated to  $H_k$  with different  $k$  pairwise commute. Let  $\frac{d}{dt_j}$  denote the vector field corresponding to  $H_{j_m}$ . Using the above lemma and formula (B.1), we obtain:

$$\frac{d}{dt_j} V = \frac{d}{dt_j} W = 0, \quad \frac{d}{dt_j} Y = 0, \quad \frac{d}{dt_j} X = -Y^{j_m-1} - Y^{j_m} X.$$

Integrating the last equation with constant  $Y = Y(0)$ , we find that

$$X(t_j) = e^{-t_j Y^{j^m}} X(0) + Y^{-1}(e^{-t_j Y^{j^m}} - 1).$$

This formula is well-defined for all  $Y$  because the function  $z^{-1}(e^{-t_k z^k} - 1)$  is analytic in  $z$ . By superposition, for  $\mathbf{t} = (t_1, t_2, \dots)$  we have:

$$X(\mathbf{t}) = e^{-\sum_{j \geq 1} t_j Y^{j^m}} X(0) + Y^{-1} \left( e^{-\sum_{j \geq 1} t_j Y^{j^m}} - 1 \right).$$

Note that  $1 + YX = e^{-\sum_{j \geq 1} t_j Y^{j^m}} (1 + YX(0))$ , so the matrices  $\text{Id}_n + Y_s X_s$ ,  $s \in I$  remain invertible for all times.

If instead of  $H_k$  one considers  $G_k := \frac{1}{k} \text{tr} Z^k$ ,  $Z = Y + X^{-1}$ , then the brackets look the same, apart from the bracket between  $z$  and  $x$  (A.6b). As a result, if  $t_j$  denotes the time flow for  $G_{j^m}$ , we obtain:

$$\frac{d}{dt_j} V = \frac{d}{dt_j} W = 0, \quad \frac{d}{dt_j} Z = 0, \quad \frac{d}{dt_j} X = -Z^{j^m} X.$$

This leads to

$$X(\mathbf{t}) = e^{-\sum_{j \geq 1} t_j Z^{j^m}} X(0), \quad Z = Z(0).$$

In both cases, it is easy to see that the dynamics preserves the moment map equations. Therefore, the associated Hamiltonian flows are complete.  $\square$

### APPENDIX C. POISSON ISOMORPHISM BETWEEN $\mathcal{C}_{n,t}^0$ AND $\mathcal{C}_{n,q}^0(m)$

In this section we prove Proposition 4.10. The map  $\xi : \mathcal{C}_{n,t}^0 \rightarrow \mathcal{C}_{n,q}^0(m)$  is described in Proposition 4.9, namely:

$$X_i = \text{Id}_n, \quad Z_i = t_i B \quad (i = 0 \dots m-2), \quad X_{m-1} = A, \quad Z_{m-1} = t_{m-1} A^{-1} B.$$

We need to check that  $\xi^*\{f, g\} = \{\xi^*f, \xi^*g\}$  for any two functions on  $\mathcal{C}_{n,q}^0(m)$ . Consider the functions  $f_\alpha := \text{tr}(X^{\alpha m})$  and  $g_\beta := \text{tr}(ZX^{1+\beta m})$ . Expressing them in terms of  $A, B$  we obtain:

$$\xi^* f_\alpha = \sum_{i \in I} \text{tr} X_{i+1} \dots X_{i+\alpha m} = m \text{tr} A^\alpha, \quad (\text{C.1a})$$

$$\xi^* g_\beta = \sum_{i \in I} \text{tr} Z_i X_i X_{i+1} \dots X_{i+\beta m} = \tau \text{tr} B A^\beta, \quad \text{where } \tau := \sum_{i \in I} t_i. \quad (\text{C.1b})$$

This implies that  $f_1, \dots, f_n, g_1, \dots, g_n$  can be used as local coordinates near a generic point of  $\mathcal{C}_{n,q}^0(m)$ . Therefore, it is sufficient to check that the brackets behave well just for these functions.

Let us rewrite the formulas from Proposition 4.4 in the algebra  $A'$  with inverted  $x$ . We use  $z := y + x^{-1}$  so that  $zx = 1 + yx$ , and rearrange (4.3a)–(4.3b) as follows:

$$\{x^a, zx^b\} = azx^{a+b} \quad \text{mod } [A', A'], \quad (\text{C.2a})$$

$$\{zx^b, zx^c\} = \sum_{t=1}^b zx^t zx^{b+c-t} - \sum_{t=1}^c zx^t zx^{b+c-t} \quad \text{mod } [A', A']. \quad (\text{C.2b})$$

Recall that we also have  $\{x^a, x^b\} = 0$  for all  $a, b$ . Substituting  $a = \alpha m$ ,  $b = 1 + \beta m$ ,  $c = 1 + \gamma m$  and taking traces using (2.10) leads to:

$$\{f_\alpha, f_\beta\} = 0, \quad \{f_\alpha, g_\beta\} = \alpha m g_{\alpha+\beta}, \quad (\text{C.3a})$$

$$\{g_\beta, g_\gamma\} = \sum_{r=0}^{\beta m} h_{r, (\beta+\gamma)m-r} - \sum_{r=0}^{\gamma m} h_{r, (\beta+\gamma)m-r}. \quad (\text{C.3b})$$

Here we used the notation  $h_{r,s} := \text{tr} ZX^{1+r} ZX^{1+s}$ . Using that  $h_{r,s} = h_{s,r}$  and assuming  $\beta < \gamma$ , the last relation can be rearranged to

$$\{g_\beta, g_\gamma\} = \sum_{r=\beta m}^{\gamma m-1} h_{r, (\beta+\gamma)m-r}. \quad (\text{C.4})$$

On the other hand, the Poisson bracket on  $\mathcal{C}_{n,t}^0$  satisfies (3.10), which in terms of  $A, B$  reads:

$$\{\mathrm{tr} A^a, \mathrm{tr} A^b\} = 0, \quad \{\mathrm{tr} A^a, \mathrm{tr} BA^b\} = a \mathrm{tr} BA^{a+b}, \quad (\text{C.5a})$$

$$\{\mathrm{tr} BA^b, \mathrm{tr} BA^c\} = \sum_{r=0}^b \mathrm{tr} BA^r BA^{b+c-r} - \sum_{r=0}^c \mathrm{tr} BA^r BA^{b+c-r} = \sum_{r=b}^{c-1} \mathrm{tr} BA^r BA^{b+c-r}. \quad (\text{C.5b})$$

It remains to check that these formulas agree with those obtained from (C.3a) and (C.4) by replacing  $f_\alpha, g_\beta$  with their pull-backs  $\xi^* f_\alpha = m \mathrm{tr} A^\alpha$  and  $\xi^* g_\beta = \tau \mathrm{tr} BA^\beta$ . For the first two relations this is obvious. For the third relation, we need to show that for  $\beta < \gamma$

$$\sum_{r=\beta m}^{\gamma m-1} \xi^* h_{r,(\beta+\gamma)m-r} = \tau^2 \sum_{p=\beta}^{\gamma-1} \mathrm{tr} BA^p BA^{\beta+\gamma-p}. \quad (\text{C.6})$$

We have  $\xi^* h_{r,s} = \mathrm{tr} XZX^r XZX^s$ , with

$$XZX^r XZX^s = \sum_{i \in I} X_i Z_i X_i \dots X_{i+r-1} X_{i+r} Z_{i+r} X_{i+r} \dots X_{i+r+s-1}.$$

By expressing this in terms of  $A, B$  and taking traces we obtain:

$$\xi^* h_{r,s} = \sum_{i \in I} t_i t_{i+r} \mathrm{tr} BA^{\phi(i,i+r)} BA^{\phi(i+r,i+r+s)},$$

where  $\phi(i,j) = \left[ \frac{j}{m} \right] - \left[ \frac{i}{m} \right]$  counts the number of integers between  $i$  and  $j-1$  congruent to  $-1$  modulo  $m$ . Here and below the indices in  $t_i$  are always treated modulo  $m$ , i.e.,  $t_{i+m} = t_i$ . Therefore, if  $r = j + pm$  then

$$\xi^* h_{r,(\beta+\gamma)m-r} = \sum_{i \in I} t_i t_{i+j} \mathrm{tr} BA^{p+\phi(i,i+j)} BA^{\beta+\gamma-p-\phi(i,i+j)}.$$

Note that for  $i, j \in I$ ,  $\phi(i,i+j)$  equals 0 or 1, depending on whether  $i+j \leq m-1$  or not. We can use this in the l.h.s. of (C.6), replacing the summation over  $r$  with the summation over  $p \in [\beta, \gamma-1]$  and  $j \in I$ , which leads to

$$\sum_{\substack{i,j \in I \\ i+j \leq m-1}} t_i t_{i+j} \sum_{p=\beta}^{\gamma-1} \mathrm{tr} BA^p BA^{\beta+\gamma-p} + \sum_{\substack{i,j \in I \\ i+j > m-1}} t_i t_{i+j} \sum_{p=\beta}^{\gamma-1} \mathrm{tr} BA^{p+1} BA^{\beta+\gamma-p-1}.$$

It is easy to see that  $\sum_{p=\beta}^{\gamma-1} \mathrm{tr} BA^{p+1} BA^{\beta+\gamma-p-1} = \sum_{p=\beta}^{\gamma-1} \mathrm{tr} BA^p BA^{\beta+\gamma-p}$ , therefore, we arrive at

$$\sum_{i,j \in I} t_i t_{i+j} \sum_{p=\beta}^{\gamma-1} \mathrm{tr} BA^p BA^{\beta+\gamma-p}.$$

It remains to notice that  $\sum_{i,j \in I} t_i t_{i+j} = (\sum_{i \in I} t_i)^2 = \tau^2$ , so the obtained expression coincides with the r.h.s. of the relation (C.6). This finishes the proof of (C.6) and of the Proposition 4.10.  $\square$

## REFERENCES

- [AKSM] Alekseev, A., Kosmann-Schwarzbach, Y., Meinrenken, E.: *Quasi-Poisson manifolds*. Canad. J. Math. 54(1), 3–29 (2002); [arXiv:math/0006168 \[math.DG\]](#).
- [AMM] Alekseev, A., Malkin, A., Meinrenken, E.: *Lie group valued moment maps*. J. Differential Geom. 48, 445–495 (1998); [arXiv:dg-ga/9707021](#).
- [BC] Berest, Yu., Chalykh, O.:  *$A_\infty$ -modules and Calogero-Moser spaces*. J. Reine Angew. Math. 607, 69–112 (2007); [arXiv:math/0410194 \[math.QA\]](#).
- [BCE] Berest, Yu., Chalykh, O., Eshmatov, F.: *Recollement of deformed preprojective algebras and the Calogero-Moser correspondence*. Mosc. Math. J. 8(1), 21–37 (2008); [arXiv:0706.3006 \[math.RT\]](#).
- [BRT] Berest, Yu., Ramadoss, A., Tang, X.: *The Picard group of a noncommutative algebraic torus*. J. Noncommut. Geom. 7(2), 335–356 (2013); [arXiv:1010.3779 \[math.QA\]](#).
- [BW1] Berest, Yu., Wilson, G.: *Automorphisms and ideals of the Weyl algebra*. Math. Ann. 318(1), 127–147 (2000); [arXiv:math/0102190 \[math.QA\]](#).
- [BW2] Berest, Yu., Wilson, G.: *Ideal classes of the Weyl algebra and noncommutative projective geometry (With an appendix by Michel Van den Bergh)*. Int. Math. Res. Not. 26, 1347–1396 (2002); [arXiv:math/0104248 \[math.AG\]](#).
- [BEF] Braverman, A., Etingof, P., Finkelberg, M.: *Cyclotomic double affine Hecke algebras (with an appendix by H. Nakajima and D. Yamakawa)*; [arXiv:1611.10216 \[math.RT\]](#).



- [BF] Baker, T., Forrester, P.: *A  $q$ -analogue of the type A Dunkl operator and integral kernel*, Int. Math. Res. Not. 14, 667–686 (1997); [arXiv:q-alg/9701039](#).
- [BGK1] Baranovsky, V., Ginzburg, V., Kuznetsov, A.: *Quiver varieties and a noncommutative  $\mathbb{P}^2$* . Compositio Math. 134 (3), 283–318 (2002); [arXiv:math/0103068 \[math.AG\]](#).
- [BGK2] Baranovsky, V., Ginzburg, V., Kuznetsov, A.: *Wilson’s Grassmannian and a noncommutative quadric*. Int. Math. Res. Not. 21, 1155–1197 (2003); [arXiv:math/0203116 \[math.AG\]](#).
- [BP] Bielawski, R., Pidstrygach, V.: *On the symplectic structure of instanton moduli spaces*. Adv. Math. 226 (3), 2796–2824 (2011); [arXiv:0812.4918 \[math.SG\]](#).
- [BFN1] Braverman, A., Finkelberg, M., Nakajima, H.: *Towards a mathematical definition of Coulomb branches of 3-dimensional  $\mathcal{N} = 4$  gauge theories, II*; [arXiv:1601.03586 \[math.RT\]](#).
- [BFN2] Braverman, A., Finkelberg, M., Nakajima, H.: *Coulomb branches of  $3d\mathcal{N} = 4$  quiver gauge theories and slices in the affine Grassmannian (with appendices by Alexander Braverman, Michael Finkelberg, Joel Kamnitzer, Ryosuke Kodera, Hiraku Nakajima, Ben Webster, and Alex Weekes)*; [arXiv:1604.03625 \[math.RT\]](#).
- [CF] Calogero, F., Françoise, J.-P.: *Solution of Certain Integrable Dynamical Systems of Ruijsenaars-Schneider Type with Completely Periodic Trajectories*. Ann. Henri Poincaré 1 (1), 173–191 (2000).
- [C] Chalykh, O.: *Macdonald polynomials and algebraic integrability*. Adv. Math. 166(2), 193–259 (2002); [arXiv:math/0212313 \[math.QA\]](#).
- [CE] Chalykh, O., Etingof, P.: *Orthogonality relations and Cherednik identities for multivariable Baker-Akhiezer functions, with an appendix by O. Chalykh*. Adv. Math. 238, 246–289 (2013); [arXiv:1111.0515 \[math.QA\]](#).
- [CN] Chalykh, O., Nijhoff, F.: *Bispectral rings of difference operators*. Russian Math. Surveys 54(3), 644–645 (1999).
- [CS] Chalykh, O., Silantyev, A.: *KP hierarchy for the cyclic quiver*. J. Math. Phys. 58, 071702 (2017); [arXiv:1512.08551 \[math.QA\]](#).
- [CB] Crawley-Boevey, W.: *Preprojective algebras, differential operators and a Conze embedding for deformations of Kleinian singularities*. Comment. Math. Helv. 74(4), 548–574 (1999).
- [CBS] Crawley-Boevey, W., Shaw, P.: *Multiplicative preprojective algebras, middle convolution and the Deligne-Simpson problem*. Adv. Math. 201, 180–208 (2006); [arXiv:math/0404186 \[math.RA\]](#).
- [DFK] Di Francesco, P., Kedem, R.:  *$(t, q)$ -deformed  $Q$ -systems, DAHA and quantum toroidal algebras via generalized Macdonald operators*; [arXiv:1704.00154 \[math-ph\]](#).
- [Esh] Eshmatov, F.: *DG-models of projective modules and Nakajima quiver varieties*. Homology, Homotopy Appl. 9(2), 177–208 (2007); [arXiv:math/0604011 \[math.QA\]](#).
- [Et] Etingof, P.: *Calogero-Moser systems and representation theory*. Zurich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zürich, x+92 (2007).
- [FK1] Fehér, L., Klimčík, C.: *On the duality between the hyperbolic Sutherland and the rational Ruijsenaars-Schneider models*. J. Phys. A: Math. Theor. 42, 185–202 (2009); [arXiv:0901.1983 \[math-ph\]](#).
- [FK2] Fehér, L., Klimčík, C.: *Poisson-Lie generalization of the Kazhdan-Kostant-Sternberg reduction*. Lett. Math. Phys. 87, 125–138 (2009); [arXiv:0809.1509 \[math-ph\]](#).
- [FK3] Fehér, L., Klimčík, C.: *Self-duality of the compactified Ruijsenaars-Schneider system from quasi-Hamiltonian reduction*. Nuclear Phys. B 860(3), 464–515 (2012); [arXiv:1101.1759 \[math-ph\]](#).
- [FM] Fehér, L., Marshall, I.: *The action-angle dual of an integrable Hamiltonian system of Ruijsenaars-Schneider-van Diejen type*. [arXiv:1702.06514 \[math-ph\]](#).
- [FG] Fehér, L., Görbe, T.F.: *The full phase space of a model in the Calogero-Ruijsenaars family*. J. Geom. Phys. 115, 139–149 (2017); [arXiv: 1603.02877 \[math-ph\]](#).
- [FGNR] Fock, V., Gorsky, A., Nekrasov, N., Rubtsov, V.: *Duality in integrable systems and gauge theories*. JHEP 07, 028 (2000); [arXiv:hep-th/9906235](#).
- [FR] Fock, V., Rosly, A.: *Poisson structure on moduli of flat connections on Riemann surfaces and the  $r$ -matrix*. In: Moscow Seminar in Mathematical Physics, volume 191 of Amer. Math. Soc. Transl. Ser. 2, 67–86. Amer. Math. Soc., Providence, RI (1999); [arXiv:math/9802054 \[math.QA\]](#).
- [GN] Gorsky, A., Nekrasov, N.: *Relativistic Calogero-Moser model as gauged WZW theory*. Nucl. Phys. B 436, 582–608 (1995); [arXiv:hep-th/9401017](#).
- [I] Iliev, P.:  *$q$ -KP hierarchy, bispectrality and Calogero-Moser systems*. J. Geom. Phys. 35(2-3), 157–182 (2000).
- [KKO] Kapustin, A., Kuznetsov, A., Orlov, D.: *Noncommutative instantons and twistor transform*. Comm. Math. Phys. 221 (2), 385–432 (2001); [arXiv:hep-th/0002193](#).
- [KKS] Kazhdan, D., Kostant, B., Sternberg, S.: *Hamiltonian group actions and dynamical systems of Calogero type*. Comm. Pure Appl. Math. 31(4), 481–507 (1978).
- [KN] R. Kodera and H. Nakajima, *Quantized Coulomb branches of Jordan quiver gauge theories and cyclotomic rational Cherednik algebras*; [arXiv:1608.00875 \[math.RT\]](#).
- [Kr1] Krichever, I.M.: *Elliptic Solutions to Difference Nonlinear Equations and Nested Bethe Ansatz Equations*. In: Calogero-Moser-Sutherland Models (ed. by J.F. van Diejen, L. Vinet), CRM Series in Mathematical Physics, 249–271. Springer New York (2000); [arXiv:solv-int/9804016](#).
- [Kr2] Krichever, I.M.: *Vector bundles and Lax equations on algebraic curves*. Comm. Math. Phys., 229(2), 229–269 (2002); [arXiv:hep-th/0108110](#).
- [KrP] Krichever, I.M., Phong, D.H.: *On the integrable geometry of soliton equations and  $N=2$  supersymmetric gauge theories*. J. Differential Geom. 45(2), 349–389 (1997).
- [KrS] Krichever, I.M., Sheinman, O.K.: *Lax operator algebras*. Funct. Anal. and Appl. 41(4), 284–294 (2007). [arXiv:math/0701648 \[math.RT\]](#).
- [M] Marshall, I.: *A new model in the Calogero-Ruijsenaars family*, Comm. Math. Phys. 338, 563–587 (2015); [arXiv:1311.4641 \[math-ph\]](#).

- [N1] Nekrasov, N.: *Infinite-dimensional algebras, many-body systems and gauge theories*. In: Moscow Seminar in Mathematical Physics, AMS Transl. Ser. 2, Vol. 191, 263–299 (1999).
- [N2] Nakajima, H.: *Instantons on ALE spaces, quiver varieties, and Kac–Moody algebras*. Duke Math. J. 76(2), 365–416, (1994).
- [O] Oblomkov, A.: *Double affine Hecke algebras and Calogero–Moser spaces*. Represent. Theory 8, 243–266 (2004); [arXiv:math/0303190](#) [[math.RT](#)].
- [P] Pusztai, B.G.: *Action-angle duality between the  $C(n)$ -type hyperbolic Sutherland and the rational Ruijsenaars–Schneider–van Diejen models*. Nucl. Phys. B 853, 139–173 (2011); [arXiv:1106.2943](#) [[math-ph](#)].
- [RS] Ruijsenaars, S.N.M., Schneider, H.: *A new class of integrable systems and its relation to solitons*. Ann. Physics 170(2), 370–405 (1986).
- [R] Ruijsenaars, S.N.M.: *Action-angle maps and scattering theory for some finite-dimensional integrable systems I. The pure soliton case*. Commun. Math. Phys. 115, 127–165 (1988).
- [VD] van Diejen, J.F.: *Difference Calogero–Moser systems and finite Toda chains*. J. Math. Phys. 36, 1299–1323 (1995).
- [VDE] van Diejen, J.F., Emsiz, E.: *Spectrum and Eigenfunctions of the Lattice Hyperbolic Ruijsenaars–Schneider System with Exponential Morse Term*. Ann. Henri Poincaré 17, 1615–1629 (2016); [arXiv:1508.03829](#) [[math-ph](#)].
- [VdB1] Van den Bergh, M.: *Double Poisson algebras*. Trans. Amer. Math. Soc., 360 no. 11, 5711–5769 (2008); [arXiv:math/0410528](#) [[math.QA](#)].
- [VdB2] Van den Bergh, M.: *Non-commutative quasi-Hamiltonian spaces*. In: Poisson geometry in mathematics and physics, volume 450 of Contemp. Math., 273–299. Amer. Math. Soc., Providence, RI (2008); [arXiv:math/0703293](#) [[math.QA](#)].
- [W] Wilson, G.: *Collisions of Calogero–Moser particles and an adelic Grassmannian (With an appendix by I. G. Macdonald)*. Invent. Math. 133(1), 1–41 (1998).
- [Y] Yamakawa, D.: *Geometry of multiplicative preprojective algebra*. Int. Math. Res. Pap. 2008, Art. ID rpn008, 77pp; [arXiv:0710.2649](#) [[math.SG](#)].

(Oleg Chalykh) SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS, LS2 9JT, UK,  
*E-mail address:* [o.chalykh@leeds.ac.uk](mailto:o.chalykh@leeds.ac.uk)

(Maxime Fairon) SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS, LS2 9JT, UK,  
*E-mail address:* [mmmfai@leeds.ac.uk](mailto:mmmfai@leeds.ac.uk)