

A properly infinite Banach $*$ -algebra with a non-zero, bounded trace

H. G. Dales, Niels Jakob Laustsen* and Charles J. Read

Abstract

A properly infinite C^* -algebra has no non-zero traces. We construct properly infinite Banach $*$ -algebras with non-zero, bounded traces, and show that there are even such algebras which are fairly ‘close’ to the class of C^* -algebras, in the sense that they can be hermitian or $*$ -semisimple.

Key words: Trace, properly infinite Banach algebra, Banach $*$ -algebra, semigroup Banach algebra.

2000 *Mathematics Subject Classification:* primary 46K05; secondary 43A20, 46H20, 16N20.

To appear in *Studia Mathematica*.

Introduction

Let \mathcal{A} be a complex algebra. The *commutator* of two elements a and b in \mathcal{A} is given by $[a, b] := ab - ba$. A *trace* on \mathcal{A} is a linear functional $\tau: \mathcal{A} \rightarrow \mathbb{C}$ satisfying $\langle ab, \tau \rangle = \langle ba, \tau \rangle$ for each a and b in \mathcal{A} . Clearly, a trace maps each sum of commutators to 0, and, conversely, an element which is mapped to 0 by each trace is necessarily a sum of commutators. It follows that the space of traces on \mathcal{A} encodes important information about the non-commutative structure of \mathcal{A} , and so it is of interest to study this space. In particular, it plays a significant role in the study of Lie derivations and other maps on Banach algebras; see [10] and [14], for example, for developments of this theory.

A unital algebra is *properly infinite* if it contains two idempotent elements which are orthogonal to each other and such that each is equivalent to the identity (see Definition 1.1, below, for details of the unexplained terminology). Fack has shown that each element in a unital, properly infinite C^* -algebra is the sum of five commutators (see [9, Theorem 2.1]), and so the zero functional is the only trace on a unital, properly infinite C^* -algebra. Recently, Pop has given a beautifully simple proof that two commutators in fact suffice (see [17]).

*Supported by the Danish Natural Science Research Council.

In another direction, the second-named author has studied commutators of bounded linear operators on Banach spaces. Let \mathfrak{X} be a Banach space, and denote by $\mathcal{B}(\mathfrak{X})$ the Banach algebra of all bounded linear operators on \mathfrak{X} . It is shown in [12, Proposition 3.7] that, in the case where \mathfrak{X} is isomorphic to an infinite direct sum of copies of itself in a certain technical sense, each bounded linear operator on \mathfrak{X} is the sum of two commutators, and so the zero functional is the only trace on $\mathcal{B}(\mathfrak{X})$ in this case. Prompted by this result, Villena raised the following question:

Let \mathfrak{X} be a Banach space such that \mathfrak{X} is isomorphic to $\mathfrak{X} \oplus \mathfrak{X}$. Is it true that the zero functional is the only trace on the Banach algebra $\mathcal{B}(\mathfrak{X})$?

This question is related to the existence of traces on properly infinite algebras because the Banach algebra $\mathcal{B}(\mathfrak{X})$ is properly infinite if and only if the Banach space \mathfrak{X} contains a complemented subspace isomorphic to $\mathfrak{X} \oplus \mathfrak{X}$ (see [13, Proposition 1.9]). Taking account of this fact and of the above-mentioned results for C^* -algebras, the following question, raised in [12, Question 3.15], emerges as a natural generalization of Villena's question:

Let \mathcal{A} be a unital, properly infinite algebra. Is it true that the zero functional is the only trace on \mathcal{A} ?

In this paper we answer this question in the negative by constructing various unital, properly infinite Banach $*$ -algebras, each with a non-zero, bounded trace. We first give an elementary example. However, this example is not semisimple, and we should like to have examples that are semisimple, and indeed 'closer' to the class of C^* -algebras. For this we give a more complicated example that is $*$ -semisimple, in the sense that it has a faithful $*$ -representation on a Hilbert space.

1 Preliminaries

In this section we give the basic definitions, facts, and conventions that apply throughout this paper, and we provide further motivation for our work. Most of the results in this section are well-known, but for completeness we have included some short and easy proofs.

Throughout, all algebras and vector spaces are assumed to be over the field \mathbb{C} of complex numbers. Whenever convenient, we give definitions only for *unital* algebras; the reason for this is that all the concrete algebras to be studied in Sections 2–4 are indeed unital.

The monograph [7] is our standard reference on Banach algebras. Note in particular that we follow the convention of [7, Definition 3.1.1] by requiring that the involution on a Banach $*$ -algebra be isometric, and that $\langle \cdot, \cdot \rangle$ denotes the pairing between a vector space and its dual space.

1.1 Definition Let \mathcal{A} be an algebra. Two elements a and b in \mathcal{A} are *orthogonal* to each other if $ab = 0 = ba$. An element p in \mathcal{A} is *idempotent* if $p^2 = p$. Two idempotent elements p and q in \mathcal{A} are *algebraically equivalent*, written $p \sim q$, if $p = ab$ and $q = ba$ for some elements a and b in \mathcal{A} .

Suppose that \mathcal{A} has an identity $1_{\mathcal{A}}$. Then \mathcal{A} is *finite* if $p = 1_{\mathcal{A}}$ is the only idempotent element in \mathcal{A} such that $p \sim 1_{\mathcal{A}}$. Otherwise, \mathcal{A} is *infinite*. The algebra \mathcal{A} is *properly infinite* if it contains two idempotent elements p_1 and p_2 which are orthogonal to each other and satisfy $p_1 \sim 1_{\mathcal{A}} \sim p_2$. If \mathcal{A} contains an idempotent element p such that $p \sim 1_{\mathcal{A}} \sim 1_{\mathcal{A}} - p$, then we say that \mathcal{A} is *properly infinite in standard form*.

We note that a unital, properly infinite algebra in standard form is indeed properly infinite. Moreover, it is easy to see that \sim defines an equivalence relation on the set of idempotent elements, and that a unital algebra \mathcal{A} is properly infinite if and only if there are elements $a_1, b_1, a_2,$ and b_2 in \mathcal{A} such that

$$b_j a_i = \begin{cases} 1_{\mathcal{A}} & \text{if } i = j \\ 0 & \text{otherwise} \end{cases} \quad (i, j = 1, 2). \quad (1.1)$$

1.2 Examples (i) Each infinite von Neumann-algebra factor is properly infinite, and each properly infinite von Neumann algebra is in standard form. (*E.g.*, see [11, Definition 6.3.1 and Lemma 6.3.3]; that our definition of properly infinite is equivalent to the one given in [11] follows from [7, Proposition 1.3.22(ii)], [11, Lemma 6.3.3], and the remarks following Definition 1.12, below.)

(ii) The Cuntz C^* -algebra \mathcal{O}_n introduced in [4] is properly infinite whenever $2 \leq n \leq \infty$, but only \mathcal{O}_2 is in standard form (see [5] or [6]).

(iii) Let \mathfrak{X} be a Banach space. The Banach algebra $\mathcal{B}(\mathfrak{X})$ is properly infinite if and only if \mathfrak{X} contains a complemented subspace isomorphic to $\mathfrak{X} \oplus \mathfrak{X}$ (see [13, Proposition 1.9]). In the case where \mathfrak{X} is isomorphic to $\mathfrak{X} \oplus \mathfrak{X}$, $\mathcal{B}(\mathfrak{X})$ is in standard form. \square

1.3 Definition A *trace* on an algebra \mathcal{A} is a linear functional $\tau: \mathcal{A} \rightarrow \mathbb{C}$ satisfying

$$\langle ab, \tau \rangle = \langle ba, \tau \rangle \quad (a, b \in \mathcal{A}).$$

For a normed algebra \mathcal{A} , we write $\mathcal{T}(\mathcal{A})$ for the set of bounded traces on \mathcal{A} . This is the *trace space* of \mathcal{A} .

1.4 Traces on quotients Let \mathcal{I} be an ideal in an algebra \mathcal{A} . There is a canonical bijective correspondence between traces τ on \mathcal{A} with $\mathcal{I} \subseteq \ker \tau$ and traces $\hat{\tau}$ on \mathcal{A}/\mathcal{I} , illustrated by the commutative diagram

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{\tau} & \mathbb{C} \\ & \searrow \pi & \nearrow \hat{\tau} \\ & \mathcal{A}/\mathcal{I} & \end{array}$$

where π denotes the quotient homomorphism. In the case where \mathcal{I} is a closed ideal in a normed algebra \mathcal{A} , a trace τ on \mathcal{A} with $\mathcal{I} \subseteq \ker \tau$ is bounded if and only if the corresponding trace $\hat{\tau}$ on \mathcal{A}/\mathcal{I} is bounded.

1.5 Definition A linear functional λ on an algebra \mathcal{A} with an identity $1_{\mathcal{A}}$ is *normalized* if $\langle 1_{\mathcal{A}}, \lambda \rangle = 1$.

1.6 Lemma Let \mathcal{A} be a unital, properly infinite algebra in standard form. Then the identity is the sum of two commutators, and there are no normalized traces on \mathcal{A} .

Proof. Take an idempotent element p in \mathcal{A} such that $p \sim 1_{\mathcal{A}} \sim 1_{\mathcal{A}} - p$, and take a_1, b_1, a_2 , and b_2 in \mathcal{A} such that $a_1 b_1 = p$, $a_2 b_2 = 1_{\mathcal{A}} - p$, and $b_1 a_1 = 1_{\mathcal{A}} = b_2 a_2$. Then we have

$$[b_1, a_1] + [b_2, a_2] = 1_{\mathcal{A}} - p + 1_{\mathcal{A}} - (1_{\mathcal{A}} - p) = 1_{\mathcal{A}}.$$

In particular, it follows that $\langle 1_{\mathcal{A}}, \tau \rangle = 0$ for each trace τ on \mathcal{A} . \square

Given a unital, properly infinite algebra, we shall now describe a general method to produce a subalgebra which is unital and properly infinite in standard form. We are particularly interested in unital, properly infinite algebras in standard form because, in Villena's original question about the existence of non-zero traces on $\mathcal{B}(\mathfrak{X})$ in the case where \mathfrak{X} is isomorphic to $\mathfrak{X} \oplus \mathfrak{X}$ (see p. 2), $\mathcal{B}(\mathfrak{X})$ is indeed properly infinite in standard form.

1.7 Definition Let p be an idempotent element in a unital algebra \mathcal{A} . The *corner* of \mathcal{A} corresponding to p is

$$p\mathcal{A}p := \{pap \mid a \in \mathcal{A}\}.$$

The corner of a unital algebra \mathcal{A} corresponding to an idempotent element p is a subalgebra of \mathcal{A} with identity p (provided that $p \neq 0$). Suppose that \mathcal{A} is a normed algebra. Then the subalgebra $p\mathcal{A}p$ is automatically closed in \mathcal{A} . However, the identity p of $p\mathcal{A}p$ may have norm strictly greater than one, in which case we equip $p\mathcal{A}p$ with its left multiplier norm

$$\|a\| := \sup\{\|ax\| \mid x \in p\mathcal{A}p, \|x\| \leq 1\} \quad (a \in p\mathcal{A}p).$$

This is an algebra norm equivalent to the original norm, and clearly $\|p\| = 1$.

1.8 Proposition Let \mathcal{A} be a unital, properly infinite algebra. Then there is an idempotent element p in \mathcal{A} such that the corner $p\mathcal{A}p$ is properly infinite in standard form.

Proof. Take idempotent elements p_1 and p_2 in \mathcal{A} which are orthogonal to each other and satisfy $p_1 \sim 1_{\mathcal{A}} \sim p_2$, and take elements a_1, b_1, a_2 , and b_2 in \mathcal{A} such that $a_i b_i = p_i$ and $b_i a_i = 1_{\mathcal{A}}$ for $i = 1, 2$. Set $p := 1_{\mathcal{A}} - p_1$, $q := p - p_2 (= 1_{\mathcal{A}} - p_1 - p_2)$, $c_1 := a_2 p$, $d_1 := p b_2$, $c_2 := a_2 a_1 b_2 + q$, and $d_2 := a_2 b_1 b_2 + q$. Then p and q are idempotent, and q, c_1, c_2, d_1 , and d_2 belong to $p\mathcal{A}p$. Direct calculations show that

$$c_1 d_1 + c_2 d_2 = p \quad \text{and} \quad d_j c_i = \begin{cases} p & \text{if } i = j \\ 0 & \text{otherwise} \end{cases} \quad (i, j = 1, 2),$$

and so the corner $p\mathcal{A}p$ is properly infinite in standard form. \square

For later reference, we record here how the spectrum of an element may change when passing to a corner. We write $\sigma_{\mathcal{A}}(a)$ for the spectrum of an element a in an algebra \mathcal{A} , and we denote by $\nu_{\mathcal{A}}(a)$ its spectral radius. When no ambiguity arises, we shall omit the subscript ‘ \mathcal{A} ’.

1.9 Lemma *Let p be a non-zero, idempotent element in a unital algebra \mathcal{A} . For each element a in $p\mathcal{A}p$,*

$$\sigma_{p\mathcal{A}p}(a) \subseteq \sigma_{\mathcal{A}}(a) \subseteq \sigma_{p\mathcal{A}p}(a) \cup \{0\}.$$

In particular, $\nu_{p\mathcal{A}p}(a) = \nu_{\mathcal{A}}(a)$. □

As explained in the introduction, it is our aim to find a unital, properly infinite Banach algebra \mathcal{A} with a non-zero, bounded trace such that \mathcal{A} is otherwise as ‘close’ as possible to the class of C^* -algebras, where ‘close’ means possessing standard properties. We shall now make this more explicit by defining the properties in which we are interested.

1.10 Definition Let \mathcal{A} be a unital $*$ -algebra. Then:

- (i) \mathcal{A} is *hermitian* if $\sigma(a)$ is contained in \mathbb{R} for each self-adjoint element a in \mathcal{A} .
- (ii) \mathcal{A} is *symmetric* if $1_{\mathcal{A}} + a^*a$ is invertible for each a in \mathcal{A} .
- (iii) A linear functional $\lambda: \mathcal{A} \rightarrow \mathbb{C}$ is *positive* if $\langle a^*a, \lambda \rangle \geq 0$ for each a in \mathcal{A} . A normalized, positive linear functional on \mathcal{A} is called a *state*. A state which is also a trace is called a *tracial state*.
- (iv) The *$*$ -radical* of \mathcal{A} is

$$*\text{-rad } \mathcal{A} := \{a \in \mathcal{A} \mid \langle a^*a, \lambda \rangle = 0 \text{ for each state } \lambda \text{ on } \mathcal{A}\}.$$

If $*\text{-rad } \mathcal{A} = \{0\}$, then \mathcal{A} is *$*$ -semisimple*.

A symmetric, unital $*$ -algebra is hermitian. The converse is true for a unital Banach algebra with an involution; this is Shirali–Ford’s theorem. The $*$ -radical of a unital $*$ -algebra \mathcal{A} is a $*$ -ideal in \mathcal{A} . In the case where \mathcal{A} is a unital Banach $*$ -algebra, the $*$ -radical of \mathcal{A} is automatically closed, it contains the radical of \mathcal{A} , and \mathcal{A} is $*$ -semisimple if and only if \mathcal{A} has a faithful, unital $*$ -representation on a Hilbert space (*e.g.*, see [7, Proposition 1.10.22(iii), Theorem 3.1.10, Proposition 1.10.16(ii), p. 347, and Theorem 3.1.17]).

It is fundamental in the theory of C^* -algebras that a unital C^* -algebra is hermitian and $*$ -semisimple (*e.g.*, see [7, Proposition 3.2.3(v) and Corollary 3.2.13]), and that a unital C^* -algebra has a non-zero, bounded trace if and only if it has a tracial state. This leads us to consider the following questions.

1.11 Questions Let \mathcal{A} be a unital, properly infinite Banach $*$ -algebra with a non-zero, bounded trace. Is it possible that:

- (i) \mathcal{A} is hermitian?
- (ii) \mathcal{A} is $*$ -semisimple?

- (iii) \mathcal{A} is both hermitian and $*$ -semisimple?
- (iv) \mathcal{A} has a tracial state?

In Sections 2 and 3, we answer Questions 1.11(i) and (ii) in the affirmative. We have not been able to resolve Questions 1.11(iii) and (iv); we can only show that none of the unital, properly infinite Banach $*$ -algebras considered in this paper provides a positive answer to any of these questions. In the case of Question 1.11(iv), this relies on the following, formally stronger, version of Definition 1.1 for $*$ -algebras.

1.12 Definition Let \mathcal{A} be a $*$ -algebra. A self-adjoint, idempotent element in \mathcal{A} is called a *projection*. Two projections p and q in \mathcal{A} are *Murray–von Neumann equivalent*, written $p \approx q$, if $p = a^*a$ and $q = aa^*$ for some element a in \mathcal{A} .

Suppose that \mathcal{A} is unital. Then the terms *$*$ -finite*, *$*$ -infinite*, *$*$ -properly infinite*, and *$*$ -properly infinite in standard form* are defined as in Definition 1.1, just with the term ‘idempotent element’ replaced by ‘projection’ and the relation \sim replaced by \approx .

We note that \approx is an equivalence relation on the set of projections. Murray–von Neumann equivalent projections are clearly algebraically equivalent, and so a unital, $*$ -properly infinite $*$ -algebra is properly infinite, and a unital, finite $*$ -algebra is $*$ -finite. None of these implications can be reversed in general, as we shall show in Section 4.

However, in a symmetric, unital $*$ -algebra, each idempotent element is algebraically equivalent to a projection and, moreover, in a C^* -algebra, two projections are Murray–von Neumann equivalent if and only if they are algebraically equivalent (*e.g.*, see [7, Propositions 1.10.21(i) and 3.2.10]). Along the same lines, one can show that a unital C^* -algebra is properly infinite (infinite, finite, respectively) if and only if it is $*$ -properly infinite ($*$ -infinite, $*$ -finite, respectively).

All our examples of unital, properly infinite Banach $*$ -algebras are in fact $*$ -properly infinite. This implies that they have no tracial states, as the following easy consequence of the Cauchy–Schwarz inequality (*e.g.*, see [7, Proposition 1.10.13(iii)]) shows.

1.13 Lemma *Let λ be a positive linear functional on a $*$ -algebra \mathcal{A} with an identity $1_{\mathcal{A}}$. Then λ is non-zero if and only if $\langle 1_{\mathcal{A}}, \lambda \rangle$ is non-zero. In particular, \mathcal{A} has a non-zero, positive trace if and only if \mathcal{A} has a tracial state. \square*

1.14 Corollary *There are no tracial states on a unital, $*$ -properly infinite $*$ -algebra.*

Proof. Let τ be a positive trace on a unital, $*$ -properly infinite $*$ -algebra \mathcal{A} . Take projections p_1 and p_2 in \mathcal{A} which are orthogonal to each other and satisfy $p_1 \approx 1_{\mathcal{A}} \approx p_2$. Then $1_{\mathcal{A}} - p_1 - p_2$ is a projection, and we have

$$\langle p_1, \tau \rangle = \langle 1_{\mathcal{A}}, \tau \rangle = \langle p_2, \tau \rangle \geq 0 \quad \text{and} \quad \langle 1_{\mathcal{A}} - p_1 - p_2, \tau \rangle \geq 0.$$

This implies that $\langle 1_{\mathcal{A}}, \tau \rangle = 0$, so that $\tau = 0$ by Lemma 1.13. \square

We conclude this preliminary section with some basic facts about corners in $*$ -algebras.

1.15 Proposition *Let \mathcal{A} be a unital $*$ -algebra.*

- (i) *For each projection p in \mathcal{A} , the corner $p\mathcal{A}p$ is a sub- $*$ -algebra of \mathcal{A} .*
- (ii) *Suppose that \mathcal{A} is $*$ -properly infinite. Then \mathcal{A} contains a projection p such that the corner $p\mathcal{A}p$ is $*$ -properly infinite in standard form.*
- (iii) *Suppose that \mathcal{A} is hermitian. Then, for each non-zero projection p in \mathcal{A} , the corner $p\mathcal{A}p$ is hermitian.*
- (iv) *Suppose that \mathcal{A} is $*$ -semisimple. Then, for each non-zero projection p in \mathcal{A} , the corner $p\mathcal{A}p$ is $*$ -semisimple.*

Proof. Clause (i) is clear, clause (ii) follows from an obvious modification of the proof of Proposition 1.8, and clause (iii) is immediate from Lemma 1.9.

Clause (iv) is a consequence of [15, Theorem 9.7.17(a)], or it can be proved easily as follows. Let p be a non-zero projection in \mathcal{A} . For each non-zero element a in $p\mathcal{A}p$, we can take a state κ on \mathcal{A} such that $\langle a^*a, \kappa \rangle$ is non-zero. Then $\lambda := \kappa|_{p\mathcal{A}p}$ is a non-zero, positive linear functional on $p\mathcal{A}p$. Lemma 1.13 implies that $\langle p, \lambda \rangle$ is non-zero, and so $\mu := \lambda / \langle p, \lambda \rangle$ is a state on $p\mathcal{A}p$ such that $\langle a^*a, \mu \rangle$ is non-zero. Hence a is not in $*$ -rad($p\mathcal{A}p$). \square

2 An elementary and hermitian example

In this section we describe an elementary method to construct a unital, properly infinite Banach algebra with a non-zero, bounded trace. Moreover, we extend this method to encompass Banach $*$ -algebras, and thus show that there exists a hermitian, unital, $*$ -properly infinite Banach $*$ -algebra with a non-zero, bounded trace.

The corner stone is the following well-known construction.

2.1 Construction Let \mathcal{A} be a Banach algebra, and let \mathcal{X} be a Banach \mathcal{A} -bimodule. The Cartesian product $\mathcal{A} \oplus \mathcal{X}$ is a Banach algebra for the coordinatewise-defined vector-space operations, the multiplication

$$(a, x)(b, y) := (ab, a \cdot y + x \cdot b) \quad (a, b \in \mathcal{A}, x, y \in \mathcal{X}),$$

and the norm

$$\|(a, x)\| := \max\{\|a\|, \|x\|\} \quad (a \in \mathcal{A}, x \in \mathcal{X})$$

(*e.g.*, see [7, p. 239]).

Suppose that \mathcal{A} has an identity $1_{\mathcal{A}}$, and that \mathcal{X} is a unital Banach \mathcal{A} -bimodule in the sense that $1_{\mathcal{A}} \cdot x = x = x \cdot 1_{\mathcal{A}}$ for each x in \mathcal{X} . Then $(1_{\mathcal{A}}, 0)$ is an identity in the Banach algebra $\mathcal{A} \oplus \mathcal{X}$.

2.2 Proposition *Let \mathcal{A} be a unital Banach algebra, and let \mathcal{X} be a unital Banach \mathcal{A} -bimodule. Then the Banach algebra $\mathcal{A} \oplus \mathcal{X}$ is properly infinite if and only if \mathcal{A} is properly infinite. In this case, $\mathcal{A} \oplus \mathcal{X}$ is in standard form if and only if \mathcal{A} is in standard form.*

Proof. Suppose that the four elements $a_1, b_1, a_2,$ and b_2 in \mathcal{A} satisfy (1.1). Then we have

$$(b_j, 0)(a_i, 0) = \begin{cases} (1_{\mathcal{A}}, 0) & \text{if } i = j \\ (0, 0) & \text{otherwise} \end{cases} \quad (i, j = 1, 2),$$

and so $\mathcal{A} \oplus \mathcal{X}$ is properly infinite.

Conversely, suppose that $(a_1, x_1), (b_1, y_1), (a_2, x_2),$ and (b_2, y_2) in $\mathcal{A} \oplus \mathcal{X}$ satisfy

$$(b_j, y_j)(a_i, x_i) = \begin{cases} (1_{\mathcal{A}}, 0) & \text{if } i = j \\ (0, 0) & \text{otherwise} \end{cases} \quad (i, j = 1, 2).$$

Then certainly $a_1, b_1, a_2,$ and b_2 satisfy (1.1), so that \mathcal{A} is properly infinite.

The final clause is proved analogously. \square

2.3 Lemma *Let \mathcal{A} be a Banach algebra, let \mathcal{X} be a Banach \mathcal{A} -bimodule, and let λ be a non-zero, bounded linear functional on \mathcal{X} . Then*

$$\tau_\lambda: (a, x) \mapsto \langle x, \lambda \rangle, \quad \mathcal{A} \oplus \mathcal{X} \rightarrow \mathbb{C}, \quad (2.1)$$

is a non-zero, bounded linear functional on $\mathcal{A} \oplus \mathcal{X}$, and τ_λ is a trace if and only if

$$\langle a \cdot x, \lambda \rangle = \langle x \cdot a, \lambda \rangle \quad (a \in \mathcal{A}, x \in \mathcal{X}). \quad (2.2)$$

Proof. It is straightforward to see that τ_λ is non-zero, bounded, and linear. By the definitions, τ_λ is a trace if and only if

$$\langle a \cdot x + y \cdot b, \lambda \rangle = \langle x \cdot a + b \cdot y, \lambda \rangle \quad (a, b \in \mathcal{A}, x, y \in \mathcal{X}). \quad (2.3)$$

This is certainly satisfied if (2.2) holds, and, conversely, taking $b = 0$ and $y = 0$ in (2.3) yields (2.2). \square

2.4 Example Let \mathcal{A} be a unital Banach algebra. The projective tensor product $\widehat{\mathcal{A} \otimes \mathcal{A}}$ of two copies of \mathcal{A} is a unital Banach \mathcal{A} -bimodule for maps which satisfy

$$a \cdot (b \otimes c) = ab \otimes c \quad \text{and} \quad (b \otimes c) \cdot a = b \otimes ca \quad (a, b, c \in \mathcal{A})$$

(*e.g.*, see [7, Corollary 2.6.5]). Take a non-zero, bounded linear functional μ on \mathcal{A} . Then the map

$$(a, b) \mapsto \langle ba, \mu \rangle, \quad \mathcal{A} \times \mathcal{A} \rightarrow \mathbb{C},$$

is a non-zero, bounded bilinear form, and so there is a non-zero, bounded linear functional $\lambda: \mathcal{A} \widehat{\otimes} \mathcal{A} \rightarrow \mathbb{C}$ such that $\langle a \otimes b, \lambda \rangle = \langle ba, \mu \rangle$ for each a and b in \mathcal{A} . It follows that λ satisfies condition (2.2) in Lemma 2.3 because

$$\langle a \cdot (b \otimes c), \lambda \rangle = \langle cab, \mu \rangle = \langle (b \otimes c) \cdot a, \lambda \rangle \quad (a, b, c \in \mathcal{A}),$$

and so the functional $\tau_\lambda: \mathcal{A} \oplus (\mathcal{A} \widehat{\otimes} \mathcal{A}) \rightarrow \mathbb{C}$ defined by (2.1) is a non-zero, bounded trace. \square

Combining Proposition 2.2 and Example 2.4 gives the following theorem which answers [12, Question 3.15] in the negative.

2.5 Theorem *Let \mathcal{A} be a unital, properly infinite Banach algebra. Then $\mathcal{A} \oplus (\mathcal{A} \widehat{\otimes} \mathcal{A})$ is a unital, properly infinite Banach algebra which has a non-zero, bounded trace. Moreover, $\mathcal{A} \oplus (\mathcal{A} \widehat{\otimes} \mathcal{A})$ is in standard form if and only if \mathcal{A} is in standard form. \square*

2.6 Remark The above construction has some obvious limitations. Indeed, let \mathcal{A} be a unital Banach algebra, and let \mathcal{X} be a unital Banach \mathcal{A} -bimodule. Then:

- (i) The Banach algebra $\mathcal{A} \oplus \mathcal{X}$ is never semisimple; the radical of $\mathcal{A} \oplus \mathcal{X}$ is given by

$$\text{rad}(\mathcal{A} \oplus \mathcal{X}) = (\text{rad } \mathcal{A}) \oplus \mathcal{X} \quad (2.4)$$

(see [7, Theorem 1.8.14(ii)]). In Section 3 we shall use an entirely different approach to find an example of a semisimple, unital, properly infinite Banach algebra with a non-zero, bounded trace.

- (ii) For each non-zero, bounded linear functional λ on \mathcal{X} , we have

$$\langle 1_{\mathcal{A} \oplus \mathcal{X}}, \tau_\lambda \rangle = \langle 0, \lambda \rangle = 0.$$

It follows that τ_λ cannot be normalized, and Lemma 1.13 implies that this construction cannot be used to resolve Question 1.11(iv).

We shall now refine the above results to show that there is a hermitian, unital, *-properly infinite Banach *-algebra with a non-zero, bounded trace.

We begin by extending Construction 2.1 to encompass Banach *-algebras.

2.7 Construction Let \mathcal{A} be a Banach *-algebra, let \mathcal{X} be a Banach \mathcal{A} -bimodule with an isometric linear involution, and suppose that the involutions on \mathcal{A} and \mathcal{X} are tied together through

$$(a \cdot x)^* = x^* \cdot a^* \quad (a \in \mathcal{A}, x \in \mathcal{X}). \quad (2.5)$$

Then the Banach algebra $\mathcal{A} \oplus \mathcal{X}$ from Construction 2.1 is a Banach *-algebra for the involution

$$(a, x)^* := (a^*, x^*) \quad (a \in \mathcal{A}, x \in \mathcal{X}). \quad (2.6)$$

2.8 Example Let \mathcal{A} be a unital Banach $*$ -algebra. The definition

$$(a \otimes b)^* := b^* \otimes a^* \quad (a, b \in \mathcal{A})$$

induces an isometric linear involution on the Banach \mathcal{A} -bimodule $\mathcal{A} \widehat{\otimes} \mathcal{A}$ from Example 2.4, and (2.5) is satisfied in this case. Thus, $\mathcal{A} \oplus (\mathcal{A} \widehat{\otimes} \mathcal{A})$ is a unital Banach $*$ -algebra for the involution given by (2.6). \square

2.9 Proposition Let \mathcal{A} be a unital Banach $*$ -algebra, and let \mathcal{X} be a unital Banach \mathcal{A} -bimodule with an isometric linear involution satisfying (2.5). Then:

- (i) $\mathcal{A} \oplus \mathcal{X}$ is hermitian if and only if \mathcal{A} is hermitian;
- (ii) $\mathcal{A} \oplus \mathcal{X}$ is $*$ -properly infinite if and only if \mathcal{A} is $*$ -properly infinite. In this case, $\mathcal{A} \oplus \mathcal{X}$ is in standard form if and only if \mathcal{A} is in standard form.

Proof. By (2.4) (or a direct computation) we have

$$\sigma_{\mathcal{A} \oplus \mathcal{X}}(a, x) = \sigma_{\mathcal{A}}(a) \quad (a \in \mathcal{A}, x \in \mathcal{X}).$$

Clause (i) is an immediate consequence of this. Clause (ii) is proved analogously to Proposition 2.2. \square

Combining this with Examples 2.4 and 2.8 enables us to answer Question 1.11(i).

2.10 Corollary Let \mathcal{A} be a hermitian, unital, $*$ -properly infinite Banach $*$ -algebra. Then $\mathcal{A} \oplus (\mathcal{A} \widehat{\otimes} \mathcal{A})$ is a hermitian, unital, $*$ -properly infinite Banach $*$ -algebra with a non-zero, bounded trace. Moreover, $\mathcal{A} \oplus (\mathcal{A} \widehat{\otimes} \mathcal{A})$ is in standard form if and only if \mathcal{A} is in standard form. \square

3 A semisimple example

3.1 Semigroup Banach algebras Let S be a semigroup. We use the monograph [3] as our standard reference on semigroups. The Banach space

$$\ell_1(S) := \left\{ f: S \rightarrow \mathbb{C} \mid \|f\|_1 := \sum_{s \in S} |f(s)| < \infty \right\},$$

equipped with the pointwise-defined vector-space operations and the norm $\|\cdot\|_1$, is a Banach algebra for the *convolution product* \star defined as follows. For each s in S , let δ_s be the point mass at s , that is,

$$\delta_s(t) := \begin{cases} 1 & \text{if } s = t \\ 0 & \text{otherwise} \end{cases} \quad (t \in S).$$

Then each element f of $\ell_1(S)$ can be uniquely expressed as an absolutely convergent sum $f = \sum_{s \in S} f(s)\delta_s$, and the convolution product is determined by the formula $\delta_s \star \delta_t := \delta_{st}$

for each s and t in S . We call $\ell_1(S)$ the *semigroup Banach algebra* of S . In the case where the semigroup S has a neutral element e , the element δ_e is an identity in the algebra $\ell_1(S)$.

The dual space of $\ell_1(S)$ is canonically identified with the Banach space

$$\ell_\infty(S) := \left\{ \lambda: S \rightarrow \mathbb{C} \mid \|\lambda\|_\infty := \sup\{|\lambda(s)| \mid s \in S\} < \infty \right\}$$

via the pairing

$$\langle f, \lambda \rangle := \sum_{s \in S} f(s)\lambda(s) \quad (f \in \ell_1(S), \lambda \in \ell_\infty(S)).$$

In particular, we regard the trace space of $\ell_1(S)$ as a subspace of $\ell_\infty(S)$:

$$\mathcal{T}(\ell_1(S)) = \left\{ \tau \in \ell_\infty(S) \mid \tau(st) = \tau(ts) \quad (s, t \in S) \right\}. \quad (3.1)$$

3.2 Semigroups with a zero element Let S be a semigroup with at least two elements. An element θ in S is a *zero element* if $s\theta = \theta = \theta s$ for each s in S . (For technical reasons, it is convenient to exclude the trivial semigroup $\{0\}$ from the class of semigroups with zero element.) Clearly, a semigroup can have at most one zero element.

Let θ be a zero element in a semigroup S . We write S^\blacksquare for $S \setminus \{\theta\}$. The closed ideal in $\ell_1(S)$ generated by δ_θ is just $\mathbb{C}\delta_\theta$. It is often preferable to consider the quotient algebra $\ell_1^\theta(S) := \ell_1(S)/\mathbb{C}\delta_\theta$ instead of $\ell_1(S)$. As a Banach space, $\ell_1^\theta(S)$ is isometrically isomorphic to $\ell_1(S^\blacksquare)$. The product, which we denote by $\widehat{\star}$, induced on $\ell_1(S^\blacksquare)$ via this isometric isomorphism is determined by

$$\delta_s \widehat{\star} \delta_t = \begin{cases} 0 & \text{if } st = \theta \\ \delta_{st} & \text{otherwise} \end{cases} \quad (s, t \in S^\blacksquare). \quad (3.2)$$

It follows from Paragraph 1.4 that the trace space of $\ell_1(S^\blacksquare)$ is given by

$$\mathcal{T}(\ell_1(S^\blacksquare)) = \left\{ \tau|_{S^\blacksquare} \mid \tau \in \mathcal{T}(\ell_1(S)) \text{ with } \tau(\theta) = 0 \right\} \subseteq \ell_\infty(S^\blacksquare). \quad (3.3)$$

3.3 Semigroups with an involution An *involution* (or an *involutional anti-automorphism* in the terminology of [3]) on a semigroup S is a map $s \mapsto s^*$, $S \rightarrow S$, such that $(s^*)^* = s$ and $(st)^* = t^*s^*$ for each s and t in S . An element s in S is *self-adjoint* if $s^* = s$.

Let S be a semigroup with an involution. Then the formula

$$f^*(s) := \overline{f(s^*)} \quad (f \in \ell_1(S), s \in S) \quad (3.4)$$

defines an isometric involution on $\ell_1(S)$, so that $\ell_1(S)$ is a Banach $*$ -algebra. Suppose that S has a zero element θ . Necessarily, θ is self-adjoint, and $\ell_1^\theta(S)$ inherits the isometric involution from $\ell_1(S)$. The ensuing isometric involution on $\ell_1(S^\blacksquare)$ is still given by the formula (3.4), above, just with f belonging to $\ell_1(S^\blacksquare)$ and s belonging to S^\blacksquare .

3.4 Definition Following Renault (see [18, p. 141f]), we denote by S_2 the *second Cuntz semigroup*, that is, S_2 is the semigroup with an involution $*$, neutral element e , zero element θ , and generators s_1, s_2, s_1^* , and s_2^* subject to the relations

$$s_i^* s_j = \begin{cases} e & \text{if } i = j \\ \theta & \text{otherwise} \end{cases} \quad (i, j = 1, 2). \quad (3.5)$$

3.5 Theorem *The unital Banach $*$ -algebra $\ell_1(S_2^\blacksquare)$ is $*$ -properly infinite.*

Proof. By (3.2) and (3.5), the elements $a_1 := \delta_{s_1}$ and $a_2 := \delta_{s_2}$ in $\ell_1(S_2^\blacksquare)$ satisfy

$$a_j^* \widehat{\star} a_i = \delta_{s_j^*} \widehat{\star} \delta_{s_i} = \begin{cases} \delta_e & \text{if } i = j \\ 0 & \text{otherwise} \end{cases} \quad (i, j = 1, 2). \quad \square$$

In particular, Corollary 1.14 shows that there are no tracial states on $\ell_1(S_2^\blacksquare)$. We shall next prove that, nevertheless, there are non-zero, bounded traces on $\ell_1(S_2^\blacksquare)$. First we shall give a specific example of a normalized, bounded trace on $\ell_1(S_2^\blacksquare)$, and then we shall characterize all the bounded traces on $\ell_1(S_2^\blacksquare)$ among the elements of $\ell_\infty(S_2^\blacksquare)$.

3.6 Definition Set

$$\mathbf{I} := \{\emptyset\} \cup \{(i_1, i_2, \dots, i_n) \mid n \in \mathbb{N}, i_1, i_2, \dots, i_n \in \{1, 2\}\},$$

and, for each \mathbf{i} in \mathbf{I} , define an element $s_{\mathbf{i}}$ in S_2 as follows:

$$s_{\mathbf{i}} := \begin{cases} e & \text{for } \mathbf{i} = \emptyset, \\ s_{i_1} s_{i_2} \cdots s_{i_n} & \text{for } \mathbf{i} = (i_1, i_2, \dots, i_n) \in \mathbf{I} \setminus \{\emptyset\}. \end{cases}$$

Moreover, for each \mathbf{i} and \mathbf{j} in \mathbf{I} , define \mathbf{ij} in \mathbf{I} by concatenation:

$$\mathbf{ij} := \begin{cases} \mathbf{i} & \text{for } \mathbf{j} = \emptyset, \\ \mathbf{j} & \text{for } \mathbf{i} = \emptyset, \\ (i_1, \dots, i_m, j_1, \dots, j_n) & \text{for } \mathbf{i} = (i_1, \dots, i_m) \text{ and } \mathbf{j} = (j_1, \dots, j_n). \end{cases}$$

Trivially, we have $s_{\mathbf{i}} s_{\mathbf{j}} = s_{\mathbf{ij}}$ for each \mathbf{i} and \mathbf{j} in \mathbf{I} . More importantly, the following fundamental result, which is essentially due to Cuntz (see [4, Lemmas 1.2 and 1.3] or [16, formulae (1.4), (4.8), and (4.9)]), holds.

3.7 Lemma (i) *For each \mathbf{i} and \mathbf{j} in \mathbf{I} ,*

$$s_{\mathbf{i}}^* s_{\mathbf{j}} = \begin{cases} s_{\mathbf{k}}^* & \text{if } \mathbf{i} = \mathbf{jk} \text{ for some } \mathbf{k} \in \mathbf{I} \\ s_{\mathbf{k}} & \text{if } \mathbf{j} = \mathbf{ik} \text{ for some } \mathbf{k} \in \mathbf{I} \\ \theta & \text{otherwise.} \end{cases}$$

(ii) For each s in S_2^\blacksquare , there are unique \mathbf{i} and \mathbf{j} in \mathbf{I} such that $s = s_{\mathbf{i}}s_{\mathbf{j}}^*$. \square

Lemma 3.7 gives a nuts-and-bolts description of the Banach-algebra structure of $\ell_1(S_2^\blacksquare)$. Indeed, clause (ii) implies that each f in $\ell_1(S_2^\blacksquare)$ has a unique representation as an absolutely convergent series of the form

$$f = \sum_{\mathbf{i}, \mathbf{j} \in \mathbf{I}} \alpha_{\mathbf{i}, \mathbf{j}} \delta_{s_{\mathbf{i}}s_{\mathbf{j}}^*}, \quad (3.6)$$

where $\alpha_{\mathbf{i}, \mathbf{j}} = f(s_{\mathbf{i}}s_{\mathbf{j}}^*)$ for each \mathbf{i} and \mathbf{j} in \mathbf{I} , and $\|f\|_1 = \sum |\alpha_{\mathbf{i}, \mathbf{j}}|$. Clause (i) then specifies how to multiply two elements represented as in (3.6).

We are now ready to construct a normalized, bounded trace on $\ell_1(S_2^\blacksquare)$.

3.8 Example Define τ in $\ell_\infty(S_2)$ by

$$\tau(s) = \begin{cases} 1 & \text{if } s = s_{\mathbf{i}}s_{\mathbf{i}}^* \text{ for some } \mathbf{i} \in \mathbf{I} \\ 0 & \text{otherwise} \end{cases} \quad (s \in S_2).$$

Then τ is a normalized, bounded linear functional on $\ell_1(S_2)$. Lemma 3.7 implies that

$$\tau(st) = \begin{cases} 1 & \text{if } s = s_{\mathbf{ij}}s_{\mathbf{kj}}^* \text{ and } t = s_{\mathbf{k}}s_{\mathbf{i}}^* \text{ for some } \mathbf{i}, \mathbf{j}, \mathbf{k} \in \mathbf{I} \\ & \text{or if } s = s_{\mathbf{i}}s_{\mathbf{j}}^* \text{ and } t = s_{\mathbf{jk}}s_{\mathbf{ik}}^* \text{ for some } \mathbf{i}, \mathbf{j}, \mathbf{k} \in \mathbf{I} \\ 0 & \text{otherwise} \end{cases} \quad (s, t \in S_2).$$

It follows that $\tau(st) = \tau(ts)$ for each s and t in S_2 , and so τ is a trace. Since $\tau(\theta) = 0$, we conclude from (3.3) that $\tau|_{S_2^\blacksquare}$ is a normalized, bounded trace on $\ell_1(S_2^\blacksquare)$. \square

As mentioned above, our next aim is to determine *all* the bounded traces on $\ell_1(S_2^\blacksquare)$.

3.9 Definition For elements s and t in a semigroup S , we write $s \wedge t$ if there are elements u and v in S such that $s = uv$ and $t = vu$.

Clearly \wedge is a symmetric relation. In general, it need not be either reflexive or transitive. In the case of interest to us, that is, the second Cuntz semigroup S_2 , \wedge is indeed reflexive (because S_2 has a neutral element), but \wedge is not transitive (for example, $s_1s_2^* \wedge \theta$ and $s_2s_1^* \wedge \theta$, but Lemma 3.7 implies that $\neg(s_1s_2^* \wedge s_2s_1^*)$).

Let $\widehat{\wedge}$ be the smallest equivalence relation containing \wedge , and let $[s]_{\widehat{\wedge}} := \{t \in S \mid t \widehat{\wedge} s\}$ denote the $\widehat{\wedge}$ -equivalence class of s in S . The equivalence relation $\widehat{\wedge}$ has previously been studied in [2] (where it is denoted by \approx) in connection with investigations of the weak amenability of $\ell_1(S)$. The following proposition explains why it is relevant for our purposes.

3.10 Proposition *Let S be a semigroup. An element τ in $\ell_\infty(S)$ is a trace on $\ell_1(S)$ if and only if $\tau(s) = \tau(t)$ whenever s and t in S satisfy $s \widehat{\wedge} t$.*

Proof. Suppose that τ is a trace. Define a relation λ_τ on S by

$$s \lambda_\tau t \iff \tau(s) = \tau(t) \quad (s, t \in S).$$

This is clearly an equivalence relation on S , and it contains λ because τ is a trace. Consequently, λ_τ contains $\widehat{\lambda}$, that is, $\tau(s) = \tau(t)$ whenever s and t in S satisfy $s \widehat{\lambda} t$.

Conversely, suppose that $\tau(s) = \tau(t)$ for each s and t in S with $s \widehat{\lambda} t$. Then, since $\widehat{\lambda}$ contains λ , we have $\tau(uv) = \tau(vu)$ for each u and v in S , and so τ is a trace by (3.1). \square

We shall now describe explicitly the $\widehat{\lambda}$ -equivalence classes in S_2 .

3.11 Lemma (i) *The $\widehat{\lambda}$ -equivalence class of θ is given by*

$$[\theta]_{\widehat{\lambda}} = \{\theta\} \cup \{s_{\mathbf{i}}s_{\mathbf{j}}^* \mid \mathbf{i} = (i_1, \dots, i_m), \mathbf{j} = (j_1, \dots, j_n) \in \mathbf{I} \setminus \{\emptyset\} \text{ with } i_p \neq j_p \text{ for some } p \leq \min\{m, n\}\}.$$

(ii) *For each \mathbf{i} in \mathbf{I} , the $\widehat{\lambda}$ -equivalence classes of $s_{\mathbf{i}}$ and $s_{\mathbf{i}}^*$ are given by*

$$[s_{\mathbf{i}}]_{\widehat{\lambda}} = \{s_{\mathbf{h}\mathbf{j}\mathbf{k}}s_{\mathbf{h}}^* \mid \mathbf{h}, \mathbf{j}, \mathbf{k} \in \mathbf{I}, \mathbf{k}\mathbf{j} = \mathbf{i}\}$$

and

$$[s_{\mathbf{i}}^*]_{\widehat{\lambda}} = \{s_{\mathbf{h}}s_{\mathbf{h}\mathbf{j}\mathbf{k}}^* \mid \mathbf{h}, \mathbf{j}, \mathbf{k} \in \mathbf{I}, \mathbf{k}\mathbf{j} = \mathbf{i}\}.$$

(iii) *Each element of S_2 belongs to either $[\theta]_{\widehat{\lambda}}$, $[e]_{\widehat{\lambda}}$, $[s_{\mathbf{i}}]_{\widehat{\lambda}}$, or $[s_{\mathbf{i}}^*]_{\widehat{\lambda}}$ for some \mathbf{i} in $\mathbf{I} \setminus \{\emptyset\}$.*

Proof. Set

$$E(\theta) = \{\theta\} \cup \{s_{\mathbf{i}}s_{\mathbf{j}}^* \mid \mathbf{i} = (i_1, \dots, i_m), \mathbf{j} = (j_1, \dots, j_n) \in \mathbf{I} \setminus \{\emptyset\} \text{ with } i_p \neq j_p \text{ for some } p \leq \min\{m, n\}\},$$

and, for each \mathbf{i} in \mathbf{I} , set

$$E(s_{\mathbf{i}}) = \{s_{\mathbf{h}\mathbf{j}\mathbf{k}}s_{\mathbf{h}}^* \mid \mathbf{h}, \mathbf{j}, \mathbf{k} \in \mathbf{I}, \mathbf{k}\mathbf{j} = \mathbf{i}\}$$

and

$$E(s_{\mathbf{i}}^*) = \{s_{\mathbf{h}}s_{\mathbf{h}\mathbf{j}\mathbf{k}}^* \mid \mathbf{h}, \mathbf{j}, \mathbf{k} \in \mathbf{I}, \mathbf{k}\mathbf{j} = \mathbf{i}\}.$$

To prove that $E(x)$ is the $\widehat{\lambda}$ -equivalence class of x for each x in the set $\{\theta, s_{\mathbf{i}}, s_{\mathbf{i}}^* \mid \mathbf{i} \in \mathbf{I}\}$, we introduce an equivalence relation $\widetilde{\lambda}$ on S_2 whose equivalence class of x is just $E(x)$, and then show that $\widetilde{\lambda}$ is equal to $\widehat{\lambda}$.

Specifically, we observe that

$$S_2 = E(\theta) \cup \bigcup_{\mathbf{i} \in \mathbf{I}} E(s_{\mathbf{i}}) \cup \bigcup_{\mathbf{i} \in \mathbf{I}} E(s_{\mathbf{i}}^*), \quad (3.7)$$

and that, for each x and y in $\{\theta, s_{\mathbf{i}}, s_{\mathbf{i}}^* \mid \mathbf{i} \in \mathbf{I}\}$, the two sets $E(x)$ and $E(y)$ are either equal or disjoint. This implies that the relation $\tilde{\lambda}$ on S_2 defined by

$$s \tilde{\lambda} t \iff s, t \in E(x) \text{ for some } x \in \{\theta, s_{\mathbf{i}}, s_{\mathbf{i}}^* \mid \mathbf{i} \in \mathbf{I}\} \quad (s, t \in S_2)$$

is an equivalence relation with equivalence classes

$$[\theta]_{\tilde{\lambda}} = E(\theta), \quad [s_{\mathbf{i}}]_{\tilde{\lambda}} = E(s_{\mathbf{i}}), \quad \text{and} \quad [s_{\mathbf{i}}^*]_{\tilde{\lambda}} = E(s_{\mathbf{i}}^*) \quad (\mathbf{i} \in \mathbf{I}). \quad (3.8)$$

For each x in $\{\theta, s_{\mathbf{i}}, s_{\mathbf{i}}^* \mid \mathbf{i} \in \mathbf{I}\}$ and each s in $E(x)$, we have $s \wedge x$, and so $\widehat{\lambda}$ contains $\tilde{\lambda}$.

For the reverse inclusion, it suffices to prove that $\tilde{\lambda}$ contains \wedge . Suppose that $s \wedge t$, where s and t belong to S_2 , and take u and v in S_2 such that $s = uv$ and $t = vu$. We split into four cases.

- (1) $s = t = \theta$: Then s and t both belong to $E(\theta)$.
- (2) $s = \theta$ and $t \neq \theta$: Since $vu = t \neq \theta$, necessarily $u \neq \theta$ and $v \neq \theta$, and, by Lemma 3.7, there are two subcases to consider.
 - (2a) $u = s_{\mathbf{h}\mathbf{i}}s_{\mathbf{j}}^*$ and $v = s_{\mathbf{k}}s_{\mathbf{h}}^*$ for some $\mathbf{h}, \mathbf{i}, \mathbf{j}, \mathbf{k}$ in \mathbf{I} : Since $\theta = s = uv = s_{\mathbf{h}\mathbf{i}}s_{\mathbf{j}}^*s_{\mathbf{k}}s_{\mathbf{h}}^*$, Lemma 3.7(i) implies that \mathbf{j} and \mathbf{k} are non-empty, say $\mathbf{j} = (j_1, \dots, j_m)$ and $\mathbf{k} = (k_1, \dots, k_n)$, with $j_p \neq k_p$ for some $p \leq \min\{m, n\}$. Then $t = vu = s_{\mathbf{k}\mathbf{i}}s_{\mathbf{j}}^*$ belongs to $E(\theta)$ because $\mathbf{k}\mathbf{i}$ and \mathbf{j} differ in the p^{th} coordinate.
 - (2b) $u = s_{\mathbf{h}}s_{\mathbf{i}}^*$ and $v = s_{\mathbf{j}}s_{\mathbf{h}\mathbf{k}}^*$ for some $\mathbf{h}, \mathbf{i}, \mathbf{j}, \mathbf{k}$ in \mathbf{I} : As above, we obtain $\mathbf{i} = (i_1, \dots, i_m)$ and $\mathbf{j} = (j_1, \dots, j_n)$ with $i_p \neq j_p$ for some $p \leq \min\{m, n\}$. It follows that $t = vu = s_{\mathbf{j}}s_{\mathbf{i}\mathbf{k}}^*$ belongs to $E(\theta)$ because \mathbf{j} and $\mathbf{i}\mathbf{k}$ differ in the p^{th} coordinate.
- (3) $s \neq \theta$ and $t = \theta$: This is similar to case (2).
- (4) $s \neq \theta$ and $t \neq \theta$: By Lemma 3.7, there are four subcases to consider.
 - (4a) $u = s_{\mathbf{h}}s_{\mathbf{j}\mathbf{i}}^*$ and $v = s_{\mathbf{j}}s_{\mathbf{h}\mathbf{k}}^*$ for some $\mathbf{h}, \mathbf{i}, \mathbf{j}, \mathbf{k}$ in \mathbf{I} : Then $s = s_{\mathbf{h}}s_{\mathbf{h}\mathbf{k}\mathbf{i}}^*$ and $t = s_{\mathbf{j}}s_{\mathbf{j}\mathbf{i}\mathbf{k}}^*$, which both belong to $E(s_{\mathbf{i}\mathbf{k}}^*)$.
 - (4b) $u = s_{\mathbf{h}\mathbf{i}}s_{\mathbf{k}\mathbf{j}}^*$ and $v = s_{\mathbf{k}}s_{\mathbf{h}}^*$ for some $\mathbf{h}, \mathbf{i}, \mathbf{j}, \mathbf{k}$ in \mathbf{I} : Then $s = s_{\mathbf{h}\mathbf{i}}s_{\mathbf{h}\mathbf{j}}^*$ and $t = s_{\mathbf{k}\mathbf{i}}s_{\mathbf{k}\mathbf{j}}^*$. By (3.7), we can take x in $\{\theta, s_{\mathbf{g}}, s_{\mathbf{g}}^* \mid \mathbf{g} \in \mathbf{I}\}$ such that $s_{\mathbf{i}}s_{\mathbf{j}}^*$ belongs to $E(x)$. By the definition of $E(x)$, we have $s_{\mathbf{f}}E(x)s_{\mathbf{f}}^* \subseteq E(x)$ for each \mathbf{f} in \mathbf{I} , and so s and t both belong to $E(x)$.
 - (4c) $u = s_{\mathbf{h}}s_{\mathbf{i}}^*$ and $v = s_{\mathbf{i}\mathbf{j}}s_{\mathbf{h}\mathbf{k}}^*$ for some $\mathbf{h}, \mathbf{i}, \mathbf{j}, \mathbf{k}$ in \mathbf{I} : This is similar to subcase (4b).
 - (4d) $u = s_{\mathbf{h}\mathbf{i}}s_{\mathbf{j}}^*$ and $v = s_{\mathbf{j}\mathbf{k}}s_{\mathbf{h}}^*$ for some $\mathbf{h}, \mathbf{i}, \mathbf{j}, \mathbf{k}$ in \mathbf{I} : Then $s = s_{\mathbf{h}\mathbf{i}\mathbf{k}}s_{\mathbf{h}}^*$ and $t = s_{\mathbf{j}\mathbf{k}\mathbf{i}}s_{\mathbf{j}}^*$, which both belong to $E(s_{\mathbf{i}\mathbf{k}}^*)$.

In each of the above cases, we see that $s \tilde{\lambda} t$, as required.

We conclude that $\tilde{\lambda} = \widehat{\lambda}$, and clauses (i) and (ii) follow from (3.8).

Clause (iii) is a consequence of (i), (ii), and (3.7). □

Combining Proposition 3.10 and Lemma 3.11 yields the following characterization of the bounded traces on $\ell_1(S_2)$ among the elements of $\ell_\infty(S_2)$.

3.12 Proposition *An element τ in $\ell_\infty(S_2)$ is a trace on $\ell_1(S_2)$ if and only if it satisfies both of the following two conditions:*

- (i) $\tau(\theta) = \tau(s_{\mathbf{i}}s_{\mathbf{j}}^*)$ for each $\mathbf{i} = (i_1, \dots, i_m)$ and $\mathbf{j} = (j_1, \dots, j_n)$ in $\mathbf{I} \setminus \{\emptyset\}$ with $i_p \neq j_p$ for some $p \leq \min\{m, n\}$;
- (ii) $\tau(s_{\mathbf{i}}) = \tau(s_{\mathbf{h}\mathbf{j}\mathbf{k}}s_{\mathbf{h}}^*)$ and $\tau(s_{\mathbf{i}}^*) = \tau(s_{\mathbf{h}}s_{\mathbf{h}\mathbf{j}\mathbf{k}}^*)$ for each $\mathbf{h}, \mathbf{i}, \mathbf{j}$, and \mathbf{k} in \mathbf{I} with $\mathbf{i} = \mathbf{k}\mathbf{j}$. \square

By (3.3) and Proposition 3.12, we finally obtain the following characterization of the bounded traces on $\ell_1(S_2^\blacksquare)$ among the elements of $\ell_\infty(S_2^\blacksquare)$.

3.13 Corollary *An element τ in $\ell_\infty(S_2^\blacksquare)$ is a trace on $\ell_1(S_2^\blacksquare)$ if and only if it satisfies both of the following two conditions:*

- (i) $\tau(s_{\mathbf{i}}s_{\mathbf{j}}^*) = 0$ for each $\mathbf{i} = (i_1, \dots, i_m)$ and $\mathbf{j} = (j_1, \dots, j_n)$ in $\mathbf{I} \setminus \{\emptyset\}$ with $i_p \neq j_p$ for some $p \leq \min\{m, n\}$;
- (ii) $\tau(s_{\mathbf{i}}) = \tau(s_{\mathbf{h}\mathbf{j}\mathbf{k}}s_{\mathbf{h}}^*)$ and $\tau(s_{\mathbf{i}}^*) = \tau(s_{\mathbf{h}}s_{\mathbf{h}\mathbf{j}\mathbf{k}}^*)$ for each $\mathbf{h}, \mathbf{i}, \mathbf{j}$, and \mathbf{k} in \mathbf{I} with $\mathbf{i} = \mathbf{k}\mathbf{j}$. \square

We shall next investigate which standard properties of a C^* -algebra that $\ell_1(S_2^\blacksquare)$ possesses. In particular, we shall show that it is $*$ -semisimple.

3.14 Definition An *inverse semigroup* is a semigroup S such that, for each element s in S , there is a unique element t in S with $s = sts$ and $t = tst$.

Let S be an inverse semigroup. Then the map $s \mapsto t$, $S \rightarrow S$, where t is the unique element in S with $s = sts$ and $t = tst$, is an involution on S .

Barnes has proved that, for each inverse semigroup S , the Banach $*$ -algebra $\ell_1(S)$ has a faithful $*$ -representation on a Hilbert space (see [1, Theorem 2.3]). Observe, however, that in the case where S is an inverse semigroup with a zero element θ , Barnes writes $\ell_1(S)$ for the Banach algebra that we denote by $\ell_1^\theta(S)$ (see [1, p. 365]), and so it follows from Barnes's work that $\ell_1(S^\blacksquare)$ is $*$ -semisimple. We note in passing that Wordingham has subsequently improved Barnes's result by showing that the left regular $*$ -representation of $\ell_1(S)$ on $\ell_2(S)$ is faithful whenever S is an inverse semigroup (see [20, Theorem 2] or [16, Theorem 2.1.1]). It is a well-known consequence of Lemma 3.7 that S_2 is an inverse semigroup (*e.g.*, see [16, p. 3]), and so we obtain the following result.

3.15 Proposition *The unital Banach $*$ -algebra $\ell_1(S_2^\blacksquare)$ is $*$ -semisimple. \square*

3.16 Remark It is also possible to give an entirely elementary proof of Proposition 3.15. Indeed, let $(e_m : m \in \mathbb{N})$ denote the canonical orthonormal basis for the Hilbert space $\ell_2(\mathbb{N})$, and set

$$R_i(e_m) := e_{3(m-1)+i} \quad (m \in \mathbb{N}, i = 1, 2).$$

These definitions extend by linearity and continuity to give isometric operators R_1 and R_2 on $\ell_2(\mathbb{N})$. Set $\pi(\delta_{s_i}) := R_i$ for $i = 1, 2$. It is straightforward to see that this definition has a unique extension to a unital $*$ -representation $\pi : \ell_1(S_2^\blacksquare) \rightarrow \mathcal{B}(\ell_2(\mathbb{N}))$, and that π is faithful.

3.17 Theorem *There is a *-semisimple, unital, *-properly infinite Banach *-algebra in standard form which has a non-zero, bounded trace. Specifically, the corner of $\ell_1(S_2^\blacksquare)$ corresponding to the projection $\delta_e - \delta_{s_1 s_1^*}$ has these properties.*

Proof. Set $p := \delta_e - \delta_{s_1 s_1^*}$, and set $\mathcal{A} := \{p \widehat{\star} f \widehat{\star} p \mid f \in \ell_1(S_2^\blacksquare)\}$. Proposition 1.15 and its proof (including the modified version of the proof of Proposition 1.8) show that \mathcal{A} is a *-semisimple, unital, *-properly infinite Banach *-algebra in standard form.

The restriction of a bounded trace to a subalgebra is itself a bounded trace, so to prove that there is a non-zero, bounded trace on \mathcal{A} , it suffices to find a bounded trace on $\ell_1(S_2^\blacksquare)$ taking a non-zero value at an element in \mathcal{A} . Let τ in $\ell_\infty(S_2)$ be defined as in Example 3.8. Then $\tau|_{S_2^\blacksquare}$ is a bounded trace on $\ell_1(S_2^\blacksquare)$, and

$$\langle p \widehat{\star} \delta_{s_2 s_2^*} \widehat{\star} p, \tau|_{S_2^\blacksquare} \rangle = \tau(s_2 s_2^*) = 1 \neq 0,$$

as desired. □

3.18 Proposition *The Banach *-algebra $\ell_1(S_2^\blacksquare)$ is not hermitian.*

Proof. Set $f := \delta_{s_1} + \delta_{s_2}$ in $\ell_1(S_2^\blacksquare)$. Then $f^* \widehat{\star} f = 2\delta_e$, and so

$$\sigma(f^* \widehat{\star} f) = \{2\} \quad \text{and} \quad \nu(f^* \widehat{\star} f) = 2.$$

On the other hand, $\|f^n\|_1 = 2^n$ for each n in \mathbb{N} because no cancellation occurs among the 2^n terms of the expression

$$f^n = \sum_{i_1, \dots, i_n=1}^2 \delta_{s_{i_1} s_{i_2} \dots s_{i_n}}.$$

This implies that $\nu(f) = 2$ by the spectral radius formula (*e.g.*, see [7, Theorem 2.3.8(iii)]).

In conclusion, we have

$$\nu(f^* \widehat{\star} f) = 2 < 4 = \nu(f)^2,$$

and so $\ell_1(S_2^\blacksquare)$ is not hermitian by [7, Proposition 3.1.8]. □

3.19 Remark In fact, the proof of [7, Proposition 3.1.8] provides an explicit example of a self-adjoint element k in $\ell_1(S_2^\blacksquare)$ whose spectrum is not contained in \mathbb{R} .

To see this, we first show that the spectrum of the element $f := \delta_{s_1} + \delta_{s_2}$ in $\ell_1(S_2^\blacksquare)$ is given by

$$\sigma(f) = \{\zeta \in \mathbb{C} \mid |\zeta| \leq 2\}. \tag{3.9}$$

The inclusion \subseteq is clear because $\|f\|_1 = 2$.

Conversely, suppose that ζ belongs to $\mathbb{C} \setminus \sigma(f)$, and take $g = \sum_{s \in S_2^\blacksquare} \alpha_s \delta_s$ in $\ell_1(S_2^\blacksquare)$ such that $(f - \zeta \delta_e) \widehat{\star} g = \delta_e$, that is,

$$\sum_{s \in S_2^\blacksquare} \alpha_s \delta_{s_1 s} + \sum_{s \in S_2^\blacksquare} \alpha_s \delta_{s_2 s} - \sum_{s \in S_2^\blacksquare} \zeta \alpha_s \delta_s = \delta_e.$$

Identifying coefficients gives

$$-\zeta\alpha_e = 1, \quad \zeta\alpha_{s_1^*} = 0 \quad (\mathbf{i} \in \mathbf{I} \setminus \{\emptyset\}), \quad \text{and } \alpha_s = \zeta\alpha_{s_i s} \quad (s \in S_2^\blacksquare, i = 1, 2).$$

It follows that $\zeta \neq 0$, $\alpha_e = -1/\zeta$, and, inductively,

$$\begin{aligned} \alpha_{s_{i_1} s_{i_2} \cdots s_{i_n}} &= -\frac{1}{\zeta^{n+1}} \quad (n \in \mathbb{N}, i_1, \dots, i_n \in \{1, 2\}), \\ \alpha_{s_i s_j^*} &= 0 \quad (\mathbf{i} \in \mathbf{I}, \mathbf{j} \in \mathbf{I} \setminus \{\emptyset\}). \end{aligned}$$

Thus, we have

$$\infty > \|g\|_1 = \frac{1}{|\zeta|} + \sum_{n=1}^{\infty} \left(\sum_{i_1, \dots, i_n=1}^2 \frac{1}{|\zeta|^{n+1}} \right) = \sum_{n=0}^{\infty} \frac{2^n}{|\zeta|^{n+1}},$$

from which we conclude that $|\zeta| > 2$. This completes the proof of (3.9).

Set $k := i(f^* - f)$ in $\ell_1(S_2^\blacksquare)$. Then k is self-adjoint and satisfies

$$(f^* + 2\delta_e) \widehat{\star} (f - 2\delta_e) = 2i(k + i\delta_e).$$

Assume towards a contradiction that $\sigma(k)$ is contained in \mathbb{R} . Then $k + i\delta_e$ is invertible, and so $f - 2\delta_e$ is left invertible. By [7, Theorem 2.3.21(ii)], this implies that 2 is not in $\partial\sigma(f)$, contradicting (3.9). Hence, we conclude that k has non-real spectrum; specifically, $-i$ belongs to $\sigma(k)$. \square

We shall now show that the corner of $\ell_1(S_2^\blacksquare)$ considered in Theorem 3.17 also fails to be hermitian. By Proposition 1.15(iii), this result improves Proposition 3.18. The strategy of the proof is exactly the same as before, but the details are somewhat more complicated.

3.20 Proposition *The corner of $\ell_1(S_2^\blacksquare)$ corresponding to the projection $\delta_e - \delta_{s_1 s_1^*}$ is not hermitian.*

Proof. Set $p := \delta_e - \delta_{s_1 s_1^*}$, and set $\mathcal{A} := \{p \widehat{\star} f \widehat{\star} p \mid f \in \ell_1(S_2^\blacksquare)\}$. By Lemma 1.9, the spectral radius of an element a in \mathcal{A} is independent of whether it is calculated in $\ell_1(S_2^\blacksquare)$ or in \mathcal{A} , and so we shall denote it just by $\nu(a)$ (without any subscript).

Set

$$g := \delta_{s_2} \widehat{\star} p + \delta_{s_2 s_1 s_2^*} + p - \delta_{s_2 s_2^*} \in \mathcal{A}.$$

Then $g^* \widehat{\star} g = 2p$, and so $\nu(g^* \widehat{\star} g) = 2$. On the other hand, an inductive argument shows that, for each integer $n \geq 2$, the n^{th} power of g is given by

$$g^n = \delta_{s_2} \widehat{\star} (\delta_{s_1} + \delta_{s_2})^{n-1} \widehat{\star} (p + \delta_{s_1 s_2^*}) + \left(\delta_e + \delta_{s_2} \widehat{\star} \left(\sum_{m=0}^{n-2} (\delta_{s_1} + \delta_{s_2})^m \right) \right) \widehat{\star} (p - \delta_{s_2 s_2^*}).$$

By Lemma 3.7(ii), the terms of this expression have pairwise disjoint supports. This implies that $\|g^n\|_1 = 3 \cdot 2^n$, and so $\nu(g) = 2$. In conclusion, we have $\nu(g^* \widehat{\star} g) = 2 < 4 = \nu(g)^2$, and the result follows from [7, Proposition 3.1.8]. \square

We conclude this section with another example of a standard property of C^* -algebras that fails for $\ell_1(S_2^\blacksquare)$.

3.21 Definition A Banach algebra \mathcal{A} is *Arens regular* if, for each pair $((a_m)_{m=1}^\infty, (b_n)_{n=1}^\infty)$ of bounded sequences in \mathcal{A} and each bounded linear functional $\lambda: \mathcal{A} \rightarrow \mathbb{C}$, we have

$$\lim_{m \rightarrow \infty} \left(\lim_{n \rightarrow \infty} \langle a_m b_n, \lambda \rangle \right) = \lim_{n \rightarrow \infty} \left(\lim_{m \rightarrow \infty} \langle a_m b_n, \lambda \rangle \right)$$

whenever both the iterated limits exist.

A number of equivalent formulations of the notion of Arens regularity is given in [7, Theorem 2.6.17]. It is well-known that every C^* -algebra is Arens regular (*e.g.*, see [7, Corollary 3.2.37]). Using a technique introduced by Young in [21], we shall now show that this is not the case for $\ell_1(S_2^\blacksquare)$.

3.22 Proposition *The Banach $*$ -algebra $\ell_1(S_2^\blacksquare)$ is not Arens regular.*

Proof. Define λ in $\ell_\infty(S_2^\blacksquare)$ by

$$\lambda(s) := \begin{cases} 1 & \text{if } s = s_1^m (s_1^*)^n \text{ for some } m, n \in \mathbb{N} \text{ with } m \leq n \\ 0 & \text{otherwise} \end{cases} \quad (s \in S_2^\blacksquare).$$

The sequences $(\delta_{s_1^m})_{m=1}^\infty$ and $(\delta_{(s_1^*)^n})_{n=1}^\infty$ in $\ell_1(S_2^\blacksquare)$ are bounded and satisfy:

$$\langle \delta_{s_1^m} \widehat{\star} \delta_{(s_1^*)^n}, \lambda \rangle = \lambda(s_1^m (s_1^*)^n) = \begin{cases} 1 & \text{for } m \leq n, \\ 0 & \text{for } m > n. \end{cases}$$

It follows that

$$\lim_{m \rightarrow \infty} \left(\lim_{n \rightarrow \infty} \langle \delta_{s_1^m} \widehat{\star} \delta_{(s_1^*)^n}, \lambda \rangle \right) = 1 \quad \text{and} \quad \lim_{n \rightarrow \infty} \left(\lim_{m \rightarrow \infty} \langle \delta_{s_1^m} \widehat{\star} \delta_{(s_1^*)^n}, \lambda \rangle \right) = 0,$$

and so $\ell_1(S_2^\blacksquare)$ is not Arens regular. □

4 A unital $*$ -algebra which is $*$ -finite and properly infinite

A unital, finite $*$ -algebra is automatically $*$ -finite, and a unital, $*$ -properly infinite $*$ -algebra is automatically properly infinite. In this section we shall show that neither of these implications can be reversed by constructing a unital $*$ -algebra which is $*$ -finite and properly infinite.

4.1 Algebraic semigroup algebras For a semigroup S , we denote by $c_{00}(S)$ the set of finitely supported functions from S to \mathbb{C} . This is a dense subalgebra of $\ell_1(S)$, termed the *algebraic semigroup algebra* of S . Clearly, $c_{00}(S) = \ell_1(S)$ if and only if S has finite cardinality. In the case where S has a zero element, $c_{00}(S^\blacksquare)$ is a dense subalgebra of $\ell_1(S^\blacksquare)$.

Suppose that S has an involution. Then $c_{00}(S)$ is closed under the involution (3.4), and is thus a dense sub- $*$ -algebra of $\ell_1(S)$. Similarly, in the case where S has a zero element, $c_{00}(S^\blacksquare)$ is a dense sub- $*$ -algebra of $\ell_1(S^\blacksquare)$.

4.2 Construction Let S be a semigroup with a neutral element e and a zero element θ . The *free product* of two copies of S with amalgamation of the neutral elements and of the zero elements is given by

$$S * S := \{e', \theta'\} \cup \{(w_1, i_1)(w_2, i_2) \cdots (w_n, i_n) \mid n \in \mathbb{N}, w_1, \dots, w_n \in S \setminus \{e, \theta\}, \text{ and } i_1, \dots, i_n \in \{1, 2\} \text{ with } i_j \neq i_{j+1} \text{ for each } j \in \{1, \dots, n-1\}\}.$$

We turn $S * S$ into a semigroup with neutral element e' and zero element θ' by defining an operation \circ as follows:

$$\theta' \circ w = \theta' = w \circ \theta', \quad e' \circ w = w = w \circ e' \quad (w \in S * S),$$

and, recursively, for $v = (v_1, i_1) \cdots (v_m, i_m)$ and $w = (w_1, j_1) \cdots (w_n, j_n)$ in $S * S \setminus \{e', \theta'\}$,

$$v \circ w := \begin{cases} \theta' & \text{for } i_m = j_1 \text{ and } v_m w_1 = \theta; \\ e' & \text{for } i_m = j_1, v_m w_1 = e, \\ & \text{and } m = n = 1; \\ (v_1, i_1) \cdots (v_{m-1}, i_{m-1}) \circ (w_2, j_2) \cdots (w_n, j_n) & \text{for } i_m = j_1, v_m w_1 = e, \\ & \text{and } (m, n) \neq (1, 1); \\ (v_1, i_1) \cdots (v_{m-1}, i_{m-1})(v_m w_1, j_1)(w_2, j_2) \cdots (w_n, j_n) & \text{for } i_m = j_1 \text{ and} \\ & v_m w_1 \in S \setminus \{e, \theta\}; \\ (v_1, i_1) \cdots (v_m, i_m)(w_1, j_1) \cdots (w_n, j_n) & \text{for } i_m \neq j_1. \end{cases}$$

(Note: in the third and fourth case, $(v_1, i_1) \cdots (v_{m-1}, i_{m-1})$ should be ignored if $m = 1$, and $(w_2, j_2) \cdots (w_n, j_n)$ should be ignored if $n = 1$.)

Suppose that S has an involution. Then we define an involution on $S * S$ by $(e')^* := e'$, $(\theta')^* := \theta'$, and

$$((w_1, i_1)(w_2, i_2) \cdots (w_n, i_n))^* := (w_n^*, i_n^*)(w_{n-1}^*, i_{n-1}^*) \cdots (w_1^*, i_1^*)$$

for each $(w_1, i_1)(w_2, i_2) \cdots (w_n, i_n)$ in $(S * S) \setminus \{e', \theta'\}$, where

$$i^* := \begin{cases} 2 & \text{for } i = 1 \\ 1 & \text{for } i = 2 \end{cases} \quad (i = 1, 2).$$

The *length* of an element w in $S * S$ is given by

$$\text{len}(w) := \begin{cases} -\infty & \text{for } w = \theta' \\ 0 & \text{for } w = e' \\ n & \text{for } w = (w_1, i_1) \cdots (w_n, i_n) \in (S * S) \setminus \{e', \theta'\}. \end{cases}$$

Clearly, for each v and w in $S * S$, we have

$$\left. \begin{aligned} \text{len}(v \circ w) &\leq \text{len}(v) + \text{len}(w) \\ \text{len}(w^*) &= \text{len}(w) \\ \text{len}(w^* \circ w) &= 2\text{len}(w). \end{aligned} \right\} \quad (4.1)$$

As is customary, we shall henceforth omit the symbol \circ and denote the semigroup operation in $S * S$ simply by juxtaposition of elements. Moreover, we shall drop the primes from e' and θ' and write e for the neutral element and θ for the zero element in $S * S$ (as well as in S).

4.3 Lemma *The only projections in $c_{00}((S_2 * S_2)^\blacksquare)$ are 0 and δ_e .*

Proof. For f in $c_{00}((S_2 * S_2)^\blacksquare) \setminus \{0, \delta_e\}$, take $w = (w_1, i_1) \cdots (w_n, i_n)$ in $(S_2 * S_2)^\blacksquare \setminus \{e\}$ of maximal length such that $f(w) \neq 0$. Then $f(w^*w) = 0$ because $\text{len}(w^*w) = 2n > n$.

On the other hand, by definition, we have

$$(f^* \widehat{\star} f)(w^*w) = \sum_{\substack{u, v \in S_2 * S_2 \\ uv = w^*w}} \overline{f(u^*)} f(v). \quad (4.2)$$

Suppose that u and v in $S_2 * S_2$ satisfy $uv = w^*w$. Then $\text{len}(u) + \text{len}(v) \geq 2n$ by (4.1). Since $f(t) = 0$ for each t in $(S_2 * S_2)^\blacksquare$ with $\text{len}(t) > n$, the only non-zero contributions to the sum on the right-hand side of (4.2) arise for $\text{len}(u) = \text{len}(v) = n$. But the only u and v of length n in $S_2 * S_2$ with $uv = w^*w$ are $u = w^*$ and $v = w$. It follows that

$$(f^* \widehat{\star} f)(w^*w) = \overline{f((w^*)^*)} f(w) = |f(w)|^2 \neq 0.$$

In conclusion, we have $f \neq f^* \widehat{\star} f$, and so f is not a projection. \square

4.4 Proposition *The unital $*$ -algebra $c_{00}((S_2 * S_2)^\blacksquare)$ is $*$ -finite and properly infinite.*

Proof. Lemma 4.3 shows that $c_{00}((S_2 * S_2)^\blacksquare)$ is $*$ -finite.

On the other hand, set $a_i := \delta_{(s_i, 1)}$ and $b_i := \delta_{(s_i^*, 1)}$ in $c_{00}((S_2 * S_2)^\blacksquare)$ for $i = 1, 2$. Then (1.1) is satisfied, and so we conclude that $c_{00}((S_2 * S_2)^\blacksquare)$ is properly infinite. \square

4.5 Remark The unital Banach $*$ -algebra $\ell_1((S_2 * S_2)^\blacksquare)$ is properly infinite because it contains $c_{00}((S_2 * S_2)^\blacksquare)$, which is properly infinite, as a unital subalgebra. We have not been able to determine whether or not $\ell_1((S_2 * S_2)^\blacksquare)$ is $*$ -properly infinite. If $\ell_1((S_2 * S_2)^\blacksquare)$ fails to be $*$ -properly infinite, then it would be of interest to determine its traces, *cf.* Question 1.11(iv).

Acknowledgements

We are grateful to John Duncan (Arkansas) for sharing with us his insights into semigroup Banach algebras and for drawing our attention to reference [19], and to Mikael Rørdam (Odense) for his advice on properly infinite C^* -algebras.

References

- [1] B. A. Barnes, Representations of the ℓ_1 -algebra of an inverse semigroup, *Trans. American Math. Soc.*, **218** (1976), 361–396.
- [2] S. Bowling and J. Duncan, First order cohomology of Banach semigroup algebras, *Semigroup Forum*, **56** (1998), 130–145.
- [3] A. H. Clifford and G. B. Preston, *The algebraic theory of semigroups*, Math. Surveys, **7**, American Math. Soc., Providence, 1961.
- [4] J. Cuntz, Simple C^* -algebras generated by isometries, *Commun. math. Phys.*, **57** (1977), 173–185.
- [5] J. Cuntz, Murray–von Neumann equivalence of projections in infinite simple C^* -algebras, *Rev. Roum. Math. Pures et Appl.*, **23** (1978), 1011–1014.
- [6] J. Cuntz, K -theory for certain C^* -algebras, *Annals of Math.*, **113** (1981), 181–197.
- [7] H. G. Dales, *Banach algebras and automatic continuity*, London Math. Soc. Monographs, **24**, Clarendon Press, Oxford, 2000.
- [8] J. Duncan and J. H. Williamson, Spectra of elements in the measure algebra of a free group, *Proc. Royal Irish Acad.*, **82A** (1982), 109–120.
- [9] T. Fack, Finite sums of commutators in C^* -algebras, *Ann. Inst. Fourier, Grenoble*, **32**, 1 (1982), 129–137.
- [10] B. E. Johnson, Symmetric amenability and the nonexistence of Lie and Jordan derivations, *Math. Proc. Camb. Phil. Soc.*, **120** (1996), 455–473.
- [11] R. V. Kadison and J. R. Ringrose, *Fundamentals of the theory of operator algebras*, Vol. I–II, Academic Press, San Diego, 1983–1986.
- [12] N. J. Laustsen, Commutators of operators on Banach spaces, *J. Operator Theory*, to appear.
- [13] N. J. Laustsen, On ring-theoretic (in)finiteness of Banach algebras of operators on Banach spaces, *Glasgow J. Math.*, to appear.
- [14] M. Mathieu and A. R. Villena, The structure of Lie derivations on C^* -algebras, preprint.
- [15] T. W. Palmer, *Banach algebras and the general theory of $*$ -algebras*, Vol. II, Cambridge University Press, 2001.
- [16] A. L. T. Paterson, *Groupoids, inverse semigroups, and their operator algebras*, Progress Math., **170**, Birkhäuser, Boston, 1999.

- [17] C. Pop, Finite sums of commutators, *Proc. American Math. Soc.*, to appear.
- [18] J. N. Renault, *A groupoid approach to C^* -algebras*, Lecture Notes Math., **793**, Springer-Verlag, New York, 1980.
- [19] J. R. Wordingham, *Topics in semigroup algebras*, PhD Thesis, University of Stirling, 1982.
- [20] J. R. Wordingham, The left regular $*$ -representation of an inverse semigroup, *Proc. American Math. Soc.*, **86** (1982), 55–58.
- [21] N. J. Young, The irregularity of multiplication in group algebras, *Quarterly J. Math. Oxford* (2), **24** (1973), 59–62.

H. G. Dales and C. J. Read
Department of Pure Mathematics
University of Leeds
Leeds LS2 9JT
England

pmt6hgd@amsta.leeds.ac.uk
read@amsta.leeds.ac.uk

N. J. Laustsen
Department of Mathematics
University of Copenhagen
Universitetsparken 5
DK-2100 Copenhagen Ø
Denmark

laustsen@math.ku.dk